

National Enteric Disease Surveillance: The *Listeria* Initiative



Surveillance System Overview: The *Listeria* Initiative

Listeria monocytogenes is estimated to cause about 1,600 illnesses each year in the United States with more than 1,500 related hospitalizations and 260 related deaths (1). *Listeria* infections (listeriosis) are nationally notifiable. Nearly all cases in persons who are not infants result from eating food contaminated with *L. monocytogenes*; newborn infants can develop listeriosis if their mothers ate contaminated food during pregnancy.

The *Listeria* Initiative is an enhanced surveillance system that collects reports of laboratory-confirmed cases of human listeriosis in the United States. Demographic, clinical, laboratory, and epidemiologic data are collected using a standardized, extended questionnaire. The *Listeria* Initiative was piloted in the Foodborne Diseases Active Surveillance Network (FoodNet) in 2004 and implemented nationwide in 2005. The number of states participating increased to 47 states and the District of Columbia by 2014. The proportion of all listeriosis cases reported to the *Listeria* Initiative continues to increase.

A main objective of the *Listeria* Initiative is to aid in the investigation of listeriosis clusters and outbreaks by decreasing the time from outbreak detection to public health intervention. Patient interviews are conducted as cases are reported, rather than after clusters are identified, to minimize the effect of recall bias on food consumption history. In addition, clinical, food, and environmental isolates of *L. monocytogenes* are subtyped. Pulsed-field gel electrophoresis (PFGE, a type of DNA fingerprinting) subtyping results are submitted to PulseNet, the National Molecular Subtyping Network for Foodborne Disease Surveillance, to identify clusters of possibly related cases. Since September 2013, state laboratories, CDC, the Food and Drug Administration (FDA), and the US Department of Agriculture's Food Safety and Inspection Service also perform whole genome sequencing on clinical, food, and environmental *L. monocytogenes* isolates. This method provides high-resolution genetic information. Whole genome sequencing can identify related illnesses that are not evident using PFGE alone. Genetically related bacteria may have different PFGE patterns, and distantly related bacteria can have the same pattern. When clusters are identified, *Listeria* Initiative data are used to rapidly conduct epidemiological analyses. The food consumption histories of patients with cluster-associated illnesses are compared with those of patients with sporadic illnesses to identify foods possibly associated with the cluster. Without the *Listeria* Initiative database, appropriate comparison data ("controls") for listeriosis investigations would be difficult to obtain through traditional methods; the population at risk for invasive listeriosis—older adults, immunocompromised persons, and pregnant women—is a small segment of the general population.

Local, state, and territorial public health professionals are encouraged to complete the *Listeria* Initiative questionnaire for all cases of laboratory-confirmed listeriosis. English and Spanish versions of the questionnaire are available at <http://www.cdc.gov/listeria/surveillance.html>. All *Listeria* isolates should be forwarded promptly to state or national laboratories for subtyping.

Overview of *Listeria*

The genus *Listeria* contains seven species (*monocytogenes*, *ivanovii*, *seeligeri*, *innocua*, *welshimeri*, *martii*, and *grayi*), two of which are pathogenic. *L. monocytogenes* is pathogenic to humans and animals; *L. ivanovii* (previously *L. monocytogenes* serotype 5) primarily infects animals and very rarely causes disease in humans.

Serotyping differentiates isolates of *Listeria* below the species level. Serotypes are designated based on the immunoreactivity of two cell surface structures, the O and H antigens. Twelve serotypes of *L. monocytogenes* (1/2a, 1/2b, 1/2c, 3a, 3b, 3c, 4a, 4b, 4c, 4d, 4e, and 7) are recognized, three of which (1/2a, 1/2b, and 4b) cause most (95%) human illness; serotype 4b is most commonly associated with outbreaks.

References

1. Scallan, E, Hoekstra RM, Angulo FJ, et al. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis* 2011; 17:7–15.

Suggested Readings

Cartwright, EJ, Jackson KA, Johnson SD, et al. Listeriosis outbreaks and associated food vehicles, United States, 1998–2008. *Emerg Infect Dis*. 2013 Jan;19:1–9.

CDC. Vital Signs: *Listeria* illnesses, deaths, and outbreaks—United States, 2009–2011. *MMWR Morb Mortal Wkly Rep* 2013; 62:448–452.

de Noordhout, CM, Devleeschauwer B, Angulo FJ, et al. The global burden of listeriosis: a systematic review and meta-analysis. *Lancet* 2014; 14:1027–8.

Gaul, L, Faraq N, Shim T, et al. Hospital-acquired listeriosis outbreak caused by contaminated diced celery—Texas, 2010. *Clin Infect Dis* 2013;56:20–6.

Heiman, KE, Garalde VB, Gronostaj M, et al. Multistate outbreak of listeriosis caused by imported cheese and evidence of cross-contamination of other cheeses, USA, 2012. *Epidemiol Infect* 2015; [Epub ahead of print].

Jackson, K, Iwamoto M, Swerdlow D. Pregnancy-associated listeriosis. *Epidemiol Infect* 2010; 138:1503–1509.

McCollum, J, Cronquist A, Silk, B, et al. Multistate outbreak of listeriosis associated with cantaloupe. *N Eng J Med* 2013; 369:944–953.

Silk, B, Date K, Jackson K, et al. Invasive listeriosis in the Foodborne Diseases Active Surveillance Network (FoodNet), 2004–2009: Further targeted prevention needed for higher-risk groups. *Clin Infect Dis* 2012; 54:S396–S404.

Swaminathan, B, Gerner-Smidt P. The epidemiology of human listeriosis. *Microb Infect* 2007; 9:1236–1243.

Voetsch, A, Angulo F, Jones T, et al. Reduction in the incidence of invasive listeriosis in foodborne diseases active surveillance network sites, 1996–2006. *Clin Infect Dis* 2007;44:513–520.