



November 2, 2001 / Vol. 50 / No. SS-4



CDC
Surveillance
Summaries

Youth Tobacco Surveillance — United States, 2000

U.S. DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Disease Control and Prevention (CDC)
Atlanta, GA 30333



The MMWR series of publications is published by the Epidemiology Program Office, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

General:Centers for Disease Control and Prevention.*CDC Surveillance Summaries*, November 2, 2001. MMWR 2001;50(No. SS-4).
Specific:[Author(s)]. [Title of particular article.] In: *CDC Surveillance Summaries*, November 2, 2001. MMWR 2001;50(No. SS-4):[inclusive page numbers].

Centers for Disease Control and Prevention Jeffrey P. Koplan, M.D., M.P.H.
Director

This report was produced as an *MMWR* serial publication in
Epidemiology Program Office Stephen B. Thacker, M.D, M.Sc.
Director

Division of Public Health Surveillance
and Informatics Daniel M. Sosin, M.D., M.P.H.
Director
Associate Editor, CDC Surveillance Summaries

Office of Scientific and Health Communications John W. Ward, M.D.
Director
Editor, MMWR Series

CDC Surveillance Summaries Suzanne M. Hewitt, M.P.A.
Managing Editor

Patricia A. McGee
Project Editor

Lynda G. Cupell
Visual Information Specialist

Michele D. Renshaw
Erica R. Shaver
Information Technology Specialists

All *MMWR* references are available on the Internet at <<http://www.cdc.gov/mmwr/>>. Use the search function to find specific articles.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

Contents

Reports Published in <i>CDC Surveillance Summaries</i> Since January 1, 1991	ii
Youth Tobacco Surveillance — United States, 2000	1
Introduction	2
Methods	3
Sampling	3
Data Collection	4
Results	5
Prevalence of Use	5
Knowledge and Attitudes	17
Access and Enforcement	20
Age and Access	22
Media and Advertising	24
Cessation	30
Secondhand Smoke	31
School	36
Discussion	38
Linking YTS Data to Program Components	39
Limitations	42
Conclusion	42
State Implementation of YTS	42
References	44
Tables	46
 State and Territorial Epidemiologists and Laboratory Directors	 Inside Back Cover

Reports Published in *CDC Surveillance Summaries* Since January 1, 1991

Subject	Responsible CIO/Agency*	Most Recent Report
Abortion	NCCDPHP	2000; Vol. 49, No. SS-11
Aging		
Health Risks	NCCDPHP	1999; Vol. 48, No. SS-8
Health-Care Services	NCCDPHP/NIP	1999; Vol. 48, No. SS-8
Health-Related Quality of Life	NCEH/NCCDPHP	1999; Vol. 48, No. SS-8
Injuries and Violence	NCIPC/NCCDPHP	1999; Vol. 48, No. SS-8
Morbidity and Mortality	NCHS/NCCDPHP	1999; Vol. 48, No. SS-8
AIDS/HIV		
AIDS-Defining Opportunistic Illnesses	NCHSTP/NCID	1999; Vol. 48, No. SS-2
Asthma	NCEH	1998; Vol. 47, No. SS-1
Behavioral Risk Factors		
State-Specific Prevalence of Selected Health Behaviors, by Race and Ethnicity	NCCDPHP	2000; Vol. 49, No. SS-2
State- and Sex-Specific Prevalence of Selected Characteristics	NCCDPHP	2000; Vol. 49, No. SS-6
Birth Defects		
Birth Defects Monitoring Program (see also Malformations)	NCEH	1993; Vol. 42, No. SS-1
Breast and Cervical Cancer	NCCDPHP	1999; Vol. 48, No. SS-6
Cardiovascular Disease	EPO/NCCDPHP	1998; Vol. 47, No. SS-5
Chancroid	NCPS	1992; Vol. 41, No. SS-3
Chlamydia	NCPS	1993; Vol. 42, No. SS-3
Cholera	NCID	1992; Vol. 41, No. SS-1
Chronic Fatigue Syndrome	NCID	1997; Vol. 46, No. SS-2
Contraception Practices	NCCDPHP	1992; Vol. 41, No. SS-4
Cytomegalovirus Disease, Congenital	NCID	1992; Vol. 41, No. SS-2
Dengue	NCID	1994; Vol. 43, No. SS-2
Developmental Disabilities	NCEH	1996; Vol. 45, No. SS-2
Diabetes Mellitus	NCCDPHP	1993; Vol. 42, No. SS-2
Dracunculiasis	NCID	1992; Vol. 41, No. SS-1
Ectopic Pregnancy	NCCDPHP	1993; Vol. 42, No. SS-6
Elderly, Hospitalizations Among	NCCDPHP	1991; Vol. 40, No. SS-1
<i>Escherichia coli</i> O157	NCID	1991; Vol. 40, No. SS-1
Evacuation Camps	EPO	1992; Vol. 41, No. SS-4
Family Planning Services at Title X Clinics	NCCDPHP	1995; Vol. 44, No. SS-2
Firearm-Related Injuries	NCIPC	2001; Vol. 50, No. SS-2
Food Safety	NCID	1998; Vol. 47, No. SS-4
Foodborne-Disease Outbreaks	NCID	2000; Vol. 49, No. SS-1
Giardiasis	NCID	2000; Vol. 49, No. SS-7
Gonorrhea and Syphilis, Teenagers	NCPS	1993; Vol. 42, No. SS-3
Hazardous Substances Emergency Events	ATSDR	1994; Vol. 43, No. SS-2

***Abbreviations**

ATSDR	Agency for Toxic Substances and Disease Registry
CIO	Centers/Institute/Offices
EPO	Epidemiology Program Office
IHPO	International Health Program Office
NCCDPHP	National Center for Chronic Disease Prevention and Health Promotion
NCEH	National Center for Environmental Health
NCEHIC	National Center for Environmental Health and Injury Control
NCHSTP	National Center for HIV, STD, and TB Prevention
NCID	National Center for Infectious Diseases
NCIPC	National Center for Injury Prevention and Control
NCPS	National Center for Prevention Services
NIOSH	National Institute for Occupational Safety and Health
NIP	National Immunization Program

Reports Published in *CDC Surveillance Summaries* Since January 1, 1991 — Continued

Subject	Responsible CIO/Agency*	Most Recent Report
Health Surveillance Systems	IHPO	1992; Vol. 41, No. SS-4
Homicide	NCEHIC	1992; Vol. 41, No. SS-3
Hysterectomy	NCCDPHP	1997; Vol. 46, No. SS-4
Influenza	NCID	2000; Vol. 49, No. SS-3
Injury		
Head and Neck	NCIPC	1993; Vol. 42, No. SS-5
In Developing Countries	NCEHIC	1992; Vol. 41, No. SS-1
Lyme Disease	NCID	2000; Vol. 49, No. SS-3
Malaria	NCID	2001; Vol. 50, No. SS-1
Measles	NCPS	1992; Vol. 41, No. SS-6
Meningococcal Disease	NCID	1993; Vol. 42, No. SS-2
Mumps	NIP	1995; Vol. 44, No. SS-3
<i>Neisseria gonorrhoeae</i> , Antimicrobial Resistance in	NCPS	1993; Vol. 42, No. SS-3
Neural Tube Defects	NCEH	1995; Vol. 44, No. SS-4
Occupational Injuries/Disease		
Asthma	NIOSH	1999; Vol. 48, No. SS-3
Silicosis	NIOSH	1997; Vol. 46, No. SS-1
Parasites, Intestinal	NCID	1991; Vol. 40, No. SS-4
Pediatric Nutrition	NCCDPHP	1992; Vol. 41, No. SS-7
Pertussis	NCPS	1992; Vol. 41, No. SS-8
Poliomyelitis	NCPS	1992; Vol. 41, No. SS-1
Postneonatal Mortality	NCCDPHP	1998; Vol. 47, No. SS-2
Pregnancy		
Pregnancy Nutrition	NCCDPHP	1992; Vol. 41, No. SS-7
Pregnancy-Related Mortality	NCCDPHP	1997; Vol. 46, No. SS-4
Pregnancy Risk Assessment Monitoring System (PRAMS)	NCCDPHP	1999; Vol. 48, No. SS-5
Pregnancy, Teenage	NCCDPHP	1993; Vol. 42, No. SS-6
Respiratory Disease	NCEHIC	1992; Vol. 41, No. SS-4
Rotavirus	NCID	1992; Vol. 41, No. SS-3
School Health Education Profiles	NCCDPHP	2000; Vol. 49, No. SS-8
Sexually Transmitted Diseases in Italy	NCPS	1992; Vol. 41, No. SS-1
Smoking		
Smoking-Attributable Mortality	NCCDPHP	1994; Vol. 43, No. SS-1
Tobacco-Control Laws, State	NCCDPHP	1999; Vol. 48, No. SS-3
Tobacco-Use Behaviors	NCCDPHP	1994; Vol. 43, No. SS-3
Youth Tobacco Surveillance	NCCDPHP	2001; Vol. 50, No. SS-4
Spina Bifida	NCEH	1996; Vol. 45, No. SS-2
Streptococcal Disease (Group B)	NCID	1992; Vol. 41, No. SS-6
Syphilis, Congenital	NCPS	1993; Vol. 42, No. SS-6
Syphilis, Primary and Secondary	NCPS	1993; Vol. 42, No. SS-3
Tetanus	NIP	1998; Vol. 47, No. SS-2
Trichinosis	NCID	1991; Vol. 40, No. SS-3
Tuberculosis	NCPS	1991; Vol. 40, No. SS-3
Vaccination Coverage		
Among Children Enrolled in Head Start Programs or Day Care Facilities or Entering School	NIP	2000; Vol. 49, No. SS-9
Influenza, Pneumococcal, and Tetanus Toxoid Vaccination (Among Adults)	NIP	2000; Vol. 49, No. SS-9
National, State, and Urban Areas (Among Children Aged 19–35 Months)	NIP	2000; Vol. 49, No. SS-9
Waterborne-Disease Outbreaks	NCID	2000; Vol. 49, No. SS-4
Years of Potential Life Lost	EPO	1992; Vol. 41, No. SS-6
Youth Risk Behaviors	NCCDPHP	2000; Vol. 49, No. SS-5
College Students	NCCDPHP	1997; Vol. 46, No. SS-6
National Alternative High Schools	NCCDPHP	1999; Vol. 48, No. SS-7

Youth Tobacco Surveillance — United States, 2000

*Office on Smoking and Health
National Center for Chronic Disease Prevention and Health Promotion
American Legacy Foundation, Washington, D.C.
CDC Foundation, Atlanta, Georgia
Macro International, Calverton, Maryland
State Youth Tobacco Survey Coordinators*

Abstract

Problem/Condition: Tobacco use is the single leading preventable cause of death in the United States, accounting for approximately 430,000 deaths each year. The prevalence of cigarette smoking nationwide among high school students increased during the 1990s, peaking during 1996–1997, then began a gradual decline. Approximately 80% of tobacco users initiate use before age 18 years. If the trend in early initiation of cigarette smoking continues, approximately 5 million children aged <18 years who are living today will die prematurely because they began to smoke cigarettes during adolescence. The economic costs associated with tobacco use ranges from \$53 billion to \$73 billion per year in medical expenses and \$47 billion in lost productivity. Because of these health and economic consequences, CDC has recommended that states establish and maintain comprehensive tobacco-control programs to reduce tobacco use among youth.

Reporting Period: January 2000 through December 2000.

Description of the System: To assist states in developing and maintaining their state-based comprehensive tobacco prevention and control programs, CDC developed the Youth Tobacco Surveillance and Evaluation System, which includes international, national, and state school-based surveys of middle school and high school students. Two components of this system are discussed in this report — the National Youth Tobacco Survey and the state Youth Tobacco Surveys. The national survey is representative of students in the 50 states and the District of Columbia; 35,828 students in 324 schools completed questionnaires in the spring of 2000. Twenty-nine state surveys were conducted in the spring and fall of 2000; state sample sizes ranged from 583 to 33,586 students. This report summarizes data from the 2000 national survey and state surveys.

Results and Interpretation: Findings from the National Youth Tobacco Survey indicate that current tobacco use ranges from 15.1% among middle school students to 34.5% among high school students. Cigarette smoking is the most prevalent form of tobacco use, followed by cigar smoking and smokeless tobacco use. Approximately one half of current cigarette smokers in middle school and high school report that they usually smoke Marlboro® cigarettes. Black students are more likely to smoke Newport® cigarettes than any other brand. More than one half of current cigarette smokers in middle school and high school report that they want to stop smoking. Nearly one fourth of middle school and high school students who have never smoked cigarettes are susceptible to initiating cigarette smoking in the next year. Exposure to secondhand smoke (e.g., environmental tobacco smoke) is substantially higher among both middle

school and high school students. During the week before the survey, approximately 9 out of 10 current cigarette smokers and one half of never cigarette smokers were in the same room with someone who was smoking cigarettes; and 8 out of 10 current cigarette smokers and 3 out of 10 never cigarette smokers rode in a car with someone who was smoking cigarettes. Approximately 70% of middle school and 57% of high school students who currently smoke cigarettes live in a home where someone smokes cigarettes. Among never cigarette smokers, approximately 3 out of 10 live in a home where someone smokes cigarettes. Approximately 69% of middle school and 58% of high school students aged <18 years who currently smoke cigarettes were not asked to show proof of age when they bought or tried to buy cigarettes. Approximately 8 out of 10 middle school and high school students have seen antismoking commercials. Eight out of 10 middle school students report having seen actors using tobacco on television or in the movies, and approximately 11% of middle school and 16% of high school students who had never used tobacco would wear or use something with a tobacco company name or picture on it. This rate increases to nearly 60% for current tobacco users.

Public Health Actions: Youth Tobacco Survey data are used by health and education officials to improve national and state programs to prevent and control youth tobacco use. Several states use the data in presentations to their state legislators to demonstrate the need for funding smoking cessation and prevention programs for youth.

INTRODUCTION

Tobacco use is the single leading preventable cause of death in the United States (1), accounting for approximately 430,000 deaths each year (2). The prevalence of cigarette smoking nationwide among high school students increased during the 1990s (3), peaking during 1996–1997, then began a gradual decline (4). Approximately 80% of tobacco users initiate use before age 18 years (5). If this trend in early initiation of cigarette smoking continues, approximately 5 million children aged <18 years who are living today will die prematurely as adults because they began to smoke cigarettes during adolescence (2). The economic costs associated with tobacco use ranges from \$50 billion to \$73 billion per year in medical expenses (6,7) and \$47 billion in lost productivity (8).

CDC recommends that states establish and maintain comprehensive tobacco-control programs to reduce tobacco use among youth (9). Surveillance and evaluation are among primary components of a comprehensive tobacco-control program. To assist states in developing and maintaining their state-based comprehensive tobacco prevention and control programs, CDC developed the Youth Tobacco Surveillance and Evaluation System, which includes international, national, and state school-based surveys of middle school and high school students. Data in this report are presented from two components of this system: the National Youth Tobacco Survey (NYTS) and the state Youth Tobacco Surveys (YTS).

The NYTS and the YTS were developed to provide states with the data necessary to support the design, implementation, and evaluation of a comprehensive tobacco-control program. Several states have data regarding the prevalence of selected tobacco use behaviors among high schools students from the Youth Risk Behavior

Surveillance System (YRBSS). The YTS supplements the YRBSS by providing more comprehensive data regarding tobacco use (bidis*, cigarettes, cigars, kreteks*, pipes, and smokeless tobacco); exposure to secondhand smoke; smoking cessation; school curriculum; minors' ability to purchase or obtain tobacco products; knowledge and attitudes about tobacco and familiarity with protobacco and antitobacco media messages; and by providing information regarding both middle school and high school students. The YTS and YRBSS use identical sampling methodologies and the same wording for questions about tobacco use to enable states to use the high school data regarding tobacco use collected by both surveys.

The NYTS was first conducted during fall 1999 then again during spring 2000 (10) and is representative of all students in the 50 states and the District of Columbia. Funding for the NYTS was provided by the American Legacy Foundation (Washington, DC), and the survey was conducted by the CDC Foundation (Atlanta, Georgia). The NYTS will be conducted every other year. State Youth Tobacco Surveys were first conducted in 1998, when three states participated; in 1999, when 13 states participated; and in 2000, when 29 states participated. Some states conduct the YTS annually, whereas others conduct the survey every other year. This report summarizes data from the 2000 NYTS and 2000 state surveys.

METHODS

Sampling

National Youth Tobacco Survey

The 2000 NYTS employed a three-stage cluster sample design to produce a nationally representative sample of public and private school students in grades 6–12. The first-stage sampling frame contained 1,307 primary sampling units (PSUs) consisting of large counties or groups of smaller, adjacent counties. Of these, 148 PSUs were selected from eight strata formed on the basis of the degree of urbanization and the relative percentage of Asian, black, and Hispanic students in the PSU. The PSUs were selected with a probability proportional to weighted school enrollment. At the second sampling stage, 360 schools from the 148 PSUs were selected with a probability proportional to weighted school enrollment. Schools with substantial numbers of Asian, black, and Hispanic students were sampled at higher rates than all other schools through the use of a weighted measure of size. The third stage of sampling consisted of randomly selecting approximately five intact classes of a required subject (e.g., English or social studies) across grades 6–12 at each participating school. All students in the selected classes were eligible to participate in the survey.

A weighting factor was applied to each student record to adjust for nonresponse and for the varying probabilities of selection, including those resulting from the

*Bidis (or beedies) are small brown cigarettes from India consisting of tobacco wrapped in a leaf and tied with a thread. Kreteks (also called clove cigarettes) are flavored cigarettes containing tobacco and clove extract. Bidis and kreteks are two emerging forms of tobacco in the United States.

oversampling of Asian, black, and Hispanic students. The number of students in other racial and ethnic groups were too low for meaningful analysis in this report.* The weights were scaled so that a) the weighted count of students was equal to the total sample size, and b) the weighted proportions of students in each grade matched national population proportions. SUDAAN (11) was used to compute 95% confidence intervals, which were used to determine the differences between subgroups at the $p < 0.05$ level. Differences between prevalence estimates were considered statistically significant if the 95% confidence intervals did not overlap. Only subgroup comparisons that were statistically significant are mentioned in this report. The NYTS is representative of students in grades 6–12 in public and private schools in all 50 states and the District of Columbia. For the NYTS, 35,828 questionnaires were completed in 324 schools. The school response rate was 90.0%, and the student response rate was 93.4%, resulting in an overall response rate of 84.1%.

State Youth Tobacco Surveys

The state YTS employed a two-stage cluster sample design to produce representative samples of students in middle schools (grades 6–8) and high schools (grades 9–12). The first-stage sampling frame included separate lists for middle schools and high schools. In all except four states, private schools were excluded, and the list consisted of all public schools containing the appropriate grades (e.g., either 6–8 or 9–12). Schools were selected with a probability proportional to the school enrollment size. At the second sampling stage, classes were randomly selected from the list of classes obtained from each participating school. All students in the selected classes were eligible to participate in the survey.

In 2000, the sample sizes for the YTS ranged from 583 to 33,586. School response rates ranged from 56.8% to 100.0%; student response rates ranged from 54.1% to 96.4%; and overall response rates ranged from 45.8% to 91.8%. All except four state surveys had overall response rates of at least 60%; these data were weighted and are considered representative of the respective population (Table 1). Data for Hawaii middle schools and Arizona, Maine, and Vermont high schools were not representative (overall response rates <60%) and therefore are not included in this report.

Data Collection

The survey was administered during one class period. Procedures were designed to protect students' privacy by assuring that student participation was anonymous and voluntary. Students completed a self-administered questionnaire in the classroom, recording their responses on an answer sheet. The core questionnaire contained 63 questions; to meet their individual needs, some states added questions. Before the survey was administered, local parental permission was obtained and institutional review board criteria were followed. The core questionnaire included questions about tobacco use (bidis, cigarettes, cigars, kreteks, pipes, and smokeless tobacco), exposure to secondhand smoke, smoking cessation, school curriculum, minors' ability to purchase or obtain tobacco products, knowledge and attitudes about tobacco, and familiarity with protobacco and antitobacco media messages.

*Estimates are not reported when <35 cases are in the denominator.

RESULTS*

Prevalence of Use

Lifetime Tobacco Use[†]

Middle school and high school students were asked about their lifetime use of cigarettes, cigars, smokeless tobacco, pipes, bidis, or kreteks. Cigarettes were the most prevalent form of tobacco used, followed by cigars and smokeless tobacco.


Middle School. Nationally, 36.3% of students had ever smoked cigarettes (Figure 1). White (33.8%), black (44.0%), and Hispanic (40.0%) students were significantly more likely than Asian students (20.8%) to have ever smoked cigarettes (Table 2). Black students were significantly more likely than white students to have ever smoked cigarettes. Cigars were the second most prevalent form of tobacco used (19.3%), with male students (24.7%) significantly more likely than female students (14.1%) to have ever smoked cigars. White (18.5%), black (21.8%), and Hispanic (22.5%) students were significantly more likely than Asian students (9.2%) to have ever smoked cigars. Smokeless tobacco was the third most prevalent form of tobacco used by students (9.5%), with male students (14.8%) significantly more likely than female students (4.2%) to have used smokeless tobacco. White students (10.7%) were significantly more likely than black (6.1%) or Asian (4.1%) students to have ever used smokeless tobacco. Hispanic (7.5%) students were significantly more likely than Asian students to have ever used smokeless tobacco. Tobacco in a pipe was the fourth most prevalent form of tobacco used by students (6.7%), with male students (9.2%) significantly more likely than female students (4.3%) to have smoked tobacco in a pipe. White students (6.5%) were significantly more likely than black students (4.4%) to have ever smoked tobacco in a pipe. Hispanic students (9.7%) were significantly more likely than white, black, or Asian (4.7%) students to have ever smoked tobacco in a pipe. Bidis were the fifth most prevalent form of tobacco used, with 4.4% of students having reported that they had ever smoked these cigarettes. Male students (5.9%) were significantly more likely than female students (3.0%) to have ever smoked bidis. Black (5.7%) and Hispanic (6.4%) students were significantly more likely than white students (3.5%) to have ever smoked bidis. Kreteks were the sixth most prevalent form of tobacco used, with 3.6% of students having reported that they had ever smoked these cigarettes. Male students (4.6%) were significantly more likely than female students (2.6%) to have ever smoked kreteks.

Twenty-six states asked students if they had ever smoked cigarettes, cigars, or used smokeless tobacco. Among these 26 states, the percentage of students who had ever smoked cigarettes ranged from 26.3% in Maryland to 53.4% in Mississippi (public schools) (median: 38.3%) (Table 3). The percentage of students who had ever smoked cigars ranged from 14.3% in New York to 31.8% in Mississippi (public schools) (median: 20.6%). The percentage of students who had ever used smokeless tobacco ranged from

*Unless otherwise noted, the data in this report are for 2000. For Mississippi, the type of school (private or public) is identified in parentheses.

[†] Lifetime use was defined by asking a) Ever tried cigarettes, even one or two puffs? b) Ever tried smoking cigars, cigarillos, or little cigars, even one or two puffs? c) Ever used chewing tobacco, snuff, or dip, such as Redman,[®] Levi Garrett,[®] Beechnut,[®] Skoal,[®] Skoal Bandits,[®] or Copenhagen[®]? d) Ever tried smoking bidis, even one or two puffs? e) Ever tried smoking any of the following: bidis, kreteks, both, neither?

FIGURE 1. Percentage of all middle school and high school students who ever used tobacco*, by type of tobacco product — National Youth Tobacco Survey, 2000



* Ever use of cigarettes was determined by asking, "Have you ever tried cigarette smoking, even one or two puffs?" Ever use of cigars was determined by asking, "Have you ever tried smoking cigars, cigarillos, or little cigars, even one or two puffs?" Ever use of smokeless tobacco was determined by asking, "Have you ever used chewing tobacco, snuff, or dip, such as Redman,[®] Levi Garrett,[®] Beechnut,[®] Skoal,[®] Skoal Bandits,[®] or Copenhagen[®]?" Ever use of bidis was determined by asking, "Have you ever tried smoking bidis, even one or two puffs?" Ever use of kreteks was determined by asking, "Have you ever tried smoking kreteks, even one or two puffs?"

3.8% in New York to 24.7% in Wyoming (median: 10.8%). Among the 24 states that asked students if they had ever smoked bidis, the percentage of students who answered "yes" to this question ranged from 4.1% in Maine to 10.1% in Arizona (median: 6.2%). Among the 22 states that asked students if they had ever smoked kreteks, the percentage of students who answered "yes" to this question ranged from 2.3% in Minnesota to 5.1% in Arizona, Tennessee, and Wyoming (median: 3.9%).

High School. Nationally, 64.0% of students had ever smoked cigarettes (Figure 1). White (64.3%), black (62.5%), and Hispanic (66.4%) students were significantly more likely than Asian students (52.4%) to have ever smoked cigarettes (Table 2). Cigars were the second most prevalent form of tobacco used (41.7%), with male students (52.2%) significantly more likely than female students (30.9%) to have ever smoked cigars. White students (44.1%) were significantly more likely than black (35.8%) or Asian (24.3%) students to have ever smoked cigars. Black and Hispanic (39.5%) students were significantly more likely than Asian students to have ever smoked cigars. Smokeless tobacco was the third most prevalent form of tobacco used by students (18.0%), with male students (29.0%) significantly more likely than female students (6.8%) to have ever used smokeless tobacco. White students (22.1%) were significantly more likely than black (7.6%), Hispanic (11.0%), or Asian (6.8%) students to have ever used smokeless tobacco. Hispanic students were significantly more likely than Asian students to have ever used smokeless tobacco. Bidis were the fourth most prevalent form of tobacco used (12.9%), with male students (15.7%) significantly more likely than female students (9.9%) to have ever smoked these cigarettes. Black (15.7%) and Hispanic (15.2%) students were significantly more likely than white (11.6%) or Asian (10.1%) students to have ever smoked bidis. Kreteks were the fifth most prevalent form of tobacco used by students (12.1%), with male students (14.0%) significantly more likely than female students (10.2%) to have ever smoked these cigarettes. White students (14.4%) were significantly more likely than black (4.2%), Hispanic (9.6%), or Asian (7.5%) students to have ever smoked kreteks. Hispanic students were significantly more likely than black students to have ever smoked these cigarettes. Tobacco in a pipe was the sixth most prevalent form of tobacco used, with 11.0% of students having reported that they had ever used this form of tobacco. Male students (16.7%) were significantly more likely than female students (5.0%) to have ever smoked tobacco in a pipe. White (12.3%) and Hispanic (10.3%) students were significantly more likely than black (5.3%) or Asian students (6.3%) to have ever smoked tobacco in a pipe.


Twenty-four states asked students if they had ever smoked cigarettes, cigars, used smokeless tobacco, or smoked bidis. Among these 24 states, the percentage of students who had ever smoked cigarettes ranged from 56.9% in Florida to 74.3% in Kentucky and West Virginia (median: 64.7%) (Table 3). The percentage of students who had ever smoked cigars ranged from 24.0% in the District of Columbia to 54.4% in Kentucky (median: 45.1%). The percentage of students who had ever used smokeless tobacco ranged from 9.9% in the District of Columbia to 38.0% in Mississippi (private schools) (median: 24.7%). The percentage of students who had ever smoked bidis ranged from 7.0% in Nebraska to 24.0% in Delaware (median: 10.5%). Among the 20 states that asked students if they had ever smoked kreteks, the percentage of students who answered "yes" to this question ranged from 3.6% in Kentucky to 13.7% in Colorado (median: 7.0%).

Current Tobacco Use

Students were asked if they had used cigarettes, cigars, smokeless tobacco, tobacco in pipes, bidis, or kreteks on ≥ 1 of the 30 days preceding the survey. Current use of any tobacco product is defined as use of cigarettes or cigars or smokeless tobacco or tobacco in a pipe or bidis or kreteks on ≥ 1 of the 30 days preceding the survey.

Middle School. Nationally, 15.1% of students were current users of any tobacco product (Figure 2), with male students (17.6%) significantly more likely than female students (12.7%) to be current users of any tobacco product (Table 4). White (14.3%), black (17.5%), and Hispanic (16.0%) students were significantly more likely than Asian (7.5%) students to currently use any tobacco product. Among current users, cigarettes were the most prevalent form of tobacco used (11.0%), with white (10.8%), black (11.2%), and Hispanic (11.4%) students significantly more likely than Asian (5.3%) students to currently smoke cigarettes. Cigars were the second most prevalent form of tobacco used (7.1%), with male students (9.7%) significantly more likely than female students (4.6%) to currently smoke cigars. Black (9.8%) and Hispanic (8.8%) students were significantly more likely than white (6.1%) or Asian (4.1%) students to currently smoke cigars. Smokeless tobacco was the third most prevalent form of tobacco used (3.6%), with male students (5.7%) significantly more likely than female students (1.5%) to currently use smokeless tobacco. Tobacco in a pipe was the fourth most prevalent form of tobacco used (3.0%), with male students (4.3%) significantly more likely than female students (1.8%) to currently smoke tobacco in a pipe. Bidis were the fifth most prevalent form of tobacco used, with 2.4% of students reporting that they currently were

FIGURE 2. Percentage of all middle school and high school students who were current users of any tobacco product,* by type of tobacco product — National Youth Tobacco Survey, 2000



* Use of cigarettes or cigars or smokeless tobacco or pipes or bidis or kreteks on ≥ 1 of the 30 days preceding the survey.

smoking these cigarettes. Male students (3.4%) were significantly more likely than female students (1.4%) to currently smoke bidis, and Hispanic students (3.6%) were significantly more likely than white students (1.9%) to currently smoke bidis. Kreteks were the sixth most prevalent form of tobacco used, with 2.1% of students reporting that they currently smoke these cigarettes. Male students (2.7%) were significantly more likely than female students (1.5%) to currently smoke kreteks.

Twenty-six states asked students if they were current users of any tobacco product. Among these 26 states, the percentage of students who were current users of any form of tobacco ranged from 10.0% in California to 28.3% in Kentucky (median: 16.1%) (Table 5). The percentage of students who currently smoke cigarettes ranged from 6.7% in California to 21.5% in Kentucky (median: 12.0%). The percentage of students who currently smoke cigars ranged from 3.7% in Minnesota to 12.4% in Alabama (median: 6.1%). Current users of smokeless tobacco ranged from 1.5% in New York to 10.1% in Mississippi (private schools) (median: 3.6%). Twenty-three states asked students if they currently smoke tobacco in a pipe or bidis. Among these 23 states, the percentage of students who reported they currently smoke tobacco in a pipe ranged from 1.7% in Kansas to 4.8% in Tennessee (median: 3.4%). The percentage of students who reported they currently smoke bidis ranged from 2.2% in New York to 5.0% in the District of Columbia and West Virginia (median: 3.5%).

High School. Nationally, 34.5% of students were current users of any tobacco product (Figure 2), with white students (38.0%) significantly more likely than black (26.5%), Hispanic (28.4%), or Asian (22.9%) students to currently use any tobacco product (Table 4). Among current users, the most prevalent form of tobacco used were cigarettes (28.0%), with white students (31.8%) significantly more likely than black (16.8%), Hispanic (22.6%), or Asian (20.6%) students to currently smoke cigarettes. Hispanic students were significantly more likely than black students to currently smoke cigarettes. Cigars were the second most prevalent form of tobacco used (14.8%), with male students (22.0%) significantly more likely than female students (7.3%) to currently smoke cigars. White (15.1%), black (15.3%), and Hispanic (13.6%) students were significantly more likely than Asian students (7.4%) to currently smoke cigars. Smokeless tobacco was the third most prevalent form of tobacco used (6.6%), with male students (11.8%) significantly more likely than female students (1.4%) to currently use smokeless tobacco. White students (8.2%) were significantly more likely than black (2.6%), Hispanic (4.0%), or Asian (1.9%) students to currently use smokeless tobacco. Kreteks were the fourth most prevalent form of tobacco used (4.2%), with male students (5.3%) significantly more likely than female students (3.0%) to currently smoke these cigarettes. White (4.5%) and Hispanic (4.0%) students were significantly more likely than black students (2.2%) to currently smoke kreteks. Bidis were the fifth most prevalent form of tobacco used (4.1%), with male students (5.4%) significantly more likely than female students (2.8%) to currently smoke these cigarettes. Hispanic students (5.7%) were significantly more likely than white (3.6%) or Asian (3.0%) students to currently smoke bidis. Tobacco in a pipe was the sixth most prevalent form of tobacco used (3.3%). Male students (5.2%) were significantly more likely than female students (1.4%) to report that they currently smoke tobacco in a pipe, and Hispanic students (4.2%) were significantly more likely than black students (2.2%) to currently smoke tobacco in a pipe.

Twenty-six states asked students if they were current users of any tobacco product. Among these 26 states, the percentage of students who were current users of any form of tobacco ranged from 21.0% in the District of Columbia to 50.4% in Mississippi (private schools) (median: 36.9%) (Table 5). The percentage of students who currently smoke cigarettes ranged from 14.7% in the District of Columbia to 41.7% in Mississippi (private schools) (median: 29.2%). The percentage of students who currently smoke cigars ranged from 4.9% in Hawaii to 20.0% in Arkansas (median: 15.1%). Current users of smokeless tobacco ranged from 3.3% in California and the District of Columbia to 19.3% in Mississippi (private schools) (median: 9.1%). Twenty-five states asked students if they currently smoke tobacco in a pipe or bidis. Among these states, the percentage of students who reported that they currently smoke tobacco in a pipe ranged from 2.4% in Delaware to 5.6% in Mississippi (private schools) (median: 4.2%). The percentage of students who reported that they currently smoke bidis ranged from 3.4% in Alabama and Kansas to 8.5% in West Virginia (median: 5.2%).

Ever Daily Cigarette Use*

Middle School. Nationally, 5.5% of students had ever smoked cigarettes daily (Figure 3). White students (6.0%) were significantly more likely than black (3.7%) or Asian (2.9%) students to have ever smoked cigarettes daily (Table 6).

Among the 26 states that asked this question, the percentage of students who had ever smoked cigarettes daily ranged from 2.7% in the District of Columbia to 14.0% in Kentucky (median: 7.5%) (Table 7).

High School. Nationally, 20.6% of students had ever smoked cigarettes daily (Figure 3). White students (23.8%) were significantly more likely than black (11.7%), Hispanic (14.1%), or Asian (15.3%) students to have ever smoked cigarettes daily (Table 6).

Among the 24 states that asked this question, the percentage of students who had ever smoked cigarettes daily ranged from 8.7% in the District of Columbia to 33.4% in West Virginia (median: 22.3%) (Table 7).

Number of Cigarettes Smoked Per Day Among Current Smokers

Middle School. Nationally, 16.1% of current cigarette smokers had smoked ≥ 6 cigarettes per day on the days they smoked (Figure 3). White smokers (18.6%) were significantly more likely than Hispanic smokers (11.3%) to smoke ≥ 6 cigarettes per day on the days they smoked (Table 6).


Among the 26 states that asked this question, the percentage who smoked ≥ 6 cigarettes per day on the days they smoked ranged from 11.1% in Texas to 27.9% in West Virginia (median: 14.8%) (Table 7).

High School. Nationally, 31.7% of current cigarette smokers had smoked ≥ 6 cigarettes per day on the days they smoked (Figure 3). White students (34.4%) were significantly more likely than black (22.6%) or Hispanic (19.0%) students to smoke ≥ 6 cigarettes per day on the days they smoked (Table 6).

Among the 24 states that asked this question, the percentage of current smokers who smoked ≥ 6 cigarettes per day on the days they smoked ranged from 18.5% in the District of Columbia to 42.0% in West Virginia (median: 30.6%) (Table 7).

*Ever smoked cigarettes daily was defined as having ever smoked at least one cigarette every day for 30 days.

FIGURE 3. Percentage of all middle school and high school student who ever smoked cigarettes daily* and current cigarette smokers who smoked ≥ 6 cigarettes per day on the days they smoked[†] — National Youth Tobacco Survey, 2000



* Students were asked, "Have you ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days?"

[†] Students were asked, "During the past 30 days, on the days you smoked, how many cigarettes did you smoke per day?"


Age of Initiation of Tobacco Use*

Cigarettes

Middle School. Nationally, 8.4% of students first smoked a cigarette before age 11 years (Figure 4), with male students (10.0%) significantly more likely than female students (6.8%) to have first smoked a cigarette before age 11 years (Table 8).

*Age of initiation was determined by asking a) How old were you when you smoked a whole cigarette for the first time? b) How old were you when you when you smoked a cigar, cigarillo, or little cigar for the first time? c) How old were you when you first used chewing tobacco, snuff, or dip for the first time? Questions on age of initiation were not asked for pipes, bidis, or kreteks.

FIGURE 4. Percentage of all middle school and high school students who first used* tobacco before age 11 years, by tobacco product — National Youth Tobacco Survey, 2000



* Age of initiation was determined by asking students these questions: "How old were you when you smoked a whole cigarette for the first time?" "How old were you when you smoked a cigar, cigarillo, or little cigar for the first time?" "How old were you when you used chewing tobacco, snuff, or dip for the first time?"

Among the 25 states that asked this question, the percentage of students who initiated cigarette smoking before age 11 years ranged from 5.7% in New York to 19.1% in Kentucky (median: 10.7%) (Table 9).

High School. Nationally, 6.7% of students first smoked a cigarette before the age of 11 years (Figure 4), with male students (8.3%) significantly more likely than female students (5.1%) to have first smoked a cigarette before age 11 years (Table 8).

Among the 23 states that asked this question, the percentage who started smoking cigarettes before age 11 years ranged from 5.8% in the District of Columbia to 14.4% in West Virginia (median: 14.3%) (Table 9).

Cigars

Middle School. Nationally, 5.2% of students smoked a cigar before age 11 years (Figure 4), with male students (6.9%) significantly more likely than female students (3.5%) to have first smoked a cigar before age 11 years (Table 8).

Among the 25 states that asked this question, the percentage of students who first smoked a cigar before the age of 11 years ranged from 3.0% in Delaware to 9.3% in Wyoming (median: 5.9%) (Table 9).

High School. Nationally, 3.1% of students first smoked a cigar before the age of 11 years (Figure 4), with male students (4.5%) significantly more likely than female students (1.7%) to report initiating cigar smoking before the age of 11 years (Table 8). White students (3.1%) were significantly more likely than Asian students (1.6%) to have initiated cigar smoking before the age of 11 years. Hispanic students (4.0%) were significantly more likely than black (2.1%) or Asian students to have initiated cigar smoking before the age of 11 years.

Among the 23 states that asked this question, the percentage of students who first smoked a cigar before age 11 years ranged from 2.6% in Delaware to 6.9% in West Virginia (median: 4.6%) (Table 9).

Smokeless Tobacco

Middle School. Nationally, 3.7% of students reported that they first used smokeless tobacco before age 11 years (Figure 4), with male students (6.0%) significantly more likely than female students (1.6%) to have first used smokeless tobacco before age 11 years (Table 8). White students (4.2%) were significantly more likely than Asian students (1.5%) to have first used smokeless tobacco before age 11 years.

Among the 25 states that asked this question, the percentage of students who initiated the use of smokeless tobacco before age 11 years ranged from 1.6% in the District of Columbia to 11.1% in Kentucky (median: 4.2%) (Table 9).

High School. Nationally, 2.9% of students reported first using smokeless tobacco before the age of 11 years (Figure 4), with male students (4.7%) significantly more likely than female students (1.0%) to report using smokeless tobacco before the age of 11 years (Table 8). White students (3.3%) were significantly more likely than black (1.8%), Hispanic (1.7%), or Asian (1.0%) students to have first used smokeless tobacco before the age of 11 years.

Among the 23 states that asked this question, the percentage ranged from 2.0% in California and Connecticut to 10.9% in Mississippi (private schools) (median: 4.5%) (Table 9).

Established Use of Tobacco Products


Smoked ≥ 100 Cigarettes in Lifetime

Middle School. Nationally, 4.1% of ever cigarette smokers had smoked ≥ 100 cigarettes during their lifetime (Figure 5), with male students (5.0%) significantly more likely than female students (3.2%) to have smoked ≥ 100 cigarettes during their lifetime. White students (4.8%) were significantly more likely than black (2.2%), Hispanic (2.7%), or Asian (2.4%) students to have smoked ≥ 100 cigarettes during their lifetime (Table 10).

Among the 24 states that asked this question, the percentage of ever cigarette smokers who smoked ≥ 100 cigarettes in their lifetime ranged from 3.5% in the District of Columbia to 21.6% in Maine (median: 13.5%) (Table 11).

High School. Nationally, 20.7% of ever cigarette smokers had smoked ≥ 100 cigarettes in their lifetime (Figure 5). White students (24.9%) were significantly more likely than black (8.6%), Hispanic (13.4%), or Asian (14.9%) students to have smoked ≥ 100 cigarettes in their lifetime (Table 10). Hispanic and Asian students were significantly more likely than black students to have smoked ≥ 100 cigarettes in their lifetime.

FIGURE 5. Percentage of middle school and high school students who ever smoked cigarettes who smoked ≥ 100 cigarettes in their lifetime and percentage of all students who frequently smoked cigarettes* — National Youth Tobacco Survey, 2000



* Smoked cigarettes on ≥ 20 of the 30 days preceding the survey.

Among the 22 states that asked this question, the percentage of ever cigarette smokers ranged from 8.4% in the District of Columbia to 45.6% in Mississippi (private schools) (median: 34.5%) (Table 11).

Frequent Use* of Tobacco

Middle School. Nationally, 2.7% of students had frequently smoked cigarettes (Figure 5). White students (3.1%) were significantly more likely than Hispanic students (1.5%) to be frequent cigarette smokers (Table 10). Students who were frequent users of cigars, smokeless tobacco, pipes, bidis, and kreteks accounted for $\leq 1\%$ of students.

Among the 26 states that asked this question, the percentage of students who were frequent cigarette smokers ranged from 0.8% in the District of Columbia to 7.4% in West Virginia (median: 3.5%) (Table 11). The percentage of students who were frequent cigar smokers ranged from 0.2% in Colorado, Minnesota, Vermont, and Wisconsin to 1.7% in Maine (median: 0.9%), though some confidence intervals overlapped. The percentage of students who frequently used smokeless tobacco ranged from 0.4% in

*Frequent use of tobacco was defined as having used a tobacco product on ≥ 20 of the 30 days preceding the survey. The survey included questions regarding frequent use of cigarettes, cigars, smokeless tobacco, and pipes.

Connecticut and the District of Columbia to 3.1% in Kentucky (median: 0.8%). The percentage of students who smoked tobacco in a pipe ranged from 0.2% in Kansas, Kentucky, Minnesota, and Wisconsin to 1.2% in Maine and Texas (median: 0.6%), though some confidence intervals overlapped. The percentage of students who frequently smoked bidis ranged from 0.1% in Vermont to 1.6% in Maine (median: 0.5%).

High School. Nationally, 13.6% of students had frequently smoked cigarettes (Figure 5). White students (16.4%) were significantly more likely than black (6.1%), Hispanic (7.4%), or Asian (9.9%) students to have frequently smoked cigarettes (Table 10). Students who were frequent users of cigars, smokeless tobacco, pipes, bidis, and kreteks, accounted for $\leq 2\%$ of students.

Among the 24 states that asked this question, the percentage of students who were frequent cigarette smokers ranged from 3.5% in the District of Columbia to 22.1% in West Virginia (median: 14.2%) (Table 11). The percentage of students who were frequent cigar smokers ranged from 0.5% in Connecticut to 2.1% in Mississippi (public schools) (median: 1.4%). The percentage of students who frequently used smokeless tobacco ranged from 0.5% in Connecticut to 8.0% in Mississippi (private schools) (median: 2.8%). The percentage of students who frequently smoked tobacco in a pipe ranged from 0.3% in South Dakota to 1.6% in New York (median: 0.8%), though their confidence intervals overlapped. The percentage of students who frequently smoked bidis ranged from 0.3% in the District of Columbia to 1.3% in Maryland, Mississippi (private schools), and New York (median: 0.7%), though some confidence intervals overlapped.

Brand of Cigarettes Usually Smoked

Current cigarette smokers were asked to identify the brand of cigarettes they usually smoked. Brand preferences were similar among current smokers in middle and high schools.


Middle School. Nationally, 47.1% of current cigarette smokers in middle school identified Marlboro® as the brand they usually smoked, followed by Newport® (22.4%), other brands (12.3%), and Camel® (6.3%) (Figure 6). Marlboro® was the brand most often smoked by white (55.6%), Hispanic (53.9%), and Asian (37.3%) students, whereas Newport® was the brand most often smoked by black students (64.1%) (Table 12).

Among the 25 states that asked this question, the percentage of current cigarette smokers who most often smoked Marlboro® ranged from 9.2% in the District of Columbia to 66.5% in Mississippi (private schools) (median: 54.6%) (Table 13). The percentage of current cigarette smokers who preferred Newport® ranged from 3.7% in Wyoming to 65.9% in the District of Columbia (median: 10.7%).

High School. Nationally, 53.3% of current cigarette smokers in high school identified Marlboro® as the brand they usually smoked, followed by Newport® (20.8%) and Camel® (11.0%) (Figure 6). Marlboro® was the brand most often smoked by white (59.4%), Hispanic (49.1%), and Asian (54.2%) students, whereas Newport® was the favored brand among black students (74.0%) (Table 12).

Among the 25 states that asked this question, the percentage of current cigarette smokers who preferred Marlboro® ranged from 5.8% in the District of Columbia to 75.7% in Mississippi (private schools) (median: 58.3%) (Table 13). The percentage of current cigarette smokers who preferred Newport® ranged from 5.4% in Hawaii to 76.3% in the District of Columbia (median: 13.2%).

FIGURE 6. Percentage of usual brand of cigarettes smoked in the 30 days preceding the survey by current cigarette smokers* in middle school and high school — National Youth Tobacco Survey, 2000



* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Includes KOOL®, Virginia Slims®, GPC®, Basic®, and Doral®.

Smoking Susceptibility*

Susceptibility Among Never Cigarette Smokers

Never cigarette smokers were classified as not susceptible to smoking cigarettes if they responded that a) they would *not* smoke a cigarette soon; and b) they would *definitely not* smoke a cigarette in the next year; and c) they would *definitely not* smoke if their closest friend offered them a cigarette. All other students were classified as susceptible to initiating cigarette smoking in the next year.

Middle School. Nationally, 24.3% of students who had never smoked cigarettes were classified as susceptible to initiating cigarette smoking in the next year (Table 14). Hispanic students (28.0%) were significantly more likely than white students (23.2%) to be susceptible to initiating cigarette smoking in the next year.

*This definition for susceptibility was developed by Pierce and colleagues (12).

Among the 26 states that asked this series of questions, the percentage of never cigarette smokers who were susceptible to initiating cigarette smoking ranged from 19.9% in Indiana to 31.8% in Wisconsin (median: 25.8%) (Table 15).

High School. Nationally, 22.5% of students who had never smoked cigarettes were classified as susceptible to initiating cigarette smoking in the next year (Table 14).

Among the 24 states that asked this series of questions, the percentage of never cigarette smokers who were susceptible to initiating cigarette smoking ranged from 17.3% in Alabama to 33.5% in Mississippi (private schools) (median: 23.5%) (Table 15).

Knowledge and Attitudes

Peer Group Use of Tobacco

One or More Closest Friends Smoke Cigarettes

Middle School. Nationally, current cigarette smokers (90.3%) were significantly more likely than students who had never smoked cigarettes (17.0%) to report that ≥ 1 of their closest friends smoked cigarettes (Figure 7). White (16.3%), black (19.7%), and Hispanic (19.0%) students who had never smoked cigarettes were significantly more likely than Asian students (9.5%) who had never smoked cigarettes to report that ≥ 1 of their closest friends smoked cigarettes (Table 16).

Among the 26 states that asked never cigarette smokers if ≥ 1 of their closest friends smoked cigarettes, the percentage of students who answered "yes" ranged from 10.7% in Minnesota to 22.2% in Mississippi (public schools) (median: 15.1%). Among the 26 states that asked current cigarette smokers if ≥ 1 of their closest friends smoked cigarettes, the percentage of students who answered "yes" ranged from 71.7% in Indiana to 95.3% in Delaware (median: 86.9%) (Table 17).


High School. Nationally, current cigarette smokers (92.3%) were significantly more likely than students who had never smoked cigarettes (33.0%) to report that ≥ 1 of their closest friends smoked cigarettes (Figure 7). Hispanic (34.6%) students who never smoked cigarettes were significantly more likely than Asian (25.6%) students to report that ≥ 1 of their closest friends smoked cigarettes (Table 16). White (93.5%), Hispanic (90.4%), and Asian (95.0%) students who currently smoke cigarettes were significantly more likely than black current cigarette smokers (82.8%) to report that ≥ 1 of their closest friends smoked cigarettes.

Among the 24 states that asked never cigarette smokers if ≥ 1 of their closest friends smoked cigarettes, the percentage of students who answered "yes" ranged from 22.3% in Mississippi (public schools) to 37.2% in Arkansas and Delaware (median: 29.6%). Among the 24 states that asked current cigarette smokers if ≥ 1 of their closest friends smoked cigarettes, the percentage of students who answered "yes" ranged from 67.3% in the District of Columbia to 96.4% in Delaware (median: 91.4%) (Table 17).

One or More Closest Friends Use Smokeless Tobacco

Middle School. Nationally, current users of smokeless tobacco (78.0%) were significantly more likely than students who had never used this product (7.3%) to report that ≥ 1 of their closest friends used smokeless tobacco (Figure 7). Among current users, white students (83.5%) were significantly more likely than black students (54.3%) to report that at least one of their closest friends used smokeless tobacco (Table 16).

FIGURE 7. Percentage of middle school and high school students with peers who use tobacco, by tobacco use status — National Youth Tobacco Survey, 2000



* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Used smokeless tobacco on ≥ 1 of the 30 days preceding the survey.

Among the 26 states that asked this question, the percentage of never smokeless tobacco users who reported that ≥ 1 of their closest friends used smokeless tobacco ranged from 4.3% in Kansas to 25.9% in West Virginia (median: 15.1%). The percentage of current smokeless tobacco users who reported that ≥ 1 of their closest friends used smokeless tobacco ranged from 52.1% in Indiana to 94.3% in New York (median: 74.0%) (Table 17).

High School. Nationally, current users of smokeless tobacco (80.7%) were significantly more likely than students who had never used this product (13.1%) to report that ≥ 1 of their closest friends used smokeless tobacco (Figure 7). Among never smokeless tobacco users, white students (16.4%) were significantly more likely than black (5.5%),

Hispanic (9.4%), or Asian (6.3%) students to report that ≥ 1 of their closest friends used smokeless tobacco, and Hispanic students were significantly more likely than black and Asian students to report that ≥ 1 of their closest friends used smokeless tobacco. Among current smokeless tobacco users, male students (82.2%) were significantly more likely than female students (67.6%) to report that ≥ 1 of their closest friends used smokeless tobacco, and white students (83.0%) were significantly more likely than black (65.0%) and Hispanic (71.7%) students to report that ≥ 1 of their closest friends used smokeless tobacco.

Among the 24 states that asked this question, the percentage of never smokeless tobacco users who reported that ≥ 1 of their closest friends used smokeless tobacco ranged from 3.8% in the District of Columbia to 41.0% in Mississippi (private schools) (median: 18.9%). The percentage of current users who reported that ≥ 1 of their closest friends used smokeless tobacco ranged from 50.2% in the District of Columbia to 90.9% in Mississippi (private schools) (median: 81.5%) (Table 17).

Social Perceptions About Cigarette Use

Think Smokers Have More Friends


Middle School. Nationally, current cigarette smokers (37.9%) were significantly more likely than students who had never smoked cigarettes (11.9%) to think that cigarette smokers have more friends (Figure 8). Among never cigarette smokers, black students (21.5%) were significantly more likely than white (9.2%) or Hispanic (14.3%) students to think that cigarette smokers have more friends, and Hispanic and Asian (15.6%) students were significantly more likely than white students to think that cigarette smokers have more friends (Table 18). Among current cigarette smokers, male students (42.6%) were significantly more likely than female students (32.7%) and black students (45.6%) were significantly more likely than white students (34.5%) to think that cigarette smokers have more friends.

Among the 26 states that asked this question, the percentage of never cigarette smokers who thought that cigarette smokers had more friends ranged from 6.2% in Kansas to 26.8% in the District of Columbia (median: 10.5%). The percentage of current cigarette smokers who thought that cigarette smokers had more friends ranged from 31.9% in Florida to 53.0% in Indiana (median: 40.7%) (Table 19).

High School. Nationally, current cigarette smokers (24.0%) were significantly more likely than students who had never smoked cigarettes (12.6%) to think cigarette smokers have more friends (Figure 8). Among never cigarette smokers, male students (15.4%) were significantly more likely than female students (9.9%) to think that cigarette smokers have more friends (Table 18). Black (21.0%), Hispanic (19.5%), and Asian (20.1%) students were significantly more likely than white students (9.0%) to think that cigarette smokers have more friends. Among current cigarette smokers, male students (29.1%) were significantly more likely than female students (18.5%) to think that cigarette smokers have more friends, and black (37.6%), Hispanic (30.7%), and Asian (39.0%) students were significantly more likely than white students (20.6%) to think that cigarette smokers have more friends.

Among the 24 states that asked this question, the percentage of never cigarette smokers who thought that cigarette smokers had more friends ranged from 8.5% in Colorado, Nebraska, and South Dakota to 24.2% in the District of Columbia

FIGURE 8. Percentage of middle school and high school students with social perceptions about cigarette use, by smoking status — National Youth Tobacco Survey, 2000



* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

(median: 13.0%). The percentage of current cigarette smokers who thought that cigarette smokers had more friends ranged from 22.2% in Delaware to 49.4% in the District of Columbia (median: 27.6%) (Table 19).


Access and Enforcement

Methods of Obtaining Cigarettes

How Current Cigarette Smokers Usually Got Cigarettes

Middle School. Nationally, current cigarette smokers aged <18 years usually got their cigarettes by having someone else buy cigarettes for them (24.4%), borrowing them from someone (22.4%), or being given them by someone aged 18 years or older (12.3%) (Figure 9). Eleven percent of current cigarette smokers reported they usually took their cigarettes from a family member or store. White (26.6%) and Asian (23.3%) students were more likely to get their cigarettes by having someone else buy them for them, whereas black (20.3%) and Hispanic (24.7%) students were more likely to get them some other way (Table 20).

FIGURE 9. How current cigarette smokers* aged <18 years in middle school and high school usually obtained cigarettes — National Youth Tobacco Survey, 2000



*Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Among the 25 states that asked this question, the percentage of current cigarette smokers who had someone else buy cigarettes for them ranged from 14.5% in the District of Columbia to 33.7% in Mississippi (private schools) (median: 23.0%). The percentage who borrowed cigarettes ranged from 19.8% in California to 38.1% in Ohio (median: 26.1%). The percentage who were given cigarettes by someone aged 18 years old or older ranged from 4.2% in Maine to 14.0% in Texas (median: 8.8%). The percentage who took their cigarettes from a family member or store ranged from 7.4% in Indiana to 19.0% in Minnesota (median: 12.0%). The percentage who got their cigarettes some other way ranged from 9.2% in Ohio to 31.2% in California (median: 19.0%) (Table 21).

High School. Nationally, current cigarette smokers aged <18 years usually got their cigarettes by buying their cigarettes in a store (32.2%), having someone else buy cigarettes for them (25.1%), or borrowing them from someone (20.9%) (Figure 9). Only 2.9% of current cigarette smokers reported they took their cigarettes from a family member or store. Male current cigarette smokers were more likely to buy their cigarettes in a store (35.9%), whereas female current cigarette smokers were more likely to have someone else buy their cigarettes (29.3%) (Table 20).

Among the 24 states that asked this question, the percentage of current cigarette smokers who usually purchased their cigarettes in a store ranged from 8.1% in South Dakota to 30.6% in Delaware (median: 20.4%). The percentage who had someone else

buy cigarettes for them ranged from 8.4% in the District of Columbia to 45.0% in South Dakota (median: 32.3%). The percentage who usually borrowed cigarettes ranged from 19.9% in Arkansas to 33.5% in Colorado (median: 24.9%) (Table 21).

Where Current Cigarette Smokers Bought Their Last Pack of Cigarettes

Cigarettes

Middle School. Nationally, 39.5% of current cigarette smokers aged <18 years bought their last pack of cigarettes at a gas station, followed by a convenience store (23.5%), a vending machine (12.6%), and a grocery store (10.4%) (Figure 10). This pattern held for all sex and race/ethnicity groups (Table 22).

Among the 20 states that asked this question, the percentage of current cigarette smokers who purchased their last pack of cigarettes at a gas station ranged from 9.5% in Colorado to 34.2% in Maine (median: 20.8%). The percentage of current cigarette smokers who purchased their last pack of cigarettes at a convenience store ranged from 5.9% in Minnesota to 19.3% in the District of Columbia (median: 12.3%) (Table 23).

High School. Nationally, 57.1% of current cigarette smokers aged <18 years bought their last pack of cigarettes at a gas station, followed by a convenience store (26.3%), a grocery store (7.7%), and a discount store (2.9%) (Figure 10). This pattern held for all sex and race/ethnicity groups (Table 22).

Among the 20 states that asked this question, the percentage of current cigarette smokers who purchased their last pack of cigarettes at a gas station ranged from 14.3 in Hawaii to 66.8% in Wisconsin (median: 45.8%). The percentage of current cigarette smokers who purchased their last pack of cigarettes at a convenience store ranged from 13.6% in Indiana to 35.9% in the District of Columbia (median: 20.9%) (Table 23).

Age and Access

Acquiring Cigarettes and Proof of Age

Proof of Age Not Required to Purchase Cigarettes

Middle School. Nationally, 69.4%* of current cigarette smokers were not asked to show proof of age when purchasing cigarettes (Figure 11 and Table 24).


Among the 25 states that asked this question, the percentage of current cigarette smokers who were not asked to show proof of age when purchasing cigarettes ranged from 56.7% in the District of Columbia to 90.5% in New Hampshire (median: 74.0%) (Table 25).

High School. Nationally, 58.1%† of current cigarette smokers were not asked to show proof of age when purchasing cigarettes (Figure 11). White (56.1%), black (64.9%), and Hispanic (67.5%) students were significantly more likely than Asian students (36.4%) to report that they were not asked to show proof of age when purchasing cigarettes (Table 24).

*Eleven percent of middle school students reported current cigarette smoking. Of this group, 7.3% obtained their cigarettes by buying them in a store, 69.4% of whom were not asked to show proof of age when making the purchase.

† Twenty-eight percent of high school students reported current cigarette smoking. Of this group, 32.3% obtained their cigarettes by buying them in a store, 58.1% of whom were not asked to show proof of age when making the purchase.

FIGURE 10. Where current cigarette smokers* aged <18 years in middle school and high school bought their last pack of cigarettes — National Youth Tobacco Survey, 2000



* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Among the 24 states that asked this question, the percentage of current cigarette smokers who were not asked to show proof of age when purchasing cigarettes ranged from 46.9% in Mississippi (private schools) to 68.3% in Nebraska (median: 59.5%) (Table 25).

Not Refused Purchase Because of Age

Middle School. Nationally, 62.4%* of current cigarette smokers aged <18 years were not refused purchase of cigarettes because of their age regardless of whether or not they were asked for proof of age (Figure 11 and Table 24).


Among the 23 states that asked this question, the percentage of current cigarette smokers aged <18 years who were not refused purchase ranged from 52.1% in California to 78.8% in Minnesota (median: 66.1%) (Table 25).

High School. Nationally, 58.1%† of current cigarette smokers aged <18 years were not refused purchase of cigarettes because of their age regardless of whether or not they were asked for proof of age (Figure 11). Female students (62.9%) were

*Eleven percent of middle school students reported current cigarette smoking. Of this group, 7.3% obtained their cigarettes by buying them in a store, 62.4% of whom were not refused purchase because of their age.

† Twenty-eight percent of high school students reported current cigarette smoking. Of this group, 32.3% obtained their cigarettes by buying them in a store, 58.1% of whom were not refused purchase because of their age.

FIGURE 11. Percentage of current cigarette smokers* aged <18 years in middle school and high school who purchased cigarettes in a store and were not asked to show proof of age or who were not refused purchase because of their age — National Youth Tobacco Survey, 2000



*Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

significantly more likely than male students (54.4%) to report that they were not refused purchase of cigarettes because of their age (Table 24).

Among the 24 states that asked this question, the percentage of current cigarette smokers aged <18 years who were not refused purchase of cigarettes because of their age ranged from 49.7% in Connecticut to 70.6% in Nebraska (median: 60.3%) (Table 25).

Media and Advertising

Exposure to Tobacco-Related Media and Advertising

Saw Antismoking Commercials on Television or Heard Them on the Radio


Middle School. Nationally, 81.5% of students saw antismoking commercials on television or heard antismoking commercials on the radio during the 30 days preceding

the survey (Figure 12), with female students (84.0%) significantly more likely than male students (78.9%) to have seen antismoking commercials on television or heard antismoking commercials on the radio (Table 26) during this time frame. White students (83.4%) were significantly more likely than black (76.5%) or Hispanic (79.2%) students to have seen or heard such commercials during this time frame.

Among the 23 states that asked this question, the percentage of students who saw antismoking commercials on television or heard antismoking commercials on the radio during the 30 days preceding the survey ranged from 72.2% in Maryland to 86.1% in Mississippi (private schools) (median: 78.9%) (Table 27).

High School. Nationally, 84.2% of students saw antismoking commercials on television or heard antismoking commercials on the radio during the 30 days preceding the survey (Figure 12). White (86.1%) and Asian (87.5%) students were significantly more likely than black (78.2%) or Hispanic (81.5%) students to have seen or heard such commercials (Table 26) during this time frame.

FIGURE 12. Percentage of all middle school and high school students who were influenced by media and advertising regarding tobacco — National Youth Tobacco Survey, 2000



Among the 21 states that asked this question, the percentage of students who saw antismoking commercials on television or heard antismoking commercials on the radio during the 30 days preceding the survey ranged from 76.3% in Ohio to 87.3% in Mississippi (private schools) and Wisconsin (median: 84.6%) (Table 27).

Saw Actors Smoking on Television or in Movies

Middle School. Nationally, 83.3% of students had seen actors using tobacco on television or in the movies (Figure 12). White students (84.4%) were significantly more likely than black (81.0%) or Hispanic (81.5%) students to have seen actors smoking on television or in the movies (Table 26).

Among the 19 states that asked this question, the percentage of students who had seen actors smoking on television or in the movies ranged from 79.3% in Connecticut to 89.7% in Kentucky and West Virginia (median: 84.1%) (Table 27).

High School. Nationally, 88.0% of students had seen actors smoking on television or in the movies (Figure 12). Female students (89.7%) were significantly more likely than male students (86.4%) to have seen actors smoking on television or in the movies (Table 26). White students (89.2%) were significantly more likely than black (84.6%) or Hispanic (86.2%) students to have seen actors smoking on television or in the movies.

Among the 19 states that asked this question, the percentage of students who had seen actors smoking on television or in the movies ranged from 86.1% in the District of Columbia to 92.1% in Iowa (median: 89.8%) (Table 27).

Saw Advertisements for Tobacco Products on Internet

Middle School. Nationally, 32.4% of students had seen advertisements for tobacco products on the Internet (Figure 12). Black students (35.0%) were significantly more likely than Asian students (28.6%) to have seen advertisements for tobacco products on the Internet (Table 26).

Among the 22 states that asked this question, the percentage of students who had seen such advertisements on the Internet ranged from 27.4% in Connecticut to 40.8% in Alabama (median: 33.9%) (Table 27).

High School. Nationally, 24.4% of students had seen advertisements for tobacco products on the Internet (Figure 12). Black (27.8%) and Hispanic (30.6%) students were significantly more likely than white students (22.5%) to have seen advertisements for tobacco products on the Internet (Table 26).


Among the 22 states that asked this question, the percentage of students who had seen such advertisements on the Internet ranged from 20.0% in Connecticut to 34.9% in Mississippi (public schools) (median: 26.3%) (Table 27).

Participation in Antitobacco Community Events

Students were asked whether they had participated in any antitobacco community events to discourage persons from using tobacco products during the preceding 12 months.

Middle School. Nationally, no difference in participation in antitobacco community events was observed between students who had never used tobacco (15.3%) and those who had ever used tobacco (15.8%) (Figure 13). Among students who had never used tobacco, black students (18.5%) were significantly more likely than Hispanic (13.3%)

FIGURE 13. Percentage of middle school and high school students who participated in antitobacco community events, by tobacco use status — National Youth Tobacco Survey, 2000



*Ever use of cigarettes *or* cigars *or* smokeless tobacco *or* pipes *or* bidis *or* kreteks.

or Asian (12.2%) students to have participated in antitobacco community events (Table 28).

Among the 26 states that asked this question, the percentage of never tobacco users who had participated in any antitobacco community events ranged from 8.9% in Maine to 40.2% in Arkansas (median: 26.4%). For students who had ever used tobacco, rates ranged from 11.9% in New York to 32.3% in the District of Columbia (median: 21.0%) (Table 29).

High School. Nationally, no difference in participation in antitobacco community events was observed between students who had never used tobacco (10.8%) and those who had ever used tobacco (9.3%) (Figure 13). Among students who had never used tobacco, female students (12.6%) were significantly more likely than male students (8.8%) to have participated in antitobacco community events (Table 28). Among students who had ever used tobacco, black (13.2%), Hispanic (11.6%), and Asian (12.6%) students were significantly more likely than white students (7.6%) to have participated in antitobacco community events.

Among the 24 states that asked this question, the percentage of never tobacco users who had participated in any antitobacco community events ranged from 9.0% in Texas to 33.7% in Arkansas (median: 19.0%). For students who had ever used tobacco, rates ranged from 8.9% in Indiana and Mississippi (private schools) to 25.7% in Hawaii (median: 14.5%) (Table 29).


Receptivity to Tobacco Company Merchandise

Bought or Received Anything with Tobacco Company Name or Picture on It

Middle School. Nationally, current tobacco users (45.5%) were significantly more likely than students who had never used tobacco (11.4%) to have bought or received anything with a tobacco company name or picture on it (e.g., sports gear, T-shirt, cigarette lighter, hat, jacket, or sunglasses that they purchased or received for free) (Figure 14). Among current tobacco users, white (50.4%) and Hispanic (45.1%) students were significantly more likely than black students (29.0%) to have bought or received anything with a tobacco company name or picture on it (Table 30).

Among the 26 states that asked this question, the percentage of never tobacco users who had bought or received anything with a tobacco company name or picture on it ranged from 8.4% in the District of Columbia to 20.5% in Maine (median: 12.9%). Among current tobacco users, rates ranged from 29.4% in the District of Columbia to 58.4% in West Virginia (median: 45.2%) (Table 31).

FIGURE 14. Percentage of middle school and high school students receptive to tobacco company merchandise*, by tobacco use status — National Youth Tobacco Survey, 2000



* For example, a cigarette lighter or T-shirt.

[†] Used cigarettes or cigars or smokeless tobacco or pipes or bidis or kreteks on ≥ 1 of the 30 days preceding the survey.

High School. Nationally, current tobacco users (34.6%) were significantly more likely than students who had never used tobacco (11.1%) to have bought or received anything with a tobacco company name or picture on it (Figure 14). Hispanic students (13.0%) were significantly more likely than black students (8.8%) to have bought or received anything with a tobacco company name or picture on it. Among current tobacco users, male students (37.3%) were significantly more likely than female students (31.0%) to have bought or received anything with a tobacco company name or picture on it (Table 30).

Among the 24 states that asked this question, the percentage of never tobacco users who had bought or received anything with a tobacco company name or picture on it ranged from 9.6% in Delaware to 17.9% in West Virginia (median: 12.3%). Among current tobacco users, rates ranged from 21.4% in the District of Columbia to 46.9% in Mississippi (private schools) (median: 38.3%) (Table 31).

Would Wear or Use Something with Tobacco Company Name or Picture on It

Middle School. Nationally, current tobacco users (57.3%) were significantly more likely than students who had never used tobacco (10.8%) to report they would wear or use anything with a tobacco company name or picture on it (e.g., cigarette lighter or T-shirt) (Figure 14). Among students who had never used tobacco, male students (15.3%) were significantly more likely than female students (7.0%) to report they would wear or use anything with a tobacco company name or picture on it (Table 30). Hispanic students (14.1%) were significantly more likely than white students (10.0%) to report they would wear or use anything with a tobacco company name or picture on it. Among current tobacco users, male students (61.7%) were significantly more likely than female students (51.3%) to report they would wear or use anything with a tobacco company name or picture on it. White (62.6%) and Hispanic (54.4%) students were significantly more likely than black students (41.4%) to report they would wear or use anything with a tobacco company name or picture on it.

Among the 26 states that asked students if they would wear or use something with a tobacco company name or picture on it, the percentage of never tobacco users ranged from 8.1% in Colorado to 18.7% in Arkansas (median: 13.1%). Among current tobacco users, rates ranged from 36.5% in the District of Columbia to 71.1% in West Virginia (median: 58.7%) (Table 31).

High School. Nationally, current tobacco users (58.9%) were significantly more likely than students who had never used tobacco (16.0%) to report that they would wear or use anything with a tobacco company name or picture on it (Figure 14). Among never tobacco users, male students (20.6%) were significantly more likely than female students (12.1%) to report that they would wear or use anything with a tobacco company name or picture on it. Hispanic students (19.0%) were significantly more likely than Asian students (12.7%) to report they would wear or use such items (Table 30). Among current tobacco users, male students (62.4%) were significantly more likely than female students (54.2%) to report that they would wear or use something with a tobacco company name or picture on it. White students (63.4%) were significantly more likely than black (40.3%), Hispanic (53.4%), or Asian (42.8%) students to report that they would wear or use such items. Hispanic students were significantly more likely than black students to report that they would wear or use such items.

Among the 24 states that asked this question, the percentage of never tobacco users who would wear or use anything with a tobacco company name or picture on it ranged from 13.5% in Colorado to 24.7% in Mississippi (private schools) (median: 17.4%). Among current tobacco users, rates ranged from 39.5% in the District of Columbia to 70.3% in Wisconsin (median: 60.7%) (Table 31).


Cessation

Cessation Attempts and Desire to Stop

Tried to Quit Smoking Cigarettes during 12 Months Preceding Survey

Middle School. Nationally, 59.9% of current cigarette smokers had tried to quit smoking cigarettes during the 12 months preceding the survey (Figure 15). Female students (65.6%) were significantly more likely than male students (54.7%) to have tried to quit smoking cigarettes in the 12 months preceding the survey (Table 32).

FIGURE 15. Percentage of current cigarette smokers* in middle school and high school who tried to quit and who want to stop smoking — National Youth Tobacco Survey, 2000



* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Among the 26 states that asked this question, the percentage of current cigarette smokers who had tried to quit smoking cigarettes in the 12 months preceding the survey ranged from 46.4% in Vermont to 63.7% in Maryland (median: 56.7%) (Table 33).

High School. Nationally, 59.3% of current cigarette smokers had tried to quit smoking cigarettes during the 12 months preceding the survey (Figure 15), with female students (63.3%) significantly more likely than male students (55.2%) to have tried to quit smoking cigarettes during the 12 months preceding the survey (Table 32). White (78.1%), black (61.5%), and Asian (70.5%) students were significantly more likely than Hispanic (67.9%) students to have tried to quit smoking cigarettes during the 12 months preceding the survey. Asian students were significantly more likely than white students to have tried to quit smoking cigarettes during the 12 months preceding the survey.

Among the 24 states that asked this question, the percentage of current cigarette smokers who had tried to quit cigarette smoking in the 12 months preceding the survey ranged from 51.3% in Mississippi (private schools) to 76.3% in New York (median: 59.2%) (Table 33).

Wants to Stop Smoking Cigarettes

Middle School. Nationally, 55.0% of current cigarette smokers said they want to stop smoking cigarettes (Figure 15 and Table 32).

Among the 26 states that asked this question, the percentage of current cigarette smokers who want to stop smoking cigarettes ranged from 41.1% in Vermont to 65.1% in the District of Columbia (median: 55.2%) (Table 33).

High School. Nationally, 61.0% of current cigarette smokers said that they want to stop smoking cigarettes (Figure 15). White (78.1%) and Asian (73.5%) students were significantly more likely than Hispanic students (57.4%) to report that they want to stop smoking cigarettes (Table 32).

Among the 24 states that asked this question, the percentage of current cigarette smokers who want to stop smoking cigarettes ranged from 49.9% in Kansas to 71.8% in Hawaii (median: 57.1%) (Table 33).


Secondhand Smoke

Exposure to Tobacco Smoke

Were in Same Room with Someone Who Was Smoking on ≥ 1 of Preceding 7 Days

Middle School. Nationally, current cigarette smokers (88.4%) were significantly more likely than never cigarette smokers (48.9%) to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Figure 16). Among never cigarette smokers, female students (51.4%) were significantly more likely than male students (46.1%) to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Table 34). White students (52.8%) were significantly more likely than black (45.8%), Hispanic (36.8%), or Asian (33.5%) students to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days. Black students were significantly more likely than Hispanic or Asian students to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days. Among current cigarette smokers, white students (92.3%)

FIGURE 16. Percentage of middle school and high school students who were in a room or who rode in a car with someone who was smoking cigarettes on ≥ 1 of preceding 7 days and who think smoke from other persons' cigarettes is harmful, by smoking status — National Youth Tobacco Survey, 2000



* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

were significantly more likely than black (80.2%) or Hispanic (84.6%) students to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days.

Among the 26 states that asked this question, the percentage of never cigarette smokers who were in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days ranged from 36.4% in California to 63.7% in West Virginia (median: 49.1%). Among current smokers, rates ranged from 77.1% in the District of Columbia and Maryland to 94.0% in West Virginia (median: 87.1%) (Table 35).

High School. Nationally, current cigarette smokers (91.6%) were significantly more likely than never cigarette smokers (56.2%) to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Figure 16). Among never cigarette smokers, white students (59.2%) were significantly more likely than Hispanic (44.6%) or Asian (49.9%) students to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Table 34). Among current cigarette smokers, female students (92.9%) were significantly more likely than male students (90.3%) to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days. White students (93.4%) were significantly more likely than black (85.1%), Hispanic (86.0%), or Asian (84.9%) students to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days.

Among the 24 states that asked this question, the percentage of never cigarette smokers who were in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days ranged from 39.7% in the District of Columbia to 71.5% in Kentucky (median: 56.5%). Among current cigarette smokers, rates ranged from 81.2% in the District of Columbia to 95.0% in Wisconsin (median: 91.3%) (Table 35).

Rode in Car with Someone Who Was Smoking on ≥ 1 of Preceding 7 Days

Middle School. Nationally, current cigarette smokers (80.6%) were significantly more likely than never cigarette smokers (31.9%) to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Figure 16). Among students who have never smoked cigarettes, female students (34.1%) were significantly more likely than male students (29.5%) to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Table 34). White (33.3%) and black (33.3%) students were significantly more likely than Hispanic (25.4%) or Asian (19.3%) students to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days. Among current cigarette smokers, female students (84.5%) were significantly more likely than male students (77.0%) to have ridden in a car with someone who was smoking cigarettes. White students (83.6%) were significantly more likely than black (74.5%) or Hispanic (74.0%) students to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days.

Among the 26 states that asked students if they rode in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days, the percentage of never cigarette smokers ranged from 21.7% in the District of Columbia to 45.5% in West Virginia (median: 34.3%). Among current cigarette smokers, rates ranged from 63.0% in New York to 87.1% in Maine (median: 79.1%) (Table 35).

High School. Nationally, current cigarette smokers (82.9%) were significantly more likely than never cigarette smokers (29.6%) to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Figure 16). Among students who have never smoked cigarettes, black students (34.0%) were significantly more likely than Hispanic (25.0%) or Asian (24.3%) students to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Table 34). Among current cigarette smokers, female students (85.8%) were significantly more likely than male students (80.2%) to have ridden in a car with someone who was smoking cigarettes; and white students (84.9%) were significantly more likely than black (77.7%), Hispanic (74.6%), or Asian (75.4%) students to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days.

Among the 24 states that asked this question, the percentage of never cigarette smokers who had ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days ranged from 19.9% in Colorado to 43.3% in Kentucky (median: 29.2%). Among current cigarette smokers, rates ranged from 70.0% in the District of Columbia to 91.3% in Wisconsin (median: 86.1%) (Table 35).

Think Smoke From Other Persons' Cigarettes is Harmful to Them

Middle School. Nationally, never cigarette smokers (92.9%) were significantly more likely than current cigarette smokers (80.0%) to think that secondhand smoke was harmful to them (Figure 16). Among never cigarette smokers, white (95.6%) and Asian (94.6%) students were significantly more likely than black (85.3%) or Hispanic (87.3%) students to think smoke from other persons' cigarettes was harmful to them (Table 34).

Among the 26 states that asked this question, the percentage of never cigarette smokers who thought that smoke from other persons' cigarettes was harmful to them ranged from 82.7% in the District of Columbia to 97.5% in Maine (median: 91.9%). Among current cigarette smokers, rates ranged from 67.6% in the District of Columbia to 88.5% in New Hampshire (median: 82.7%) (Table 35).

High School. Nationally, never cigarette smokers (94.7%) were significantly more likely than current cigarette smokers (89.8%) to think that smoke from other persons' cigarettes was harmful to them (Figure 16). Among never cigarette smokers, female students (96.2%) were significantly more likely than male students (93.0%) to think smoke from other persons' cigarettes was harmful to them; white (96.6%) and Asian (95.6%) students were significantly more likely than black (87.3%) or Hispanic (90.8%) students to think that smoke from other persons' cigarettes was harmful to them. Among current cigarette smokers, female students (92.9%) were significantly more likely than male students (86.9%) to think smoke from other persons' cigarettes was harmful to them; white students (91.9%) were significantly more likely than black (79.7%) or Hispanic (85.4%) students, and Asian students (89.4%) were significantly more likely than black students to think that smoke from other persons' cigarettes was harmful to them (Table 34).

Among the 24 states that asked students if they thought that smoke from other persons' cigarettes is harmful to them, the percentage of never cigarette smokers ranged from 87.1% in the District of Columbia to 97.2% in Delaware and Iowa (median: 94.9%). Among current smokers, rates ranged from 76.2% in the District of Columbia to 93.8% in Wisconsin (median: 89.9%) (Table 35).

Exposure to Tobacco Use at Home

Anyone in Home Smokes Cigarettes


Middle School. Nationally, current cigarette smokers (70.1%) were significantly more likely than students who had never smoked cigarettes (32.4%) to live in a home where someone smokes cigarettes (Figure 17). Black never cigarette smokers (36.5%) were significantly more likely than Hispanic never cigarette smokers (28.7%) to live in a home where someone smokes cigarettes (Table 36). Among current cigarette smokers, white students (73.8%) were significantly more likely than black (63.7%) or Hispanic (62.0%) students to live in a home where someone smokes cigarettes.

Among the 26 states that asked this question, the percentage of never cigarette smokers who lived in a home where someone smoked cigarettes ranged from 25.2% in Wyoming to 43.8% in Arkansas (median: 35.4%). Among current cigarette smokers, rates ranged from 57.1% in California to 77.5% in Maine (median: 68.9%) (Table 37).

High School. Nationally, current cigarette smokers (56.7%) were significantly more likely than students who had never smoked cigarettes (27.5%) to live in a home where someone smoked (Figure 17 and Table 36).

Among the 24 states that asked this question, the percentage of never cigarette smokers ranged from 20.8% in Colorado to 40.5% in Mississippi (public schools) (median: 28.7%). Among current cigarette smokers, rates ranged from 43.0% in Mississippi (private schools) to 64.9% in Delaware (median: 56.7%) (Table 37).

FIGURE 17. Percentage of middle school and high school students who were exposed to tobacco use at home, by tobacco status — National Youth Tobacco Survey, 2000



* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

† Used smokeless tobacco on ≥1 of the 30 days preceding the survey.

Anyone in Home Uses Smokeless Tobacco

Middle School. Nationally, current users of smokeless tobacco (47.7%) were significantly more likely than students who had never used smokeless tobacco (7.4%) to live in a home where someone used smokeless tobacco (Figure 17). Among never users, white students (8.8%) were significantly more likely than black (5.2%), Hispanic (3.6%), or Asian (3.2%) students to live in a home where someone used smokeless tobacco (Table 36).

Among the 26 states that asked this question, the percentage of never users ranged from 3.7% in California and New York to 23.9% in West Virginia (median: 9.8%). Among current users, rates ranged from 29.1% in Delaware to 61.3% in Arkansas (median: 45.4%) (Table 37).

High School. Nationally, current users of smokeless tobacco (31.8%) were significantly more likely than students who had never used smokeless tobacco (6.7%) to live in a home where someone used smokeless tobacco (Figure 17). Among never users, female students (7.8%) were significantly more likely than male students (5.2%) to live in a home where someone used smokeless tobacco, and white students (7.7%) were significantly more likely than Hispanic (3.9%) or Asian (3.4%) students to live in a home where someone used smokeless tobacco (Table 36).

Among the 24 states that asked this question, the percentage of never users who lived in a home where someone used smokeless tobacco ranged from 3.9% in Delaware to 22.7% in West Virginia (median: 9.3%). Among current users, rates ranged from 21.5% in Hawaii to 50.5% in Mississippi (private schools) (median: 38.1%) (Table 37).

School

Students Who Practiced Ways to Say "No" to Tobacco as Part of School Curriculum

Practice of Refusal Skills


Middle School. Nationally, 41.9% of students practiced ways to say "no" to cigarettes as part of the school curriculum (Figure 18 and Table 38).

Among the 25 states that asked this question, the percentage of students who practiced ways to say "no" to tobacco in school ranged from 22.1% in Mississippi (private schools) to 54.9% in Maine (median: 42.0%) (Table 39).

High School. Nationally, 14.7% of students practiced ways to say "no" to cigarettes as part of the school curriculum (Figure 18). Black (21.0%), Hispanic (18.5%), and Asian (18.6%) students were significantly more likely than white students (12.4%) to have practiced ways to say "no" to tobacco in school (Table 38).

Among the 23 states that asked this question, the percentage of students who practiced ways to say "no" to tobacco in school ranged from 9.3% in Mississippi (private schools) to 23.7% in Hawaii (median: 15.0%) (Table 39).

FIGURE 18. Percentage of all middle school and high school students who practiced ways to say “no” to tobacco as part of school curriculum and who smoked cigarettes or used smokeless tobacco on school property during the 30 days preceding the survey — National Youth Tobacco Survey, 2000



Students Who Smoked Cigarettes on School Property During Preceding 30 Days

Cigarettes

Middle School. Nationally, 3.0% of students smoked cigarettes on school property in the 30 days preceding the survey (Figure 18). Male students (3.6%) were significantly more likely than female students (2.4%) to have smoked cigarettes on school property in the 30 days preceding the survey (Table 38).

Among the 25 states that asked this question, the percentage of students who smoked cigarettes on school property during the 30 days preceding the survey ranged from 1.7% in Kansas to 6.3% in Kentucky (median: 3.9%) (Table 39).

High School. Nationally, 9.4% of students smoked cigarettes on school property during the 30 days preceding the survey (Figure 18). Male students (11.1%) were significantly more likely than female students (7.7%) to have smoked cigarettes on school

property during the 30 days preceding the survey, and white students (10.4%) were significantly more likely than black students (5.7%) to have smoked cigarettes on school property during the preceding 30 days (Table 38).

Among the 24 states that asked this question, the percentage of students who had smoked cigarettes on school property during the 30 days preceding the survey ranged from 6.4% in California to 18.6% in Kentucky (median: 10.7%) (Table 39).

Students Who Used Smokeless Tobacco on School Property During Preceding 30 Days

Smokeless Tobacco

Middle School. Nationally, 2.0% of students used smokeless tobacco on school property during the 30 days preceding the survey (Figure 18). Male students (3.3%) were significantly more likely than female students (0.8%) to have used smokeless tobacco on school property during the 30 days preceding the survey (Table 38).

Among the 26 states that asked this question, the percentage of students who had used smokeless tobacco on school property during the 30 days preceding the survey ranged from 1.1% in New Hampshire to 5.1% in Wyoming (median: 2.1%) (Table 39).

High School. Nationally, 4.0% of students used smokeless tobacco on school property in the 30 days preceding the survey (Figure 18). Male students (7.3%) were significantly more likely than female students (0.6%) to have used smokeless tobacco on school property during the 30 days preceding the survey, and white students (4.5%) were significantly more likely than black (2.6%) or Asian (1.8%) students to have used smokeless tobacco on school property during the 30 days preceding the survey (Table 38).

Among the 24 states that asked this question, the percentage of students who had used smokeless tobacco on school property during the 30 days preceding the survey ranged from 2.3% in Delaware to 11.9% in Mississippi (private schools) (median: 5.1%) (Table 39).

DISCUSSION

CDC recommends that states establish comprehensive tobacco-control programs that include nine elements (9):

- **community programs to reduce tobacco use***;
- chronic disease programs to reduce the burden of tobacco-related disease;
- **school programs**;
- **enforcement**;
- statewide programs;
- **countermarketing**;
- **smoking cessation programs**;
- **surveillance and evaluation**; and

*Bold, italic type indicates the six elements of Best Practices for Tobacco Control Programs (9) that are covered in the state Youth Tobacco Surveys. States can use survey data to monitor these six elements.

- administration and management.

Survey data can be used as part of an essential surveillance and evaluation system to document the extent of the problem and monitor and document the effectiveness of comprehensive tobacco-control programs. Surveillance should include periodic and ongoing monitoring of tobacco-related attitudes, behaviors, health outcomes, and the prevalence of protobacco influences (e.g., advertising, promotions, and events that glamorize tobacco use). Evaluation elements should measure the affect of program elements on attitudes, behaviors, and policies.

Linking YTS Data to Program Components

States can use this report in developing and reporting on six of the nine elements of their comprehensive tobacco-control programs. National estimates for these elements provide an initial marker by which states can compare their current data. In the future, states can a) monitor their trends across time, a step that could lead to program development and modifications; b) compare their state data with those of other states to identify successful programs; and c) compare their state data with national estimates.

The following data from the NYTS are an example of how states can report youth surveillance and evaluation data within the context of a comprehensive tobacco-control program.

Surveillance

The rate of current cigarette smoking documented by the 2000 NYTS is consistent with the most recent data from the national Youth Risk Behavior Survey (13) and the Monitoring the Future Survey (4), both conducted in 1999. For the NYTS and the Monitoring the Future Survey, rates are similar for students in grades 8, 10, and 12. For the NYTS and the national Youth Risk Behavior Survey, rates are similar for students in grades 9, 10, 11, and 12.

Community Programs

Community programs to reduce tobacco use should focus on four goals: a) prevention of the initiation of tobacco use among young persons; b) cessation for current users of tobacco; c) protection from secondhand smoke; and d) elimination of disparities in tobacco use among populations. CDC encourages states to analyze and report their data regarding racial and ethnic disparities when possible. NYTS results support the need for these goals.

- Young persons have a high exposure to secondhand smoke
 - During the previous week, almost nine out of 10 current cigarette smokers and approximately one half of never cigarette smokers were in the same room with someone who was smoking.
 - During the previous week, eight out of 10 current cigarette smokers and three out of 10 never cigarette smokers rode in a car with someone who was smoking cigarettes.
 - Seven out of 10 middle school students who currently smoke cigarettes and six out of 10 high school students who currently smoke cigarettes live in a

home where someone smokes cigarettes. This is compared with three out of 10 never cigarette smokers in both middle school and high school who live in a home where someone smokes cigarettes.

- Approximately 90% of young persons think smoke from other people's cigarettes is harmful to them.
- Only 16% of middle school students and 9% of high school students who have ever used tobacco participated in a community event to discourage persons from using tobacco.
- Among high school students, black students have significantly lower rates of current cigarette smoking than white and Hispanic students — a finding consistent with reports from previous national surveys (3,4,10). In middle school, however, NYTS data document that rates of current cigarette smoking among black students do not differ significantly from rates among white and Hispanic students.

School Programs

School programs that prevent the initiation of tobacco use are a critical element of comprehensive tobacco-control programs because the majority of persons start smoking before age 18 years (5,14). Several studies have documented that effective school-based tobacco prevention programs, in conjunction with community intervention programs such as media campaigns and price increases, can significantly reduce or delay adolescent smoking (5,14–17).

- Inclusion of tobacco refusal skills as part of school curricula is needed as well as enforcement of smoke-free school policies.
 - Only 42% of middle school and 15% of high school students were taught ways to say “no” to tobacco as part of their curriculum in school during the preceding year.
 - A total of 3% of middle school and approximately 9% of high school students smoked cigarettes on school property during the preceding 30 days.

Enforcement

Enforcement of tobacco-control policies enhances their efficacy by deterring violators and by sending a message to the public that community leaders believe that these policies are important (18). The two primary areas addressed by local and state policies that require enforcement strategies are restrictions on minors' access to tobacco and restrictions on smoking (clean indoor air) policies.

- In the majority of tobacco-control policies, restrictions on minors' access are not enforced.
 - Approximately 70% of middle school and 60% of high school students who currently smoke cigarettes and are age <18 years were not asked to show proof of age when they purchased cigarettes.
 - Approximately 60% of middle school and high school students who currently smoke cigarettes and are age <18 years were not refused purchase of cigarettes because of age.

Countermarketing

Tobacco advertising and promotion activities can both stimulate adult consumption and increase the risk of youth initiation (5,19). Children buy the most heavily advertised cigarette brands (20) and are three times more affected by advertising than are adults (21). Countermarketing activities can promote cigarette smoking cessation among young persons and decrease the likelihood that they will begin cigarette smoking. Countermarketing messages can also substantially influence public support for tobacco-control intervention and build a supportive climate for school and community efforts.

- A high percentage of young persons are exposed to tobacco advertising and promotion activities.
 - Approximately 85% of middle school and high school students saw actors using tobacco on television or in movies.
 - Approximately one third of middle school students and one fourth of high school students saw advertisements for tobacco on the Internet.
 - One out of 10 middle school and high school students who have never used tobacco bought or received something with a tobacco company name or picture on it. The rate more than triples for current tobacco users.
 - Approximately 15% of middle school and high school students who have never used tobacco would wear or use something with a tobacco company name or picture on it. The rate increases to nearly 60% among students who currently use tobacco.
- Young persons who smoke cigarettes have strong brand preferences.
 - Approximately one half of middle school and high school students who currently smoke cigarettes report that they usually smoke Marlboro® cigarettes.
 - Black students are more likely to smoke Newport® cigarettes than any other brand.
- Exposure to countermarketing messages is high. Approximately eight out of 10 middle school and high school students saw an antismoking commercial on television or heard one on the radio.

Smoking Cessation

Quitting cigarette smoking can produce a increased short-term public health benefit than any other element of a comprehensive tobacco-control program. The NYTS and state YTS data demonstrate the need for programs that help adolescents to quit cigarette smoking. However, little is known about the effectiveness of these interventions among adolescents (22). With no interventions proven effective for adolescents, states should carefully evaluate any programs that are implemented.

- Effective youth cessation programs are needed.
 - Approximately 60% of middle school and high school students who are current cigarette smokers tried to quit during the preceding 12 months.

- More than one half of middle school and high school students who are current cigarette smokers reported that they want to stop smoking cigarettes.

Limitations

The findings in this report have at least three limitations. First, these data apply only to youth who attend middle school or high school and are not representative of all persons in this age group. However, in 1997, only 4% of 16-year-olds and 6% of 17-year-olds who had not completed high school were not enrolled in a high school program (23). The dropout rate for young adults aged 16–24 years varies substantially by race/ethnicity (7.6%, white; 13.4%, black; and 25.3% Hispanic) (23). Second, the NYTS and the majority of state surveys were conducted during the spring semester, whereas a small number of states conducted surveys during the fall semester. Within each grade, the fall school population is approximately 6 months younger than the spring school population. This difference can be expected to lead to higher estimates of ever tobacco use in the spring surveys and also might lead to higher estimates of current use. Future analysis of data from the fall 1999 NYTS and spring 2000 NYTS will provide a measure of this effect. Third, the data are all based on self-reports, possibly leading to under- or overreporting of behavior. Although the extent of this under- or overreporting of behavior cannot be determined, some YTS questions have been analyzed and demonstrate good test-retest reliability (24).

CONCLUSION

State Implementation of YTS

After completing their 2000 surveys, Florida, Mississippi, and Texas have data for 3 years, and Arkansas, Kansas, and Tennessee have data for 2 years to begin examining trends. A report recently published by Florida documents a substantial decline in youth tobacco use — including current and lifetime use of cigarettes, cigars, and smokeless tobacco — during the preceding 3 years (25). Substantial shifts in attitudes toward an antitobacco direction also occurred in Florida during this time. YTS findings in Florida have led to the adoption of youth-oriented programs by other states.

The YTS provides states with a mechanism for responding to emerging concerns within tobacco control. For example, in 1998, Texas was the first state to include questions regarding tobacco use in a pipe. After Texas reported a 8.5% prevalence rate of pipe use among middle school students, other states realized the public health significance of monitoring pipe use among adolescents. Similarly, in 1999, New Jersey was the first state to include questions regarding use of bidis and kreteks. After finding 11.0% of middle school students and 21.3% of high school students had ever smoked bidis, and 6.3% of middle school students and 11.1% of high school students had ever smoked kreteks, questions regarding these products were added to the 2000 core YTS questionnaire.

Data from the YTS have been used in several states to generate legislative support for allocation of dollars from the Master Settlement Agreement to support tobacco control and prevention efforts. The survey can then be used to report on the effectiveness of the programs developed under the Master Settlement Agreement.

A distinct feature of the YTS is that it allows states to obtain substate-level estimates to support their tobacco-control efforts. For example, after conducting the survey in 1999 to provide baseline data to support the youth component of their tobacco-control program, Tennessee expanded their 2000 YTS sample design to provide data for each of the state's 13 health regions to support local tobacco-control efforts. The scope of such efforts continues to grow, and during fall 2000, Maryland conducted the largest YTS to date, successfully collecting middle school and high school data in 23 counties and the city of Baltimore.

Localities (e.g., city, community, county, district, tribal councils, and U.S. territories) have also conducted the survey independent of the state(s). For example, separate surveys have been conducted in large cities such as New York City and Detroit; in seven smaller cities and counties in Wisconsin (Cameron, Florence County, Franklin County, Neenah, St. Francis, Western Racine County, and Waupaca); by two tribal councils (Great Lakes Inter-Tribal Council and Inter-Tribal Council of Arizona); and by four U.S. territories (American Virgin Islands, Commonwealth of the Northern Mariana Islands, Federated States of Micronesia, and Palau).

Finally, the YTS is an integral component of surveillance and evaluation plans implemented by states. New Jersey has developed a logic model to outline the links between multiple data sources and their programmatic objectives. In New Jersey, the YTS is one source of data for monitoring progress toward initial, intermediate, and long-term outcome objectives for their comprehensive tobacco-control program. The YTS continues to serve as a critical surveillance and evaluation tool for the design, implementation, and evaluation of comprehensive tobacco-control and prevention programs.

Uses of YTS Data

For states that have participated in the survey and for states that plan to participate in the future, the state YTS and NYTS are primary sources of data for planning, monitoring and evaluating the youth component of state-based tobacco-control programs. The following are examples of how data can be used:

- Prevalence estimates from the YTS identify where a state stands in comparison with Healthy People 2010 objectives. As programs mature, repeating the YTS can indicate whether or not the state has moved toward meeting their objectives. In addition, these estimates highlight the greatest problems (e.g., rate of cigarette smoking is much higher than cigar smoking) and aid in focusing programmatic efforts.
- Data collected regarding knowledge and attitudes identify key perceptions concerning social context and influence and harm from tobacco. This information can be used to direct programs that focus on topics regarding peer groups and awareness of the dangers of tobacco use.
- General questions regarding exposure to tobacco use in the media can be used to monitor the degree of exposure to pro- or antitobacco messages and gauge social context. State-specific questions regarding a particular advertisement campaign can help determine the reach and duration of these messages.
- Questions regarding minors' access determine where and how youth are obtaining tobacco products. Data can be used to monitor compliance with the

Synar Amendment, which bans the sale of tobacco products to youth aged <18 years. Also, programmatic and policy efforts can target the persons or vendors who are most likely to provide or sell tobacco to underaged youth.

- School curriculum and policy play an important role in reducing youth tobacco use. Related YTS questions identify needs by assessing the extent to which students are being taught antitobacco lessons as part of their curriculum. In addition, the data indicate whether schools have been effective in enforcing rules that ban tobacco use on campus.
- YTS questions regarding exposure to secondhand smoke determine where students are exposed and to what extent. These data are often used to develop key messages for media campaigns that aim to reduce exposure.
- Data regarding cessation attempts indicate the level of desire to quit as well as of addiction. This information assesses the need to provide students with programs that help them to quit using tobacco.

References

1. McGinnis JM, Foege WH. Actual causes of death in the United States. *JAMA* 1993;270:2207–12.
2. CDC. Smoking-attributable mortality and years of potential life lost—United States 1984. *MMWR* 1997;46:444–51.
3. CDC. Tobacco use among high school students—United States, 1997. *MMWR* 1998;47:229–33.
4. Johnston LD, O'Malley PM, Bachman JG. Monitoring the Future Survey. Ann Arbor, MI: University of Michigan, 1999. Available at <http://monitoringthefuture.org/data/99data.html#1999data_cigs>. Accessed August 15, 2000.
5. Department of Health and Human Services. Preventing tobacco use among young people: a report of the Surgeon General. Atlanta, GA: Department of Health and Human Services, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1994. [Reprinted with corrections, July 1994.]
6. Miller VP, Ernst C, Collin F. Smoking-attributable medical care costs in the USA. *Soc Sci Med* 1999;48:375–91.
7. Miller LS, Zhang X, Rice DP, Max W. State estimates of total medical expenditures attributable to cigarette smoking, 1993. *Public Health Rep* 1998;113:447–58.
8. CDC. Cigarette smoking-attributable mortality and years of potential life lost—United States, 1990. *MMWR* 1993;42:645–9.
9. CDC. Best Practices for comprehensive tobacco control programs, August 1999. Atlanta, GA: Department of Health and Human Services, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1999.
10. CDC. Tobacco use among middle and high school students—United States, 1999. *MMWR* 2000;49:49–53.
11. Shah BV, Barnwell BG, Bieler GS. Software for the Statistical Analysis of Correlated Data (SUDAAN): user's manual, release 7.5, 1997 [Software documentation]. Research Triangle Park, NC: Research Triangle Institute, 1997.
12. Pierce JP, Choi WS, Gilpin EA, Farkas AJ, Berry CC. Tobacco industry promotion of cigarettes and adolescent smoking. *JAMA* 1998;279:511–5.
13. Kann L, Kinchen SA, Williams BI, et al. Youth risk behavior surveillance—United States, 1999. In: CDC Surveillance Summaries (June 9). *MMWR* 2000;49(No. SS-5).
14. Institute of Medicine. Growing up tobacco free. Lynch BS, Bonnie RJ, eds. Washington, DC: National Academy Press, 1994:143–74.

15. Glynn TJ. Essential elements of school-based smoking prevention programs. *J Sch Health* 1989;59:181–8.
16. Bruvold WH. A meta-analysis of the California school-based risk reduction program. *J Drug Educ* 1990;20:139–52.
17. Rooney BL, Murray DM. A meta-analysis of smoking prevention programs after adjustment for errors in the unit of analysis. *Health Education Quarterly* 1996;23:48–68.
18. Forster JL, Woldson M. Youth access to tobacco: policies and politics. *Annu Rev Public Health* 1998;19:203–35.
19. US Food and Drug Administration. 21 CFR Parts 801, 803, 804, and 897. Regulations restricting the sale and distribution of cigarettes and smokeless tobacco products to protect children and adolescents; final rule. *Federal Register* 1996;60:155:41314–75.
20. CDC. Changes in the cigarette brand preference of adolescent smokers—United States, 1989–1993. *MMWR* 1994;43:577–81.
21. Pollay RW, Siddarth S, Siegel M, et al. The last straw? Cigarette advertising and realized market shares among youths and adults, 1979–1993. *Journal of Marketing* 1996;60:1–16.
22. Fiore MC, Bailey WC, Cohen SJ, et al. Treating tobacco use and dependence: a clinical practice guideline. Rockville, MD: Department of Health and Human Services, Public Health Service, 2000; publication no. AHRQ 00-0032.
23. US Department of Education. Dropout rates in the United States, 1997. Washington, DC: US Department of Education, National Center for Education Statistics, Office of Educational Research and Improvement, 1999; publication no. NCES 99-082.
24. Brener ND, Collins JL, Kann L, Warren CW, Williams BI. Reliability of the Youth Risk Behavior Survey questionnaire. *Am J Epidemiol* 1995;141:575–80.
25. Bauer U, Johnson T. Assessing the impact of Florida's pilot program on tobacco control, 1998–2000: a comprehensive analysis of data from the Florida Youth Tobacco Survey. Vol. 3, report 2. Tallahassee, FL: Florida Department of Health, 2001.

**TABLE 1. Sample sizes and response rates for middle schools and high schools, by state—
State Youth Tobacco Surveys, 2000**

States	Student completed sample size	Response rate (%)		
		School	Student	Overall
Middle school				
Alabama	1,576	87.8	84.3	73.96
Arizona	2,063	82.0	80.4	65.9
Arkansas	1,582	94.0	70.5	66.2
California	2,910	83.3	92.6	77.1
Colorado	1,506	71.9	86.2	62.0
Connecticut	2,089	70.7	86.4	61.1
Delaware	4,654	97.1	83.2	80.8
District of Columbia	1,205	95.2	81.3	77.4
Florida	14,311	91.4	89.7	82.0
Hawaii	1,045	100.0	54.1	54.1
Indiana	1,516	81.2	89.7	72.9
Iowa	1,834	86.7	87.0	75.4
Kansas	1,609	84.0	83.8	70.4
Kentucky	1,282	74.0	83.3	61.6
Maine	1,370	74.2	83.3	61.8
Maryland	33,586	99.4	84.8	84.3
Minnesota	4,751	79.3	87.9	69.7
Mississippi — Private	1,282	70.0	95.32	66.7
Mississippi — Public	1,864	85.0	91.4	77.7
New Hampshire	1,525	80.0	87.3	69.9
New York	1,996	90.9	94.2	85.6
Ohio	1,728	87.8	84.5	74.2
Tennessee	10,779	96.0	75.5	72.4
Texas	2,112	95.2	96.4	91.8
Vermont	1,501	76.0	88.4	67.2
West Virginia	1,757	86.0	79.5	68.4
Wisconsin	1,440	77.6	88.6	68.7
Wyoming	1,843	80.0	89.1	71.3
High school				
Alabama	1,575	93.8	80.9	75.8
Arizona	1,399	60.0	74.5	47.7
Arkansas	1,394	82.0	76.2	62.5
California	3,029	93.6	92.2	86.3
Colorado	1,552	83.0	86.3	71.6
Connecticut	2,200	82.1	74.3	61.0
Delaware	3,516	92.6	72.2	66.8
District of Columbia	1,359	92.9	64.5	59.9
Florida	9,484	91.2	89.2	81.4
Hawaii	1,511	100.0	61.3	61.3
Indiana	1,416	86.0	85.0	73.1
Iowa	1,445	90.0	74.7	67.2
Kansas	1,612	86.0	80.3	69.0
Kentucky	1,313	81.5	82.1	67.0
Maine	583	56.8	80.6	45.8
Maryland	22,381	100.0	89.4	89.4
Minnesota	7,625	74.0	84.6	62.6
Mississippi — Private	1,284	70.0	91.3	63.9
Mississippi — Public	1,701	82.5	89.3	73.7
Nebraska	3,268	85.4	94.4	80.6
New York	2,161	80.8	90.2	72.9
Ohio	1,280	87.8	78.2	68.6
South Dakota	1,529	95.8	84.0	80.5
Tennessee	9,959	89.0	77.4	68.8
Texas	2,293	91.3	93.3	85.2
Vermont	952	60.0	81.9	49.2
West Virginia	1,617	80.0	84.1	67.3
Wisconsin	1,307	88.0	83.5	73.5

TABLE 2. Percentage of all middle school and high school students who ever used* cigarettes, cigars, smokeless tobacco, pipes, bidis, or kreteks, by sex and race/ethnicity—National Youth Tobacco Survey, 2000

	<u>Cigarettes</u>	<u>Cigars</u>	<u>Smokeless tobacco</u>	<u>Pipes</u>	<u>Bidis</u>	<u>Kreteks</u>
	% (95% CI) [†]	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school						
Sex						
Male	38.3 (±2.7)	24.7 (±2.2)	14.8 (±2.7)	9.2 (±1.1)	5.9 (±0.8)	4.6 (±0.7)
Female	34.2 (±2.6)	14.1 (±1.4)	4.2 (±0.7)	4.3 (±0.6)	3.0 (±0.5)	2.6 (±0.5)
Race/ethnicity						
White	33.8 (±2.9)	18.5 (±1.9)	10.7 (±2.2)	6.5 (±0.9)	3.5 (±0.5)	3.5 (±0.6)
Black	44.0 (±3.8)	21.8 (±3.4)	6.1 (±1.5)	4.4 (±0.9)	5.7 (±1.1)	2.7 (±0.7)
Hispanic	40.0 (±3.6)	22.5 (±2.5)	7.5 (±1.2)	9.7 (±1.6)	6.4 (±1.4)	4.4 (±1.0)
Asian	20.8 (±4.0)	9.2 (±2.8)	4.1 (±1.8)	4.7 (±2.0)	4.1 (±2.0)	3.6 (±1.4)
Total (middle school)	36.3 (±2.4)	19.3 (±1.6)	9.5 (±1.5)	6.7 (±0.7)	4.4 (±0.5)	3.6 (±0.5)
High school						
Sex						
Male	65.3 (±1.9)	52.2 (±2.1)	29.0 (±2.6)	16.7 (±1.1)	15.7 (±1.4)	14.0 (±1.3)
Female	62.5 (±2.3)	30.9 (±1.8)	6.8 (±1.0)	5.0 (±0.6)	9.9 (±1.2)	10.2 (±1.3)
Race/ethnicity						
White	64.3 (±2.5)	44.1 (±2.3)	22.1 (±2.0)	12.3 (±0.9)	11.6 (±1.4)	14.4 (±1.5)
Black	62.5 (±2.8)	35.8 (±3.5)	7.6 (±1.5)	5.3 (±1.2)	15.7 (±2.2)	4.2 (±1.2)
Hispanic	66.4 (±2.6)	39.5 (±2.7)	11.0 (±2.0)	10.3 (±1.3)	15.2 (±2.1)	9.6 (±1.6)
Asian	52.4 (±4.3)	24.3 (±3.2)	6.8 (±1.7)	6.3 (±1.9)	10.1 (±2.1)	7.5 (±2.2)
Total (high school)	64.0 (±2.0)	41.7 (±1.8)	18.0 (±1.7)	11.0 (±0.7)	12.9 (±1.2)	12.1 (±1.1)

* Ever use of cigarettes was determined by asking, "Have you ever tried cigarette smoking, even one or two puffs?" Ever use of cigars was determined by asking, "Have you ever tried smoking cigars, cigarillos, or little cigars, even one or two puffs?" Ever use of smokeless tobacco was determined by asking, "Have you ever used chewing tobacco, snuff, or dip, such as Redman,[®] Levi Garrett,[®] Beechnut,[®] Skoal,[®] Skoal Bandits,[®] or Copenhagen?[®]" Ever use of bidis was determined by asking, "Have you ever tried smoking bidis, even one or two puffs?" Ever use of kreteks was determined by asking, "Have you ever tried smoking kreteks, even one or two puffs?"

[†] Confidence interval.

TABLE 3. Percentage of all middle school and high school students who ever used* cigarettes, cigars, smokeless tobacco, bidis, or kreteks, by state—State Youth Tobacco Surveys, 2000

	Cigarettes		Cigars		Smokeless tobacco		Bidis		Kreteks	
	%	(95% CI) [†]	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school										
Alabama	51.2	(±5.0)	29.4	(±2.8)	19.2	(±3.9)	6.3	(±1.6)	4.9	(±1.4)
Arizona	37.7	(±3.6)	21.9	(±2.3)	13.2	(±1.8)	10.1	(±2.5)	5.1	(±2.1)
Arkansas	47.8	(±4.1)	25.3	(±2.7)	19.9	(±3.4)	6.4	(±1.2)	3.9	(±0.9)
California	26.9	(±4.3)	14.7	(±2.8)	5.6	(±1.1)	5.1	(±1.7)	NA [§]	
Colorado	31.9	(±6.3)	20.0	(±4.0)	10.2	(±2.6)	4.2	(±1.1)	3.5	(±1.0)
Connecticut	31.7	(±4.2)	19.6	(±2.0)	7.7	(±1.8)	6.2	(±1.5)	2.7	(±1.1)
Delaware	44.3	(±3.7)	20.0	(±1.5)	7.7	(±1.4)	9.1	(±1.5)	3.2	(±0.6)
District of Columbia	36.3	(±4.0) [¶]	18.4	(±2.5)	8.6	(±1.8)	10.0	(±2.2)	3.3	(±1.1)
Florida	30.7	(±1.7)	16.9	(±1.2)	7.0	(±0.6)	4.6	(±0.5)	NA	
Indiana	34.1	(±6.3)	17.8	(±3.3)	10.4	(±2.5)	5.4	(±1.5)	2.8	(±1.2)
Iowa	39.3	(±5.1)	20.6	(±2.2)	13.8	(±2.7)	6.2	(±1.1)	4.7	(±0.7)
Kansas	28.6	(±5.0)	16.2	(±3.4)	10.0	(±2.4)	5.2	(±1.5)	3.3	(±0.9)
Kentucky	50.0	(±5.3)	29.0	(±3.5)	22.0	(±4.2)	6.5	(±1.9)	4.3	(±1.4)
Maine	38.6	(±5.7)	19.0	(±3.9)	6.6	(±2.3)	4.1	(±1.8)	4.8	(±1.9)
Maryland	26.3	(±2.5)	17.2	(±1.3)	9.7	(±0.8)	7.6	(±0.9)	3.6	(±0.7)
Minnesota	33.3	(±2.1)	18.3	(±1.5)	12.4	(±1.3)	4.7	(±1.2)	2.3	(±0.5)
Mississippi — Private	41.7	(±4.2)	22.5	(±2.4)	22.2	(±3.8)	NA		NA	
Mississippi — Public	53.4	(±3.3)	31.8	(±2.5)	19.7	(±2.7)	NA		NA	
New Hampshire	34.4	(±4.9)	21.2	(±2.1)	7.5	(±1.5)	5.9	(±0.8)	5.0	(±1.0)
New York	29.2	(±6.8)	14.3	(±5.2)	3.8	(±2.4)	4.4	(±1.3)	NA	
Ohio	38.3	(±5.3)	23.8	(±3.8)	13.4	(±3.8)	6.4	(±1.9)	3.3	(±0.9)
Tennessee	45.5	(±2.9)	25.5	(±1.9)	18.4	(±1.7)	7.1	(±0.9)	5.1	(±0.9)
Texas	44.1	(±6.0)	24.2	(±3.2)	10.3	(±1.9)	5.1	(±1.8)	NA	
Vermont	36.2	(±4.3)	20.1	(±2.6)	11.1	(±3.0)	7.0	(±1.8)	2.5	(±0.7)
West Virginia	46.9	(±3.5)	25.2	(±3.1)	21.0	(±1.9)	5.9	(±1.4)	4.1	(±1.0)
Wisconsin	39.4	(±4.6)	22.4	(±2.6)	12.9	(±2.0)	6.6	(±1.8)	3.9	(±1.1)
Wyoming	46.3	(±3.9)	26.3	(±2.9)	24.7	(±2.9)	7.3	(±1.6)	5.1	(±1.3)
Median	38.3		20.6		10.8		6.2		3.9	
High School										
Alabama	69.7	(±2.5)	47.4	(±3.5)	24.7	(±3.4)	7.6	(±1.8)	5.8	(±1.3)
Arkansas	70.5	(±4.6)	50.3	(±3.7)	28.4	(±5.0)	9.4	(±2.3)	6.9	(±1.7)
California	60.5	(±4.3)	37.7	(±3.3)	11.8	(±5.0)	19.3	(±3.4)	NA [§]	
Colorado	61.6	(±5.1)	45.1	(±4.2)	23.5	(±3.5)	16.7	(±3.0)	13.7	(±2.7)
Connecticut	63.0	(±4.7)	43.0	(±4.3)	15.2	(±2.6)	13.3	(±1.9)	4.0	(±1.0)
Delaware	66.4	(±1.9)	37.5	(±2.0)	11.7	(±1.5)	24.0	(±1.9)	8.1	(±1.6)
District of Columbia	58.1	(±3.1)	24.0	(±3.1)	9.9	(±2.0)	19.7	(±3.2)	5.2	(±1.6)
Florida	56.9	(±2.3)	35.6	(±2.3)	12.4	(±1.3)	13.5	(±1.2)	NA	
Hawaii	63.3	(±3.9)	27.5	(±2.9)	10.4	(±2.0)	13.2	(±2.4)	11.0	(±2.5)
Indiana	65.3	(±4.1)	45.4	(±4.5)	20.0	(±4.8)	8.8	(±2.0)	9.1	(±1.8)
Iowa	63.4	(±2.6)	43.5	(±4.0)	27.4	(±3.2)	9.3	(±2.8)	10.6	(±1.9)
Kansas	60.4	(±5.1)	42.8	(±4.6)	25.1	(±3.9)	8.3	(±1.7)	3.8	(±1.0)
Kentucky	74.3	(±4.2)	54.4	(±3.3)	32.8	(±3.7)	9.4	(±1.9)	3.6	(±1.0)
Maryland	57.9	(±1.1)	36.5	(±0.9)	15.2	(±0.7)	16.7	(±0.8)	7.2	(±0.5)
Minnesota	64.7	(±2.1)	45.8	(±2.1)	29.2	(±2.3)	9.4	(±1.4)	9.1	(±1.4)
Mississippi — Private	70.5	(±3.9)	50.7	(±3.0)	38.0	(±4.3)	10.5	(±2.5)	6.5	(±2.0)
Mississippi — Public	71.5	(±2.2)	48.8	(±4.2)	25.0	(±4.2)	10.5	(±1.3)	3.7	(±0.9)
Nebraska	62.4	(±4.1)	42.4	(±2.9)	26.2	(±2.5)	7.0	(±1.3)	5.9	(±1.4)
New York	62.9	(±4.8)	34.4	(±6.8)	14.5	(±5.0)	17.8	(±2.4)	NA	
Ohio	68.7	(±2.4)	52.1	(±3.9)	26.0	(±2.4)	13.3	(±2.5)	9.0	(±2.1)
South Dakota	64.8	(±4.7)	45.3	(±4.0)	30.4	(±4.4)	7.7	(±1.5)	7.0	(±1.9)
Tennessee	69.3	(±1.9)	49.0	(±2.6)	29.9	(±2.4)	12.1	(±1.1)	10.8	(±1.1)
Texas	64.7	(±4.6)	43.1	(±5.0)	19.8	(±6.2)	9.5	(±2.7)	NA	
West Virginia	74.3	(±4.1)	52.2	(±4.0)	31.0	(±3.6)	10.9	(±1.9)	5.2	(±1.0)
Wisconsin	64.9	(±4.5)	47.3	(±3.9)	23.4	(±4.8)	10.5	(±2.3)	10.8	(±1.3)
Median	64.7		45.1		24.7		10.5		7.0	

* Ever use of cigarettes was determined by asking, "Have you ever tried cigarette smoking, even one or two puffs?" Ever use of cigars was determined by asking, "Have you ever tried smoking cigars, cigarillos, or little cigars, even one or two puffs?" Ever use of smokeless tobacco was determined by asking, "Have you ever used chewing tobacco, snuff, or dip, such as Redman,® Levi Garrett,® Beechnut,® Skoal,® Skoal Bandits,® or Copenhagen?®" Ever use of both bidis and kreteks was determined by asking students to choose one of the following: a) I have tried smoking bidis; or b) I have tried smoking kreteks; or c) I have tried smoking both bidis and kreteks; or d) I have never smoked bidis or kreteks."

[†] Confidence interval.

[§] Question was not asked.

[¶] Washington, D.C., middle school sample only included 7th and 8th graders.

TABLE 4. Percentage of all middle school and high school students who were current users of any tobacco product,* cigarettes, cigars, smokeless tobacco, pipes, bidis, or kreteks, by sex and race/ethnicity — National Youth Tobacco Survey, 2000

	<u>Any tobacco*</u>	<u>Cigarettes</u>	<u>Cigars</u>	<u>Smokeless tobacco</u>	<u>Pipes</u>	<u>Bidis</u>	<u>Kreteks</u>
	% (95% CI) [†]	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school							
Sex							
Male	17.6 (±2.2)	11.7 (±1.7)	9.7 (±1.5)	5.7 (±1.8)	4.3 (±0.7)	3.4 (±0.6)	2.7 (±0.5)
Female	12.7 (±1.5)	10.2 (±1.3)	4.6 (±0.8)	1.5 (±0.3)	1.8 (±0.4)	1.4 (±0.3)	1.5 (±0.3)
Race/ethnicity							
White	14.3 (±1.9)	10.8 (±1.6)	6.1 (±1.1)	3.9 (±1.3)	2.7 (±0.5)	1.9 (±0.4)	1.9 (±0.4)
Black	17.5 (±3.0)	11.2 (±2.0)	9.8 (±2.5)	2.4 (±0.7)	2.2 (±0.7)	2.9 (±0.8)	1.7 (±0.5)
Hispanic	16.0 (±2.0)	11.4 (±1.7)	8.8 (±1.4)	2.9 (±0.7)	5.3 (±1.1)	3.6 (±0.9)	2.6 (±0.8)
Asian	7.5 (±2.6)	5.3 (±2.3)	4.1 (±1.9)	1.7 (±1.2)	2.8 (±1.5)	2.9 (±1.7)	2.3 (±1.4)
Total (middle school)	5.1 (±1.5)	11.0 (±1.2)	7.1 (±1.0)	3.6 (±0.9)	3.0 (±0.4)	2.4 (±0.4)	2.1 (±0.4)
High School							
Sex							
Male	39.1 (±2.2)	28.8 (±1.9)	22.0 (±1.5)	11.8 (±1.7)	5.2 (±0.7)	5.4 (±0.6)	5.3 (±0.7)
Female	29.8 (±1.9)	27.3 (±2.0)	7.3 (±0.9)	1.4 (±0.4)	1.4 (±0.3)	2.8 (±0.4)	3.0 (±0.5)
Race/ethnicity							
White	38.0 (±2.3)	31.8 (±2.1)	15.1 (±1.2)	8.2 (±1.2)	3.3 (±0.5)	3.6 (±0.5)	4.5 (±0.6)
Black	26.5 (±3.6)	16.8 (±3.0)	15.3 (±2.9)	2.6 (±0.9)	2.2 (±0.8)	4.9 (±1.0)	2.2 (±0.7)
Hispanic	28.4 (±2.5)	22.6 (±2.4)	13.6 (±1.6)	4.0 (±1.2)	4.2 (±0.9)	5.7 (±1.1)	4.0 (±0.8)
Asian	22.9 (±3.7)	20.6 (±3.5)	7.4 (±2.1)	1.9 (±0.9)	2.5 (±1.1)	3.0 (±1.3)	3.2 (±1.4)
Total (high school)	34.5 (±1.9)	28.0 (±1.7)	14.8 (±1.1)	6.6 (±0.9)	3.3 (±0.4)	4.1 (±0.4)	4.2 (±0.5)

* Use of cigarettes or cigars or smokeless tobacco or pipes or bidis or kreteks on ≥ 1 of the 30 days preceding the survey.

[†] Confidence interval.

TABLE 5. Percentage of all middle school and high school students who were current users of any tobacco product*, cigarettes, cigars, smokeless tobacco, pipes, or bidis, by state—State Youth Tobacco Surveys, 2000

	Any tobacco*	Cigarettes	Cigars	Smokeless tobacco	Pipes	Bidis
	% (95% CI) [†]	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school						
Alabama	26.5 (±3.8)	19.1 (±3.2)	12.4 (±2.0)	8.2 (±2.3)	3.9 (±1.4)	4.0 (±1.1)
Arizona	17.1 (±2.4)	11.4 (±1.4)	6.1 (±1.1)	3.9 (±1.5)	4.3 (±1.4)	4.9 (±1.4)
Arkansas	22.4 (±2.6)	15.8 (±2.1)	8.5 (±1.4)	7.9 (±1.9)	4.3 (±1.2)	4.4 (±1.2)
California	10.0 (±2.3)	6.7 (±1.7)	5.2 (±1.4)	2.1 (±0.6)	4.0 (±0.9)	2.8 (±1.1)
Colorado	13.6 (±3.6)	8.8 (±2.8)	5.3 (±2.0)	2.4 (±0.8)	2.7 (±0.9)	3.2 (±1.2)
Connecticut	13.1 (±3.2)	9.8 (±2.8)	6.1 (±1.6)	2.2 (±0.9)	2.6 (±1.1)	3.7 (±1.2)
Delaware	17.8 (±1.3)	15.2 (±1.6)	8.1 (±0.8)	3.4 (±0.7)	2.9 (±0.7)	NA [§]
District of Columbia	14.2 (±2.5)	9.4 (±2.1)	4.7 (±1.5)	2.3 (±1.0)	3.0 (±1.1)	5.0 (±1.5)
Florida	14.7 (±1.2) [¶]	11.1 (±1.0)	7.5 (±0.8)	3.2 (±0.4)	NA	2.8 (±0.4)
Indiana	15.3 (±3.3)	9.8 (±2.7)	5.2 (±1.4)	4.1 (±1.4)	2.7 (±0.9)	4.4 (±1.0)
Iowa	16.4 (±3.0)	11.8 (±2.7)	5.6 (±1.0)	4.2 (±1.6)	3.6 (±0.6)	3.5 (±0.9)
Kansas	12.0 (±2.7)	8.1 (±2.0)	5.1 (±1.6)	2.7 (±1.2)	1.7 (±0.9)	2.5 (±1.0)
Kentucky	28.3 (±4.4)	21.5 (±3.9)	10.7 (±2.5)	9.4 (±3.3)	4.3 (±1.4)	3.7 (±1.1)
Maine	16.0 (±3.9)	14.5 (±3.9)	6.1 (±2.6)	3.2 (±1.3)	3.1 (±1.5)	2.8 (±1.5)
Maryland	11.8 (±1.5)	7.3 (±1.0)	4.7 (±0.7)	2.2 (±0.4)	2.8 (±0.7)	3.5 (±0.8)
Minnesota	12.6 (±1.9)	9.1 (±1.3)	3.7 (±0.8)	2.2 (±0.5)	2.7 (±0.8)	2.8 (±0.9)
Mississippi — Private	19.9 (±2.2)	14.2 (±2.3)	7.5 (±1.6)	10.1 (±2.4)	NA	NA
Mississippi — Public	25.7 (±2.9)	18.1 (±2.3)	12.1 (±2.2)	7.5 (±2.0)	NA	NA
New Hampshire	15.2 (±2.7)	12.0 (±2.6)	4.3 (±1.0)	2.2 (±0.6)	2.8 (±0.7)	3.0 (±1.0)
New York	11.8 (±4.3)	9.3 (±3.8)	5.0 (±2.5)	1.5 (±1.0)	2.4 (±1.1)	2.2 (±1.0)
Ohio	18.7 (±4.6)	13.7 (±3.5)	8.1 (±2.0)	5.1 (±2.5)	3.5 (±1.0)	3.4 (±1.0)
Tennessee	23.2 (±2.4)	16.6 (±1.9)	9.3 (±1.3)	6.9 (±1.2)	4.3 (±0.7)	4.5 (±0.6)
Texas	17.6 (±3.6)	13.9 (±3.1)	8.5 (±1.8)	3.8 (±1.0)	4.8 (±1.8)	3.4 (±1.3)
Vermont	15.3 (±2.2)	11.9 (±2.1)	4.8 (±1.1)	3.3 (±1.9)	3.4 (±0.9)	3.0 (±0.9)
West Virginia	25.3 (±3.0)	18.1 (±2.5)	8.3 (±1.9)	9.1 (±1.9)	3.7 (±0.9)	5.0 (±0.9)
Wisconsin	16.1 (±3.0)	12.2 (±2.2)	5.9 (±1.6)	3.6 (±1.1)	3.9 (±1.2)	3.5 (±1.5)
Wyoming	21.7 (±2.6)	14.8 (±2.0)	8.1 (±1.4)	8.6 (±1.7)	3.7 (±1.1)	3.5 (±1.0)
Median	16.1	12.0	6.1	3.6	3.4	3.5
High school						
Alabama	37.6 (±3.2)	30.2 (±3.5)	16.6 (±2.2)	10.2 (±2.7)	2.9 (±0.9)	3.4 (±0.9)
Arkansas	43.8 (±4.3)	35.8 (±4.6)	20.0 (±3.0)	11.9 (±2.9)	4.9 (±1.4)	5.2 (±1.3)
California	27.8 (±4.1)	21.6 (±4.0)	11.7 (±1.6)	3.3 (±1.4)	3.4 (±0.7)	5.5 (±1.1)
Colorado	34.4 (±5.3)	25.3 (±5.3)	15.1 (±2.2)	9.0 (±2.9)	4.8 (±1.6)	6.3 (±1.6)
Connecticut	32.4 (±4.7)	25.6 (±4.9)	12.7 (±2.8)	4.1 (±0.9)	4.2 (±1.3)	6.0 (±1.0)
Delaware	31.2 (±1.6)	27.1 (±1.6)	12.9 (±1.4)	3.4 (±0.9)	2.4 (±0.6)	NA [§]
District of Columbia	21.0 (±2.9)	14.7 (±2.5)	7.1 (±1.9)	3.3 (±1.8)	4.2 (±1.8)	6.9 (±2.0)
Florida	29.8 (±2.2) [¶]	22.6 (±1.8)	16.1 (±1.2)	5.4 (±0.7)	NA	5.4 (±0.6)
Hawaii	27.1 (±4.2)	24.5 (±3.8)	4.9 (±1.3)	3.4 (±1.6)	3.7 (±1.7)	5.2 (±2.0)
Indiana	36.9 (±3.7)	31.6 (±3.3)	15.4 (±2.2)	6.9 (±2.2)	3.7 (±1.2)	4.1 (±1.2)
Iowa	39.0 (±2.8)	32.7 (±2.5)	14.5 (±2.7)	10.6 (±2.2)	4.1 (±1.4)	4.1 (±0.9)
Kansas	33.6 (±4.2)	26.1 (±3.7)	14.8 (±3.0)	8.8 (±2.6)	3.5 (±1.3)	3.4 (±1.1)
Kentucky	46.2 (±3.6)	37.4 (±4.2)	19.1 (±2.5)	12.7 (±2.7)	4.2 (±1.2)	4.1 (±0.9)
Maryland	29.9 (±1.0)	23.7 (±0.9)	13.0 (±0.7)	5.0 (±0.4)	5.2 (±0.4)	7.3 (±0.5)
Minnesota	38.7 (±2.4)	32.4 (±2.3)	13.0 (±1.3)	10.2 (±1.2)	5.0 (±0.7)	4.8 (±0.9)
Mississippi — Private	50.4 (±5.0)	41.7 (±5.3)	18.8 (±3.3)	19.3 (±4.2)	5.6 (±2.5)	6.4 (±2.6)
Mississippi — Public	39.6 (±3.6)	29.2 (±3.6)	18.6 (±2.9)	9.1 (±2.3)	4.1 (±1.2)	5.5 (±1.2)
Nebraska	35.8 (±3.3)	29.0 (±3.2)	12.4 (±1.7)	10.3 (±1.9)	3.8 (±0.7)	3.8 (±0.9)
New York	32.8 (±6.7)	26.8 (±5.9)	11.5 (±3.7)	6.0 (±2.9)	4.5 (±1.7)	5.9 (±1.7)
Ohio	41.1 (±3.3)	33.4 (±3.1)	18.1 (±3.0)	9.1 (±2.0)	3.4 (±1.0)	6.9 (±1.8)
South Dakota	39.2 (±3.9)	32.6 (±3.6)	13.5 (±2.8)	11.5 (±2.7)	3.5 (±1.2)	3.8 (±1.3)
Tennessee	41.1 (±3.0)	32.4 (±2.8)	18.1 (±1.5)	12.7 (±1.9)	5.4 (±1.2)	6.4 (±1.0)
Texas	34.6 (±5.7)	28.1 (±5.8)	18.2 (±3.3)	8.3 (±3.6)	4.4 (±1.3)	3.7 (±1.0)
West Virginia	47.9 (±3.8)	38.5 (±3.4)	17.2 (±2.9)	15.3 (±2.8)	5.0 (±1.4)	8.5 (±1.7)
Wisconsin	39.4 (±3.5)	32.9 (±2.5)	16.1 (±3.0)	7.6 (±2.9)	4.7 (±1.4)	5.2 (±1.3)
Median	36.9	29.2	15.1	9.1	4.2	5.2

* Use of cigarettes or cigars or smokeless tobacco or pipes or bidis during ≥ 1 of the 30 days preceding the survey.

[†] Confidence interval.

[§] Question was not asked.

[¶] Pipe use was not included because this question was not asked.

TABLE 6. Percentage of all middle school and high school students who ever smoked cigarettes daily* and current cigarette smokers who smoked ≥ 6 cigarettes per day[†] on the days they smoked, by sex and race/ethnicity — National Youth Tobacco Survey, 2000

	<u>All students who ever smoked cigarettes daily*</u>		<u>Current smokers who smoked ≥ 6 cigarettes per day[†]</u>	
	%	(95% CI) [§]	%	(95% CI)
Middle school				
Sex				
Male	6.0	(± 1.0)	18.6	(± 3.5)
Female	5.0	(± 0.8)	13.3	(± 3.5)
Race/ethnicity				
White	6.0	(± 1.1)	18.6	(± 3.7)
Black	3.7	(± 0.9)	11.0	(± 5.7)
Hispanic	4.8	(± 0.9)	11.3	(± 3.1)
Asian	2.9	(± 1.2)	19.1	(± 13.3)
Total (middle school)	5.5	(± 0.8)	16.1	(± 2.8)
High school				
Sex				
Male	20.8	(± 1.6)	33.9	(± 2.5)
Female	20.3	(± 1.8)	29.4	(± 2.7)
Race/ethnicity				
White	23.8	(± 1.8)	34.4	(± 2.3)
Black	11.7	(± 2.4)	22.6	(± 6.0)
Hispanic	14.1	(± 2.1)	19.0	(± 3.9)
Asian	15.3	(± 3.3)	26.9	(± 7.5)
Total (high school)	20.6	(± 1.5)	31.7	(± 2.1)

* Students were asked, "Have you ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days?"

[†] Students were asked, "During the past 30 days, on the days you smoked, how many cigarettes did you smoke per day?"

[§] Confidence interval.

TABLE 7. Percentage of all middle school and high school students who ever smoked cigarettes daily* and current cigarette smokers who smoked ≥ 6 cigarettes per day on the days that they smoked, by state — State Youth Tobacco Surveys, 2000

	All students who ever smoked cigarettes daily*		Current smokers who smoked ≥ 6 cigarettes per day on the days that they smoked†	
	% (95% CI) [§]		% (95% CI)	
Middle school				
Alabama	8.4	(± 1.9)	14.6	(± 3.0)
Arizona	6.5	(± 1.1)	10.9	(± 4.8)
Arkansas	10.1	(± 1.7)	23.1	(± 5.9)
California	2.9	(± 1.0)	11.2	(± 4.4)
Colorado	5.7	(± 1.9)	11.5	(± 5.0)
Connecticut	5.4	(± 1.7)	13.1	(± 6.1)
Delaware	8.5	(± 1.4)	21.7	(± 3.1)
District of Columbia	2.7	(± 0.9)	12.9	(± 7.1)
Florida	7.0	(± 0.8)	18.1	(± 2.4)
Indiana	6.6	(± 2.1)	13.3	(± 5.5)
Iowa	7.9	(± 1.7)	20.6	(± 7.0)
Kansas	4.5	(± 1.6)	14.8	(± 7.4)
Kentucky	14.0	(± 3.1)	19.6	(± 6.4)
Maine	10.3	(± 3.8)	22.6	(± 11.4)
Maryland	4.6	(± 0.7)	14.4	(± 2.6)
Minnesota	6.3	(± 1.0)	14.0	(± 5.3)
Mississippi — Private	5.9	(± 1.9)	12.5	(± 4.4)
Mississippi — Public	8.7	(± 2.1)	13.4	(± 4.9)
New Hampshire	7.5	(± 2.1)	19.4	(± 5.5)
New York	4.0	(± 2.3)	13.5	(± 10.1)
Ohio	8.1	(± 2.2)	15.9	(± 5.4)
Tennessee	9.3	(± 1.0)	18.9	(± 2.6)
Texas	5.3	(± 0.9)	11.1	(± 5.0)
Vermont	7.7	(± 1.3)	19.8	(± 5.9)
West Virginia	12.3	(± 2.0)	27.9	(± 5.3)
Wisconsin	7.7	(± 2.0)	17.3	(± 6.9)
Wyoming	10.2	(± 1.8)	19.6	(± 5.2)
Median	7.5		14.8	
High school				
Alabama	22.0	(± 3.3)	29.9	(± 5.4)
Arkansas	28.9	(± 4.2)	41.3	(± 5.7)
California	13.6	(± 4.2)	19.0	(± 6.1)
Colorado	18.5	(± 4.1)	24.1	(± 5.0)
Connecticut	20.7	(± 3.7)	29.8	(± 6.0)
Delaware	23.4	(± 1.8)	38.4	(± 3.7)
District of Columbia	8.7	(± 2.4)	18.5	(± 6.7)
Florida	19.1	(± 1.8)	28.3	(± 2.6)
Hawaii	19.8	(± 2.6)	25.2	(± 5.2)
Indiana	25.9	(± 3.7)	39.2	(± 6.9)
Iowa	23.7	(± 2.5)	29.7	(± 4.3)
Kansas	21.0	(± 3.9)	27.7	(± 4.8)
Kentucky	30.9	(± 3.5)	40.9	(± 5.1)
Maryland	18.4	(± 0.8)	30.6	(± 1.7)
Minnesota	24.9	(± 2.1)	31.6	(± 3.4)
Mississippi — Private	28.3	(± 3.6)	30.7	(± 5.4)
Mississippi — Public	22.6	(± 3.7)	31.2	(± 6.5)
Nebraska	20.6	(± 2.3)	27.3	(± 5.0)
New York	21.6	(± 5.0)	40.6	(± 6.2)
Ohio	28.3	(± 3.4)	37.1	(± 4.4)
South Dakota	22.3	(± 3.0)	28.3	(± 3.8)
Tennessee	26.0	(± 2.9)	38.2	(± 2.7)
Texas	17.8	(± 5.4)	26.0	(± 9.0)
West Virginia	33.4	(± 3.6)	42.0	(± 5.3)
Wisconsin	25.6	(± 3.7)	36.9	(± 5.6)
Median	22.3		30.6	

* Students were asked, "Have you ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days?"

† Students were asked, "During the past 30 days, on the days you smoked, how many cigarettes did you smoke per day?"

§ Confidence interval.

TABLE 8. Percentage of all middle school and high school students who first used* cigarettes, cigars, or smokeless tobacco before age 11 years, by sex and race/ethnicity — National Youth Tobacco Survey, 2000

	<u>Cigarettes</u>		<u>Cigars</u>		<u>Smokeless tobacco</u>	
	%	(95% CI) [†]	%	(95% CI)	%	(95% CI)
Middle school						
Sex						
Male	10.0	(±1.6)	6.9	(±1.2)	6.0	(±1.8)
Female	6.8	(±0.8)	3.5	(±0.4)	1.6	(±0.4)
Race/ethnicity						
White	8.3	(±1.4)	5.1	(±1.0)	4.2	(±1.4)
Black	8.4	(±1.2)	4.7	(±0.8)	2.5	(±0.8)
Hispanic	8.7	(±1.4)	5.8	(±1.0)	2.6	(±0.8)
Asian	5.4	(±2.2)	3.0	(±1.8)	1.5	(±1.2)
Total (middle school)	8.4	(±1.0)	5.2	(±0.8)	3.7	(±1.0)
High school						
Sex						
Male	8.3	(±0.8)	4.5	(±0.6)	4.7	(±0.8)
Female	5.1	(±0.6)	1.7	(±0.4)	1.0	(±0.2)
Race/ethnicity						
White	6.9	(±0.8)	3.1	(±0.4)	3.3	(±0.6)
Black	5.3	(±1.0)	2.1	(±0.6)	1.8	(±0.6)
Hispanic	7.1	(±1.0)	4.0	(±0.8)	1.7	(±0.6)
Asian	4.7	(±1.6)	1.6	(±0.8)	1.0	(±0.6)
Total (high school)	6.7	(±0.6)	3.1	(±0.4)	2.9	(±0.4)

* Age of initiation was determined by asking students these questions: "How old were you when you smoked a whole cigarette for the first time?" "How old were you when you smoked a cigar, cigarillo, or little cigar for the first time?" "How old were you when you used chewing tobacco, snuff, or dip for the first time?"

† Confidence interval.

TABLE 9. Percentage of all middle school and high school students who first used* cigarettes, cigars, or smokeless tobacco before age 11 years[†], by state — State Youth Tobacco Surveys, 2000

	Cigarettes		Cigars		Smokeless tobacco	
	%	(95% CI [§])	%	(95% CI)	%	(95% CI)
Middle school						
Alabama	13.3	(±2.2)	9.0	(±1.6)	8.7	(±1.8)
Arizona	11.3	(±1.5)	6.5	(±1.6)	4.3	(±1.9)
Arkansas	14.8	(±1.8)	8.1	(±1.8)	8.9	(±2.1)
California	5.9	(±1.2)	3.9	(±1.1)	2.2	(±0.7)
Colorado	9.5	(±3.3)	5.8	(±1.2)	3.4	(±1.3)
Connecticut	7.5	(±1.7)	4.6	(±1.0)	1.7	(±0.7)
Delaware	12.1	(±1.7)	3.0	(±0.8)	5.3	(±0.6)
District of Columbia	6.6	(±1.6)	3.2	(±1.1)	1.6	(±0.8)
Florida	NA [¶]		NA		NA	
Indiana	10.2	(±2.2)	5.0	(±1.5)	4.0	(±1.6)
Iowa	14.0	(±2.5)	6.7	(±1.2)	5.8	(±1.8)
Kansas	10.4	(±2.2)	5.9	(±1.5)	4.7	(±1.1)
Kentucky	19.1	(±3.1)	9.1	(±1.5)	11.1	(±2.7)
Maine	12.0	(±3.5)	5.4	(±1.9)	2.5	(±1.1)
Maryland	8.9	(±1.0)	4.4	(±0.6)	2.2	(±0.3)
Minnesota	9.6	(±2.1)	5.1	(±0.8)	3.1	(±0.7)
Mississippi — Private	12.3	(±1.3)	5.2	(±1.9)	8.3	(±2.7)
Mississippi — Public	14.8	(±2.0)	8.6	(±1.7)	9.5	(±1.6)
New Hampshire	7.9	(±1.4)	3.9	(±1.0)	2.4	(±0.8)
New York	5.7	(±2.7)	4.5	(±1.3)	1.8	(±1.5)
Ohio	10.5	(±2.5)	6.5	(±1.4)	4.8	(±1.5)
Tennessee	13.3	(±1.3)	7.9	(±0.8)	8.6	(±1.1)
Texas	10.2	(±2.6)	7.8	(±1.8)	4.1	(±1.3)
Vermont	9.6	(±2.1)	5.9	(±1.4)	3.3	(±0.9)
West Virginia	15.6	(±2.3)	7.8	(±2.0)	9.8	(±1.5)
Wisconsin	10.9	(±2.5)	6.4	(±1.5)	4.2	(±1.2)
Wyoming	16.2	(±2.4)	9.3	(±1.2)	7.5	(±1.3)
Median	10.7		5.9		4.2	
High school						
Alabama	9.4	(±2.0)	3.5	(±1.0)	5.8	(±1.6)
Arkansas	14.1	(±2.5)	6.0	(±1.5)	8.9	(±3.0)
California	6.7	(±1.9)	2.9	(±1.2)	2.0	(±1.7)
Colorado	9.4	(±2.5)	4.9	(±1.2)	4.1	(±1.5)
Connecticut	6.3	(±1.2)	3.9	(±1.1)	2.0	(±0.6)
Delaware	8.1	(±1.1)	2.6	(±0.5)	2.1	(±0.5)
District of Columbia	5.8	(±1.4)	2.8	(±1.0)	3.6	(±1.2)
Florida	NA [¶]		NA		NA	
Hawaii	8.3	(±1.9)	3.6	(±1.2)	2.4	(±1.2)
Indiana	10.2	(±2.0)	4.6	(±1.8)	3.8	(±1.5)
Iowa	10.0	(±2.7)	4.3	(±1.5)	5.5	(±2.2)
Kansas	10.8	(±2.7)	4.1	(±1.2)	5.3	(±1.7)
Kentucky	14.1	(±1.9)	5.4	(±1.3)	8.1	(±2.2)
Maryland	9.3	(±0.6)	4.5	(±0.3)	3.1	(±0.3)
Minnesota	9.1	(±0.9)	4.8	(±0.7)	4.9	(±0.7)
Mississippi — Private	9.9	(±2.4)	6.7	(±1.7)	10.9	(±2.0)
Mississippi — Public	11.1	(±2.0)	5.2	(±1.3)	9.2	(±3.0)
Nebraska	9.7	(±1.3)	4.8	(±0.6)	4.7	(±1.0)
New York	6.4	(±1.8)	2.9	(±1.0)	2.1	(±0.9)
Ohio	10.9	(±2.2)	4.6	(±1.6)	3.9	(±1.3)
South Dakota	11.7	(±1.7)	4.2	(±1.1)	6.7	(±1.3)
Tennessee	10.9	(±1.0)	5.7	(±0.9)	7.9	(±0.9)
Texas	7.7	(±2.9)	4.3	(±1.4)	4.3	(±2.3)
West Virginia	14.4	(±2.0)	6.9	(±2.0)	10.6	(±2.1)
Wisconsin	10.6	(±2.6)	4.6	(±1.5)	3.5	(±1.5)
Median	9.8		4.6		4.5	

* Age of initiation was determined by asking students these questions: "How old were you when you smoked a whole cigarette for the first time?" "How old were you when you smoked a cigar, cigarillo, or little cigar for the first time?" "How old were you when you used chewing tobacco, snuff, or dip for the first time?"

[†] Age of initiation was <10 years.

[§] Confidence interval.

[¶] Question was not asked.

TABLE 10. Percentage of middle school and high school students who ever smoked cigarettes who smoked ≥ 100 cigarettes in their lifetime and percentage of all students who frequently used* cigarettes, cigars, smokeless tobacco, pipes, bidis, or kreteks, by sex and race/ethnicity — National Youth Tobacco Survey, 2000

	Ever smokers who have smoked ≥ 100 cigarettes in lifetime	All students who frequently use tobacco*					
		Cigarettes % (95% CI) [†]	Cigars % (95% CI)	Smokeless tobacco % (95% CI)	Pipes % (95% CI)	Bidis % (95% CI)	Kreteks % (95% CI)
Middle school							
Sex							
Male	5.0 (±1.0)	3.3 (±0.7)	1.4 (±0.3)	1.7 (±0.7)	1.1 (±0.3)	0.7 (±0.2)	0.6 (±0.2)
Female	3.2 (±0.7)	2.2 (±0.5)	0.5 (±0.2)	0.4 (±0.2)	0.3 (±0.1)	0.3 (±0.2)	0.2 (±0.1)
Race/ethnicity							
White	4.8 (±1.0)	3.1 (±0.7)	0.8 (±0.3)	1.1 (±0.5)	0.5 (±0.2)	0.4 (±0.2)	0.4 (±0.2)
Black	2.2 (±0.7)	2.0 (±0.8)	1.1 (±0.5)	0.8 (±0.4)	0.7 (±0.4)	0.8 (±0.4)	0.5 (±0.3)
Hispanic	2.7 (±0.7)	1.5 (±0.5)	1.2 (±0.4)	0.8 (±0.4)	1.0 (±0.4)	0.7 (±0.4)	0.6 (±0.3)
Asian	2.4 (±1.3)	1.6 (±1.0)	0.3 (±0.4)	0.3 (±0.4)	1.1 (±0.9)	0.7 (±0.6)	0.2 (±0.3)
Total (middle school)	4.1 (±0.7)	2.7 (±0.5)	0.9 (±0.2)	1.0 (±0.4)	0.7 (±0.2)	0.5 (±0.1)	0.4 (±0.1)
High school							
Sex							
Male	21.9 (±1.6)	13.8 (±1.3)	1.8 (±0.4)	3.8 (±0.8)	0.9 (±0.3)	0.8 (±0.2)	0.7 (±0.2)
Female	19.4 (±1.9)	13.4 (±1.5)	0.4 (±0.1)	0.2 (±0.1)	0.3 (±0.1)	0.1 (±0.1)	0.3 (±0.1)
Race/ethnicity							
White	24.9 (±1.9)	16.4 (±1.5)	0.8 (±0.2)	2.5 (±0.6)	0.4 (±0.1)	0.3 (±0.1)	0.3 (±0.1)
Black	8.6 (±1.7)	6.1 (±1.5)	1.8 (±0.8)	0.5 (±0.4)	0.9 (±0.5)	0.7 (±0.4)	0.7 (±0.4)
Hispanic	13.4 (±2.1)	7.4 (±1.6)	1.4 (±0.4)	1.3 (±0.6)	1.1 (±0.5)	0.9 (±0.4)	0.8 (±0.5)
Asian	14.9 (±3.1)	9.9 (±3.1)	0.6 (±0.5)	0.9 (±0.5)	0.6 (±0.5)	0.3 (±0.3)	0.4 (±0.4)
Total (high school)	20.7 (±1.5)	13.6 (±1.2)	1.1 (±0.2)	2.0 (±0.5)	0.6 (±0.2)	0.5 (±0.1)	0.5 (±0.1)

* Use of cigarettes or cigars or smokeless tobacco or pipes or bidis or kreteks on ≥ 20 of the 30 days preceding the survey.

[†] Confidence interval.

TABLE 11. Percentage of middle school and high school students who ever smoked cigarettes who smoked ≥ 100 cigarettes in their lifetime and percentage of all students who frequently used* cigarettes, cigars, smokeless tobacco, pipes, or bidis, by state — State Youth Tobacco Surveys, 2000

	Ever smokers who have smoked ≥ 100 cigarettes in lifetime	All students who frequently use tobacco*				
	% (95% CI) [†]	Cigarettes	Cigars	Smokeless tobacco	Pipes	Bidis
		% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school						
Alabama	13.0 (± 3.1)	3.5 (± 1.5)	0.9 (± 0.5)	1.8 (± 1.1)	0.6 (± 0.5)	0.7 (± 0.7)
Arizona	10.7 (± 2.9)	2.3 (± 0.8)	0.5 (± 0.4)	0.7 (± 0.6)	0.6 (± 0.5)	0.5 (± 0.5)
Arkansas	15.3 (± 3.3)	5.4 (± 1.4)	1.5 (± 0.6)	2.0 (± 0.9)	1.1 (± 0.7)	1.1 (± 0.6)
California	5.9 (± 2.8)	1.1 (± 0.5)	0.9 (± 0.2)	0.6 (± 0.2)	0.9 (± 0.2)	0.6 (± 0.2)
Colorado	8.8 (± 3.2)	2.2 (± 1.1)	0.2 (± 0.2)	0.6 (± 0.3)	0.4 (± 0.3)	0.4 (± 0.3)
Connecticut	9.6 (± 4.0)	2.5 (± 1.2)	0.8 (± 0.6)	0.4 (± 0.4)	0.5 (± 0.4)	0.5 (± 0.4)
Delaware	12.8 (± 1.7)	4.1 (± 0.7)	0.9 (± 0.4)	0.8 (± 0.5)	0.8 (± 0.4)	NA [§]
District of Columbia	3.5 (± 1.9)	0.8 (± 0.6)	0.4 (± 0.4)	0.4 (± 0.4)	0.4 (± 0.4)	0.6 (± 0.5)
Florida	NA	2.9 (± 0.5)	1.0 (± 0.2)	0.9 (± 0.2)	NA	0.8 (± 0.8)
Indiana	9.6 (± 3.5)	2.2 (± 1.1)	0.6 (± 0.4)	1.0 (± 0.5)	0.5 (± 0.3)	0.6 (± 0.5)
Iowa	13.9 (± 2.4)	3.5 (± 1.3)	0.4 (± 0.3)	0.8 (± 0.5)	0.4 (± 0.3)	0.3 (± 0.3)
Kansas	11.1 (± 2.6)	1.9 (± 0.8)	0.3 (± 0.2)	0.5 (± 0.6)	0.2 (± 0.2)	0.2 (± 0.2)
Kentucky	20.3 (± 4.6)	6.7 (± 2.5)	1.3 (± 0.8)	3.1 (± 0.7)	0.2 (± 0.3)	0.2 (± 0.4)
Maine	21.6 (± 8.3)	4.4 (± 2.3)	1.7 (± 1.3)	1.2 (± 0.8)	1.2 (± 0.8)	1.6 (± 1.3)
Maryland	8.5 (± 1.6)	1.6 (± 0.3)	0.5 (± 0.2)	0.5 (± 0.2)	0.5 (± 0.2)	0.4 (± 0.2)
Minnesota	12.1 (± 2.4)	2.5 (± 0.4)	0.2 (± 0.1)	0.5 (± 0.3)	0.2 (± 0.1)	0.2 (± 0.1)
Mississippi — Private	13.5 (± 3.7)	2.9 (± 0.7)	0.4 (± 0.5)	2.3 (± 1.2)	NA	NA
Mississippi — Public	9.9 (± 2.6)	3.8 (± 1.1)	0.9 (± 0.6)	1.5 (± 0.9)	NA	NA
New Hampshire	16.6 (± 3.9)	4.2 (± 1.2)	0.7 (± 0.5)	0.6 (± 0.4)	0.6 (± 0.4)	0.4 (± 0.4)
New York	9.8 (± 5.9)	2.2 (± 1.8)	1.1 (± 0.6)	0.9 (± 0.6)	1.0 (± 0.8)	1.0 (± 0.8)
Ohio	13.7 (± 3.8)	3.6 (± 1.4)	0.9 (± 0.6)	0.7 (± 0.6)	0.7 (± 0.5)	0.4 (± 0.3)
Tennessee	15.7 (± 2.0)	4.4 (± 0.7)	0.9 (± 0.4)	1.6 (± 0.5)	0.6 (± 0.2)	0.7 (± 0.3)
Texas	8.3 (± 2.8)	2.5 (± 1.2)	1.6 (± 0.8)	1.2 (± 0.6)	1.2 (± 0.7)	0.7 (± 0.5)
Vermont	13.9 (± 4.0)	3.7 (± 1.3)	0.2 (± 0.2)	0.6 (± 0.7)	0.3 (± 0.3)	0.1 (± 0.1)
West Virginia	20.2 (± 2.9)	7.4 (± 1.8)	1.0 (± 0.6)	2.6 (± 1.1)	0.7 (± 0.5)	0.9 (± 0.6)
Wisconsin	13.7 (± 4.4)	3.6 (± 1.2)	0.2 (± 0.2)	0.5 (± 0.5)	0.2 (± 0.2)	0.5 (± 0.5)
Wyoming	17.4 (± 3.1)	5.1 (± 1.1)	0.9 (± 0.4)	1.9 (± 0.8)	0.8 (± 0.5)	0.5 (± 0.3)
Median	13.5	3.5	0.9	0.8	0.6	0.5
High school						
Alabama	29.9 (± 4.9)	13.5 (± 2.7)	1.5 (± 0.6)	3.6 (± 1.1)	0.6 (± 0.4)	0.5 (± 0.3)
Arkansas	41.8 (± 5.3)	21.0 (± 3.7)	1.5 (± 0.8)	4.2 (± 1.8)	0.8 (± 0.5)	0.9 (± 0.7)
California	21.7 (± 6.3)	7.3 (± 3.0)	1.0 (± 0.3)	0.9 (± 0.6)	0.5 (± 0.2)	0.6 (± 0.2)
Colorado	28.0 (± 4.9)	11.1 (± 3.3)	0.8 (± 0.5)	2.6 (± 1.0)	0.9 (± 0.6)	0.9 (± 0.6)
Connecticut	30.4 (± 5.7)	12.3 (± 3.6)	0.5 (± 0.4)	0.5 (± 0.4)	0.5 (± 0.3)	0.5 (± 0.3)
Delaware	32.8 (± 2.5)	15.2 (± 1.7)	1.6 (± 0.5)	1.1 (± 0.3)	0.8 (± 0.3)	NA [§]
District of Columbia	8.4 (± 3.2)	3.5 (± 1.5)	1.0 (± 0.7)	0.8 (± 0.6)	0.8 (± 0.6)	0.3 (± 0.4)
Florida	NA	10.5 (± 1.2)	1.7 (± 0.3)	1.8 (± 0.3)	NA	1.0 (± 0.2)
Hawaii	27.0 (± 3.0)	10.3 (± 2.3)	0.8 (± 0.5)	0.8 (± 0.6)	0.6 (± 0.4)	0.8 (± 0.6)
Indiana	38.9 (± 4.2)	17.1 (± 2.5)	0.9 (± 0.5)	2.7 (± 1.3)	1.0 (± 0.6)	0.8 (± 0.4)
Iowa	37.0 (± 5.1)	14.4 (± 2.4)	1.5 (± 1.2)	3.5 (± 1.2)	1.0 (± 0.9)	0.9 (± 0.6)
Kansas	34.5 (± 5.7)	12.7 (± 2.8)	1.3 (± 0.7)	3.1 (± 1.5)	0.5 (± 0.4)	0.5 (± 0.4)
Kentucky	42.8 (± 4.2)	21.3 (± 3.3)	1.5 (± 0.5)	4.9 (± 1.7)	0.6 (± 0.6)	0.6 (± 0.4)
Maryland	26.8 (± 1.1)	10.8 (± 0.7)	1.7 (± 0.2)	1.3 (± 0.2)	1.3 (± 0.2)	1.3 (± 0.2)
Minnesota	39.5 (± 2.9)	16.9 (± 1.8)	0.7 (± 0.3)	3.0 (± 0.6)	1.0 (± 0.2)	0.6 (± 0.2)
Mississippi — Private	45.6 (± 4.1)	18.5 (± 3.2)	2.0 (± 1.5)	8.0 (± 2.3)	1.5 (± 1.6)	1.3 (± 1.4)
Mississippi — Public	29.5 (± 5.9)	13.8 (± 3.0)	2.1 (± 1.0)	3.9 (± 1.3)	0.9 (± 0.4)	0.5 (± 0.4)
Nebraska	33.3 (± 2.7)	13.2 (± 1.7)	1.0 (± 0.3)	2.6 (± 0.8)	0.9 (± 0.3)	0.6 (± 0.4)
New York	32.1 (± 6.0)	14.2 (± 4.3)	1.4 (± 0.7)	1.7 (± 1.3)	1.6 (± 1.1)	1.3 (± 0.9)
Ohio	39.2 (± 3.8)	20.7 (± 3.0)	1.6 (± 0.8)	3.2 (± 1.7)	0.6 (± 0.3)	1.1 (± 0.7)
South Dakota	35.1 (± 3.4)	15.3 (± 2.4)	0.9 (± 0.5)	3.6 (± 1.4)	0.3 (± 0.3)	0.4 (± 0.3)
Tennessee	36.4 (± 3.4)	17.4 (± 2.1)	1.5 (± 0.4)	4.6 (± 0.9)	0.9 (± 0.2)	0.6 (± 0.2)
Texas	29.3 (± 6.0)	11.3 (± 4.3)	1.3 (± 0.5)	2.8 (± 1.7)	0.8 (± 0.5)	0.6 (± 0.4)
West Virginia	44.0 (± 3.7)	22.1 (± 3.3)	1.5 (± 0.6)	5.9 (± 1.7)	1.0 (± 0.7)	1.1 (± 0.6)
Wisconsin	41.3 (± 3.6)	19.8 (± 3.0)	1.2 (± 0.8)	2.2 (± 1.3)	0.9 (± 0.6)	0.9 (± 0.7)
Median	34.5	14.2	1.4	2.8	0.8	0.7

* Smoked cigarettes on ≥ 20 of the 30 days preceding the survey; smoked cigars on ≥ 20 of the 30 days preceding the survey; used smokeless tobacco on ≥ 20 of the 30 days preceding the survey; smoked tobacco in a pipe on ≥ 20 of the 30 days preceding the survey; or smoked bidis on ≥ 20 of the 30 days preceding the survey.

[†] Confidence interval.

[§] Question was not asked.

TABLE 12. Usual brand of cigarettes smoked by current cigarette smokers* in middle school and high school during the 30 days preceding the survey (percentage distribution), by sex and race/ethnicity— National Youth Tobacco Survey, 2000

	Marlboro®	Newport®	Camel®	Other†	No usual brand	Total§
Middle school						
Sex						
Male	45.7	22.0	8.0	14.4	9.9	100.0
Female	48.6	22.8	4.4	12.8	11.4	100.0
Race/ethnicity						
White	55.6	14.1	7.5	11.8	10.9	100.0
Black	8.8	64.1	1.7	19.7	5.7	100.0
Hispanic	53.9	13.1	5.3	13.1	14.5	100.0
Asian	37.3	15.5	12.7	17.2	17.2	100.0
Total (middle school)	47.1	22.4	6.3	12.3	10.6	100.0
High school						
Sex						
Male	52.2	19.8	12.7	8.8	6.5	100.0
Female	54.6	22.0	9.2	7.5	6.7	100.0
Race/ethnicity						
White	59.4	13.5	12.8	7.7	6.6	100.0
Black	8.4	74.0	2.2	9.1	6.4	100.0
Hispanic	49.1	26.9	7.0	10.2	6.7	100.0
Asian	54.2	31.4	1.1	8.0	5.4	100.0
Total (high school)	53.3	20.8	11.0	8.1	6.6	100.0

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Includes Kool,® Virginia Slims,® GPC,® Basic,® and Doral.®

§ Some totals might not add to 100% because of rounding.

TABLE 13. Usual brand of cigarettes smoked by current cigarette smokers* in middle school and high school during the 30 days preceding the survey (percentage distribution), by state — State Youth Tobacco Surveys, 2000

	Marlboro®	Newport®	Camel®	Other†	No usual brand	Total§
Middle school						
Alabama	46.2	24.0	4.6	15.0	10.3	100.0
Arizona	50.7	6.8	8.3	12.6	21.6	100.0
Arkansas	58.7	11.4	6.3	12.4	11.2	100.0
California	53.1	8.4	6.4	18.5	13.6	100.0
Colorado	46.8	9.4	11.5	12.6	19.7	100.0
Connecticut	35.9	38.0	3.4	10.6	12.0	100.0
Delaware	24.1	60.2	2.3	6.8	6.6	100.0
District of Columbia	9.2	65.9	2.0	11.4	11.6	100.0
Florida	NA¶	NA	NA	NA	NA	NA
Indiana	44.2	22.8	10.9	7.3	14.7	100.0
Iowa	58.8	13.4	5.2	9.6	13.0	100.0
Kansas	56.3	8.3	3.0	17.9	14.5	100.0
Kentucky	63.5	8.5	4.3	16.5	7.2	100.0
Maine	57.7	4.1	7.3	18.5	12.4	100.0
Maryland	24.6	44.7	5.9	12.5	12.2	100.0
Minnesota	59.0	9.6	7.6	11.4	12.4	100.0
Mississippi — Private	66.5	5.6	8.0	10.1	9.7	100.0
Mississippi — Public	35.6	33.3	6.6	16.6	7.9	100.0
New Hampshire	60.1	8.7	9.0	10.3	11.9	100.0
New York	43.6	29.0	6.6	8.9	11.8	100.0
Ohio	48.9	19.7	8.5	9.5	13.3	100.0
Tennessee	56.0	13.5	4.4	15.1	11.1	100.0
Texas	56.7	7.6	4.4	16.4	14.9	100.0
Vermont	59.9	5.1	9.1	13.8	12.1	100.0
West Virginia	59.1	10.0	7.7	11.7	11.5	100.0
Wisconsin	41.1	29.9	6.5	11.8	10.8	100.0
Wyoming	56.7	3.7	13.1	12.7	13.7	100.0
Median	54.6	10.7	6.6	12.4	12.0	
High school						
Alabama	52.0	28.7	6.5	7.0	5.8	100.0
Arkansas	64.0	17.7	7.7	7.3	3.3	100.0
California	55.9	11.7	12.8	11.2	8.4	100.0
Colorado	61.8	8.4	12.2	7.5	10.1	100.0
Connecticut	44.4	30.0	12.6	5.2	7.8	100.0
Delaware	32.4	55.1	4.3	3.7	4.4	100.0
District of Columbia	5.8	76.3	2.7	8.5	6.7	100.0
Florida	NA¶	NA	NA	NA	NA	NA
Hawaii	40.4	5.4	2.1	45.8	6.3	100.0
Indiana	51.9	16.8	20.5	4.8	6.0	100.0
Iowa	71.9	6.5	10.5	5.5	5.6	100.0
Kansas	66.6	8.4	10.6	5.4	9.0	100.0
Kentucky	72.7	10.5	5.4	6.9	4.5	100.0
Maryland	35.8	41.0	7.4	7.0	8.7	100.0
Minnesota	57.3	7.3	24.0	4.9	6.4	100.0
Mississippi — Private	75.7	5.8	10.1	5.5	3.0	100.0
Mississippi — Public	53.0	30.5	5.4	5.6	5.5	100.0
Nebraska	67.2	9.9	9.8	5.0	8.1	100.0
New York	44.7	28.6	4.5	15.5	6.7	100.0
Ohio	60.0	15.7	10.3	7.3	6.6	100.0
South Dakota	61.4	5.7	19.7	4.6	8.6	100.0
Tennessee	63.8	14.8	9.4	5.9	6.1	100.0
Texas	66.8	10.1	6.3	9.5	7.2	100.0
West Virginia	59.3	9.4	15.5	9.1	6.7	100.0
Wisconsin	49.5	20.1	18.3	6.8	5.3	100.0
Median	58.3	13.2	10.0	6.8	6.5	

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Includes Virginia Slims,® GPC,® Basic,® and Doral.®

§ Some totals might not add to 100% because of rounding.

¶ Question was not asked.

TABLE 14. Percentage of middle school and high school students who never smoked cigarettes, by smoking intentions, susceptibility* to initiate cigarette smoking, by sex and race/ethnicity — National Youth Tobacco Survey, 2000

	Will not try a cigarette soon	Will definitely not smoke a cigarette in the next year	Would definitely not smoke a cigarette if best friend offered a cigarette	Not susceptible	Susceptible
	% (95% CI)[†]	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school					
Sex					
Male	96.2 (±0.6)	84.0 (±1.5)	83.1 (±1.5)	75.1 (±1.6)	24.9 (±1.6)
Female	96.0 (±0.7)	83.4 (±1.6)	84.4 (±1.4)	76.3 (±1.7)	23.7 (±1.7)
Race/ethnicity					
White	96.7 (±0.6)	83.8 (±1.6)	84.1 (±1.4)	76.8 (±1.7)	23.2 (±1.7)
Black	95.1 (±1.4)	85.6 (±2.0)	83.7 (±1.9)	74.7 (±2.3)	25.3 (±2.3)
Hispanic	94.3 (±1.2)	81.1 (±1.9)	82.5 (±1.9)	72.0 (±2.2)	28.0 (±2.2)
Asian	94.9 (±2.3)	82.8 (±4.6)	80.9 (±4.4)	74.0 (±4.9)	26.0 (±4.9)
Total (middle school)	96.1 (±0.5)	83.7 (±1.3)	83.7 (±1.1)	75.7 (±1.4)	24.3 (±1.4)
High school					
Sex					
Male	97.7 (±0.6)	84.1 (±1.7)	84.9 (±1.6)	78.3 (±2.1)	21.7 (±2.1)
Female	96.5 (±0.7)	81.0 (±1.9)	84.1 (±1.5)	76.7 (±2.0)	23.3 (±2.0)
Race/ethnicity					
White	97.5 (±0.6)	81.9 (±2.0)	83.9 (±1.6)	77.9 (±2.2)	22.1 (±2.2)
Black	96.2 (±1.4)	85.9 (±2.8)	88.3 (±2.3)	78.9 (±3.3)	21.1 (±3.3)
Hispanic	96.0 (±1.5)	80.1 (±3.2)	82.1 (±3.4)	72.3 (±3.9)	27.7 (±3.9)
Asian	96.1 (±1.8)	85.5 (±3.4)	86.8 (±3.2)	79.8 (±4.3)	20.2 (±4.3)
Total (high school)	97.1 (±0.5)	82.5 (±1.5)	84.5 (±1.2)	77.5 (±1.6)	22.5 (±1.6)

* Students were considered not susceptible if they answered, "No, will not try a cigarette soon" and "Definitely no, will not smoke a cigarette in the next year" and "Definitely no, would not smoke a cigarette if best friend offered a cigarette." All other students were considered susceptible.

† Confidence interval.

TABLE 15. Percentage of middle school and high school students who never smoked cigarettes, by smoking intentions, susceptibility* to initiate cigarette smoking, and state—State Youth Tobacco Surveys, 2000

	<u>Will not try a cigarette soon</u>	<u>Will definitely not smoke a cigarette in the next year</u>	<u>Would definitely not smoke a cigarette if best friend offered</u>	<u>Not susceptible</u>	<u>Susceptible</u>
	% (95% CI) [†]	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school					
Alabama	93.6 (±1.7)	81.0 (±3.8)	82.7 (±3.0)	72.2 (±3.9)	27.8 (±3.9)
Arizona	94.0 (±1.4)	81.6 (±1.9)	83.1 (±2.5)	72.8 (±3.1)	27.2 (±3.1)
Arkansas	96.1 (±1.1)	85.2 (±2.2)	87.2 (±1.7)	78.0 (±2.4)	22.0 (±2.4)
California	95.9 (±1.2)	83.3 (±3.6)	83.3 (±2.2)	74.2 (±2.8)	25.8 (±2.8)
Colorado	96.0 (±1.0)	84.5 (±2.7)	85.1 (±2.4)	77.1 (±2.8)	22.9 (±2.8)
Connecticut	93.2 (±1.9)	79.7 (±2.7)	83.3 (±2.0)	72.7 (±3.3)	27.3 (±3.3)
Delaware	95.5 (±1.0)	83.6 (±2.4)	84.6 (±2.2)	73.6 (±2.8)	26.4 (±2.8)
District of Columbia	93.2 (±1.7)	78.4 (±2.9)	82.7 (±3.0)	69.8 (±3.8)	30.2 (±3.8)
Florida	95.1 (±0.5)	82.7 (±1.1)	84.7 (±1.0)	76.9 (±1.3)	23.1 (±1.3)
Indiana	95.9 (±1.3)	86.4 (±2.0)	87.4 (±2.4)	80.1 (±2.1)	19.9 (±2.1)
Iowa	96.1 (±1.2)	84.2 (±3.1)	84.1 (±3.1)	75.1 (±4.6)	24.9 (±4.6)
Kansas	95.9 (±1.3)	85.8 (±2.5)	84.3 (±2.6)	76.0 (±3.2)	24.0 (±3.2)
Kentucky	95.8 (±1.8)	85.4 (±2.6)	87.3 (±2.3)	78.9 (±3.2)	21.1 (±3.2)
Maine	98.0 (±1.4)	82.2 (±4.9)	82.5 (±4.6)	74.7 (±5.1)	25.3 (±5.1)
Maryland	94.7 (±0.6)	84.5 (±1.7)	85.5 (±1.3)	74.9 (±1.9)	25.1 (±1.9)
Minnesota	95.4 (±0.9)	81.3 (±1.7)	82.7 (±1.9)	73.6 (±2.0)	26.4 (±2.0)
Mississippi — Private	94.7 (±2.0)	80.0 (±5.2)	82.2 (±5.2)	73.4 (±6.7)	26.6 (±6.7)
Mississippi — Public	94.2 (±2.1)	81.0 (±3.5)	82.4 (±3.8)	73.2 (±3.7)	26.8 (±3.7)
New Hampshire	95.6 (±1.5)	78.0 (±3.9)	81.8 (±3.5)	71.8 (±3.8)	28.2 (±3.8)
New York	94.8 (±1.5)	84.2 (±5.0)	84.4 (±4.3)	76.5 (±4.7)	23.5 (±4.7)
Ohio	95.5 (±1.4)	79.8 (±3.4)	82.5 (±3.0)	72.0 (±4.2)	28.0 (±4.2)
Tennessee	94.4 (±0.9)	82.4 (±1.4)	85.1 (±1.6)	76.1 (±1.8)	23.9 (±1.8)
Texas	94.8 (±1.7)	82.6 (±4.4)	84.7 (±3.9)	74.8 (±4.5)	25.2 (±4.5)
Vermont	95.2 (±1.8)	79.8 (±3.1)	82.3 (±3.6)	72.7 (±3.7)	27.3 (±3.7)
West Virginia	96.3 (±1.3)	83.1 (±2.6)	84.0 (±2.1)	74.8 (±3.1)	25.2 (±3.1)
Wisconsin	93.3 (±1.7)	77.7 (±3.8)	78.7 (±3.8)	68.2 (±4.4)	31.8 (±4.4)
Wyoming	95.2 (±1.6)	82.0 (±2.6)	83.4 (±2.7)	74.2 (±2.9)	25.8 (±2.9)
Median	95.2	82.4	83.4	74.2	25.8
High school					
Alabama	96.6 (±1.7)	87.3 (±3.4)	89.4 (±3.1)	82.7 (±4.2)	17.3 (±4.2)
Arkansas	96.9 (±1.6)	83.2 (±3.2)	86.0 (±3.7)	75.8 (±5.0)	24.2 (±5.0)
California	95.4 (±1.4)	80.0 (±3.7)	83.4 (±2.4)	74.3 (±3.9)	25.7 (±3.9)
Colorado	97.7 (±1.2)	83.3 (±3.3)	85.3 (±2.6)	80.2 (±3.4)	19.8 (±3.4)
Connecticut	97.4 (±1.1)	82.1 (±3.7)	87.0 (±2.8)	76.1 (±4.3)	23.9 (±4.3)
Delaware	99.0 (±0.5)	87.6 (±2.4)	87.2 (±2.4)	77.9 (±3.4)	22.1 (±3.4)
District of Columbia	96.2 (±2.0)	83.7 (±3.6)	85.9 (±3.7)	77.5 (±4.2)	22.5 (±4.2)
Florida	96.3 (±0.7)	83.8 (±1.4)	85.4 (±1.2)	79.2 (±1.7)	20.8 (±1.7)
Hawaii	96.7 (±1.9)	84.6 (±3.5)	85.6 (±3.6)	77.1 (±3.8)	22.9 (±3.8)
Indiana	95.5 (±1.5)	83.3 (±3.7)	86.2 (±3.3)	78.0 (±4.5)	22.0 (±4.5)
Iowa	95.9 (±1.8)	82.5 (±3.0)	85.7 (±3.6)	76.3 (±3.8)	23.7 (±3.8)
Kansas	96.2 (±1.8)	83.1 (±3.7)	84.4 (±3.4)	77.0 (±5.1)	23.0 (±5.1)
Kentucky	98.9 (±1.1)	83.8 (±4.7)	85.4 (±3.8)	79.6 (±5.3)	20.4 (±5.3)
Maryland	95.3 (±0.6)	80.5 (±1.1)	82.7 (±1.0)	72.7 (±1.2)	27.3 (±1.2)
Minnesota	96.4 (±0.8)	80.0 (±1.9)	84.2 (±1.6)	74.2 (±2.1)	25.8 (±2.1)
Mississippi — Private	97.1 (±2.4)	71.1 (±8.0)	81.3 (±6.0)	66.5 (±8.5)	33.5 (±8.5)
Mississippi — Public	97.3 (±1.9)	82.5 (±3.7)	87.3 (±4.5)	75.1 (±5.3)	24.9 (±5.3)
Nebraska	95.0 (±1.2)	80.4 (±2.6)	82.7 (±2.7)	73.7 (±2.6)	26.3 (±2.6)
New York	96.4 (±0.8)	84.3 (±2.0)	83.9 (±2.9)	76.3 (±2.5)	23.7 (±2.5)
Ohio	97.1 (±1.5)	82.0 (±4.5)	84.1 (±3.5)	77.2 (±4.5)	22.8 (±4.5)
South Dakota	95.4 (±1.5)	82.7 (±4.9)	83.0 (±4.3)	75.3 (±5.1)	24.7 (±5.1)
Tennessee	96.4 (±0.9)	83.3 (±2.3)	86.4 (±2.4)	78.3 (±2.9)	21.7 (±2.9)
Texas	96.3 (±1.7)	80.6 (±3.5)	82.8 (±3.4)	76.6 (±4.1)	23.4 (±4.1)
West Virginia	96.6 (±2.1)	80.1 (±5.0)	81.3 (±5.2)	73.3 (±4.7)	26.7 (±4.7)
Wisconsin	97.2 (±1.6)	81.9 (±3.8)	83.1 (±4.2)	76.5 (±4.3)	23.5 (±4.3)
Median	96.4	82.7	85.3	76.5	23.5

* Students were considered not susceptible if they answered "No, will not try a cigarette soon" and "Definitely no, will not smoke a cigarette in the next year" and "Definitely no, would not smoke a cigarette if best friend offered a cigarette." All other students were considered susceptible.

† Confidence interval.

TABLE 16. Percentage of middle school and high school students with peers who use tobacco, by tobacco use status, sex, and race/ethnicity — National Youth Tobacco Survey, 2000

	≥1 Closest friends smoke cigarettes		≥1 Closest friends use smokeless tobacco	
	Never cigarette smoker	Current cigarette smoker*	Never smokeless tobacco user	Current smokeless tobacco user†
	% (95% CI‡)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school				
Sex				
Male	16.2 (±1.7)	90.3 (±2.0)	8.0 (±1.3)	80.6 (± 8.5)
Female	17.6 (±2.0)	92.4 (±2.6)	6.6 (±1.1)	68.0 (±10.0)
Race/ethnicity				
White	16.3 (±2.0)	91.5 (±2.0)	7.7 (±1.5)	83.5 (± 7.1)
Black	19.7 (±2.6)	87.4 (±4.9)	5.3 (±1.2)	54.3 (±13.8)
Hispanic	19.0 (±2.7)	87.9 (±4.4)	7.4 (±1.3)	72.5 (± 9.6)
Asian	9.5 (±3.6)	¶	6.2 (±2.8)	¶
Total (middle school)	17.0 (±1.5)	90.3 (±2.0)	7.3 (±1.0)	78.0 (± 7.1)
High school				
Sex				
Male	32.7 (±2.8)	91.4 (±1.2)	14.6 (±2.0)	82.2 (± 3.3)
Female	33.2 (±2.5)	93.2 (±0.9)	12.0 (±1.6)	67.6 (± 8.3)
Race/ethnicity				
White	33.8 (±3.4)	93.5 (±0.9)	16.4 (±2.1)	83.0 (± 3.4)
Black	30.6 (±4.3)	82.8 (±4.3)	5.5 (±1.6)	65.0 (±11.4)
Hispanic	34.6 (±3.1)	90.4 (±2.4)	9.4 (±1.4)	71.7 (± 8.2)
Asian	25.6 (±5.4)	95.0 (±2.9)	6.3 (±1.6)	¶
Total (high school)	33.0 (±2.5)	92.3 (±0.9)	13.1 (±1.6)	80.7 (± 3.1)

* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

† Used smokeless tobacco on ≥1 of the 30 days preceding the survey.

‡ Confidence interval.

¶ n<35.

TABLE 17. Percentage of middle school and high school students with peers who use tobacco*, by tobacco use status and state — State Youth Tobacco Surveys, 2000

	≥1 Closest friends smoke cigarettes		≥1 Closest friends use smokeless tobacco	
	Never cigarette smoker	Current cigarette smoker†	Never smokeless tobacco user	Current smokeless tobacco user‡
	% (95% CI¶)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school				
Alabama	19.6 (±4.0)	92.0 (± 4.0)	18.7 (±2.4)	79.9 (± 8.4)
Arizona	15.1 (±2.6)	87.6 (± 4.2)	15.1 (±1.9)	66.6 (± 9.6)
Arkansas	18.0 (±3.7)	88.4 (± 4.7)	22.4 (±4.5)	79.9 (± 9.6)
California	12.2 (±1.8)	82.8 (± 7.8)	14.5 (±1.4)	74.0 (±14.7)
Colorado	10.8 (±2.5)	76.1 (±10.3)	5.9 (±1.4)	**
Connecticut	14.4 (±2.4)	86.4 (± 6.0)	11.1 (±2.0)	67.4 (±21.0)
Delaware††	21.0 (±2.3)	95.3 (± 2.2)	19.7 (±2.0)	72.4 (± 8.8)
District of Columbia	14.7 (±3.1)	72.1 (± 9.1)	5.5 (±1.4)	**
Florida	16.3 (±1.1)	86.4 (± 2.4)	6.4 (±0.7)	71.8 (± 4.6)
Indiana	11.9 (±4.0)	71.7 (± 6.4)	6.2 (±2.1)	52.1 (±15.0)
Iowa	13.2 (±3.2)	87.8 (± 5.1)	15.5 (±3.0)	84.9 (± 7.8)
Kansas	10.8 (±2.9)	73.0 (± 6.6)	4.3 (±1.4)	57.9 (±18.6)
Kentucky	19.5 (±4.4)	93.4 (± 3.3)	21.8 (±4.4)	87.0 (± 6.2)
Maine	18.5 (±5.1)	74.2 (± 6.1)	7.4 (±3.3)	60.5 (±17.1)
Maryland	11.8 (±1.1)	84.3 (± 3.4)	11.9 (±1.1)	65.7 (± 6.8)
Minnesota	10.7 (±1.6)	85.8 (± 3.6)	11.3 (±1.7)	61.4 (± 9.1)
Mississippi — Private	15.4 (±2.7)	92.5 (± 4.2)	22.9 (±3.2)	84.9 (± 6.7)
Mississippi — Public	22.2 (±4.6)	86.6 (± 4.8)	25.2 (±3.5)	77.0 (± 8.4)
New Hampshire	12.9 (±3.5)	85.5 (± 5.9)	15.1 (±3.2)	75.4 (±14.6)
New York	15.2 (±4.2)	87.7 (± 6.2)	14.5 (±1.8)	94.3 (± 7.8)
Ohio	13.2 (±3.0)	88.9 (± 5.1)	17.2 (±3.2)	72.8 (±11.1)
Tennessee	15.7 (±1.8)	86.9 (± 2.5)	19.7 (±2.1)	79.3 (± 4.3)
Texas	18.2 (±4.8)	90.4 (± 3.7)	17.3 (±2.5)	70.7 (±12.3)
Vermont	17.1 (±2.8)	88.2 (± 5.8)	16.7 (±2.0)	77.4 (±25.7)
West Virginia	22.1 (±3.1)	90.3 (± 4.0)	25.9 (±4.0)	82.4 (± 6.0)
Wisconsin	14.3 (±4.3)	89.5 (± 4.5)	14.8 (±2.5)	83.7 (±11.9)
Wyoming	14.0 (±2.8)	86.3 (± 5.1)	19.1 (±2.4)	77.8 (± 6.1)
Median	15.1	86.9	15.1	74.0
High school				
Alabama	32.2 (±4.1)	89.8 (± 3.0)	25.1 (±3.5)	82.7 (± 8.5)
Arkansas	37.2 (±6.0)	90.6 (± 3.7)	27.3 (±4.3)	83.4 (± 4.8)
California	29.5 (±4.1)	91.4 (± 3.3)	18.9 (±3.1)	78.8 (± 9.7)
Colorado	26.2 (±3.8)	81.5 (± 4.4)	15.5 (±2.1)	72.5 (±10.5)
Connecticut	29.9 (±6.5)	92.8 (± 3.7)	14.8 (±2.2)	81.5 (± 8.1)
Delaware††	37.2 (±3.6)	96.4 (± 1.4)	25.3 (±2.4)	84.5 (± 5.9)
District of Columbia	25.5 (±4.6)	67.3 (± 8.2)	3.8 (±1.2)	50.2 (±17.4)
Florida	29.6 (±2.0)	91.9 (± 1.2)	9.2 (±1.1)	78.7 (± 4.4)
Hawaii	25.5 (±5.7)	76.4 (± 6.4)	9.9 (±2.1)	61.6 (±16.3)
Indiana	27.3 (±4.0)	81.6 (± 3.5)	11.9 (±2.0)	86.8 (± 7.5)
Iowa	31.8 (±6.8)	91.7 (± 3.3)	22.6 (±2.8)	86.5 (± 6.2)
Kansas	26.7 (±3.7)	77.7 (± 4.3)	17.3 (±3.2)	64.4 (± 7.5)
Kentucky	36.2 (±6.0)	93.7 (± 1.9)	31.1 (±4.3)	90.4 (± 4.2)
Maryland	27.3 (±1.3)	88.4 (± 1.1)	17.1 (±0.7)	75.5 (± 3.1)
Minnesota	27.4 (±2.4)	92.2 (± 1.4)	20.2 (±1.9)	82.5 (± 3.2)
Mississippi — Private	44.3 (±6.8)	90.2 (± 5.0)	41.0 (±3.9)	90.9 (± 3.2)
Mississippi — Public	22.3 (±4.3)	86.9 (± 3.7)	27.3 (±3.6)	78.6 (± 7.3)
Nebraska	26.2 (±3.1)	82.9 (± 2.6)	15.8 (±2.5)	73.7 (± 4.8)
New York	31.3 (±3.9)	91.5 (± 2.3)	18.3 (±4.8)	77.8 (± 8.3)
Ohio	27.1 (±4.0)	92.7 (± 2.3)	22.0 (±3.6)	72.9 (±10.7)
South Dakota	30.8 (±4.5)	79.7 (± 4.5)	18.8 (±3.3)	79.0 (± 5.2)
Tennessee	29.9 (±2.9)	92.4 (± 1.3)	26.0 (±3.0)	82.8 (± 3.6)
Texas	27.2 (±3.5)	94.4 (± 1.9)	21.6 (±4.7)	86.2 (± 3.7)
West Virginia	30.3 (±4.7)	92.5 (± 2.7)	34.7 (±5.0)	82.9 (± 4.9)
Wisconsin	32.1 (±5.5)	94.5 (± 2.5)	17.9 (±3.4)	87.7 (± 7.9)
Median	29.6	91.4	18.9	81.5

* Students were asked, "How many of your *four* closest friends smoke cigarettes?" and "How many of your *four* closest friends use chewing tobacco, snuff, or dip?"

† Smoked cigarettes on ≥1 of the 30 days preceding the survey.

‡ Use of smokeless tobacco on ≥1 of the 30 days preceding the survey.

¶ Confidence interval.

** n<35.

†† For the survey in Delaware, students were asked, "How many of your closest friends smoke cigarettes?" and "How many of your closest friends use chewtobacco, snuff, or dip?"

TABLE 18. Percentage of middle school and high school students with perception that smokers have more friends, by cigarette smoking status, sex, and race/ethnicity — National Youth Tobacco Survey, 2000

	Think cigarette smokers have more friends	
	Never cigarette smoker	Current cigarette smoker*
	% (95% CI)[†]	% (95% CI)
Middle school		
Sex		
Male	12.5 (±1.5)	42.6 (±4.0)
Female	11.3 (±1.2)	32.7 (±3.5)
Race/ethnicity		
White	9.2 (±1.1)	34.5 (±3.2)
Black	21.5 (±2.3)	45.6 (±6.4)
Hispanic	14.3 (±1.8)	42.5 (±5.9)
Asian	15.6 (±4.4)	§
Total (middle school)	11.9 (±1.1)	37.9 (±2.7)
High school		
Sex		
Male	15.4 (±1.7)	29.1 (±2.4)
Female	9.9 (±1.5)	18.5 (±1.9)
Race/ethnicity		
White	9.0 (±1.2)	20.6 (±1.7)
Black	21.0 (±3.0)	37.6 (±5.9)
Hispanic	19.5 (±2.8)	30.7 (±4.3)
Asian	20.1 (±4.4)	39.0 (±7.2)
Total (high school)	12.6 (±1.4)	24.0 (±1.6)

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Confidence interval.

§ n<35.

TABLE 19. Percentage of middle school and high school students with perception that smokers have more friends, by smoking status and state — State Youth Tobacco Surveys, 2000

	Think smokers have more friends	
	Never smoker % (95% CI) [†]	Current smoker* % (95% CI)
Middle school		
Alabama	17.2 (±3.8)	43.5 (± 5.8)
Arizona	10.6 (±2.1)	43.3 (± 9.2)
Arkansas	11.2 (±3.4)	40.0 (± 5.2)
California	14.0 (±2.1)	34.6 (± 6.3)
Colorado	9.1 (±2.0)	44.5 (± 7.3)
Connecticut	9.8 (±2.0)	41.3 (± 7.4)
Delaware	13.7 (±2.2)	37.2 (± 6.7)
District of Columbia	26.8 (±3.1)	48.5 (± 9.6)
Florida	16.9 (±2.4)	31.9 (± 3.7)
Indiana	8.0 (±2.2)	53.0 (±12.0)
Iowa	7.6 (±1.8)	38.5 (± 5.6)
Kansas	6.2 (±1.4)	33.6 (±11.3)
Kentucky	12.1 (±2.3)	39.1 (± 9.5)
Maine	8.6 (±3.4)	37.9 (±12.8)
Maryland	13.1 (±1.4)	46.0 (± 4.9)
Minnesota	8.8 (±1.5)	41.5 (± 6.2)
Mississippi — Private	14.8 (±2.7)	41.5 (± 8.0)
Mississippi — Public	21.0 (±4.1)	45.9 (± 6.7)
New Hampshire	10.1 (±2.4)	40.7 (± 6.9)
New York	13.8 (±3.5)	37.0 (± 9.6)
Ohio	9.7 (±2.3)	40.7 (± 5.8)
Tennessee	10.5 (±1.2)	38.7 (± 3.4)
Texas	10.4 (±3.1)	32.9 (± 6.3)
Vermont	10.5 (±2.9)	48.6 (± 8.4)
West Virginia	8.2 (±2.2)	38.6 (± 7.3)
Wisconsin	8.8 (±2.9)	42.2 (±10.3)
Wyoming	9.5 (±2.0)	42.2 (± 6.2)
Median	10.5	40.7
High school		
Alabama	14.9 (±4.5)	26.5 (± 3.5)
Arkansas	13.0 (±4.3)	25.2 (± 4.2)
California	16.1 (±3.9)	25.0 (± 3.6)
Colorado	8.5 (±2.5)	26.1 (± 5.0)
Connecticut	13.7 (±4.9)	28.7 (± 6.7)
Delaware	12.9 (±2.3)	22.2 (± 3.0)
District of Columbia	24.2 (±4.5)	49.4 (± 7.8)
Florida	19.2 (±3.0)	28.6 (± 3.5)
Hawaii	22.0 (±4.2)	36.9 (± 6.2)
Indiana	9.5 (±2.6)	26.8 (± 3.9)
Iowa	9.5 (±3.8)	26.0 (± 3.4)
Kansas	9.5 (±2.4)	22.6 (± 5.4)
Kentucky	11.4 (±3.4)	28.0 (± 3.8)
Maryland	17.0 (±1.3)	34.5 (± 1.6)
Minnesota	13.2 (±2.6)	31.1 (± 2.7)
Mississippi — Private	19.1 (±3.9)	24.8 (± 3.7)
Mississippi — Public	21.8 (±4.1)	28.3 (± 5.5)
Nebraska	8.5 (±1.4)	28.0 (± 3.6)
New York	17.4 (±5.0)	27.6 (± 5.8)
Ohio	12.4 (±4.1)	24.3 (± 5.3)
South Dakota	8.5 (±3.2)	28.3 (± 4.7)
Tennessee	11.3 (±1.5)	30.3 (± 2.9)
Texas	13.4 (±3.9)	24.4 (± 4.7)
West Virginia	10.6 (±3.4)	28.8 (± 3.6)
Wisconsin	11.1 (±2.4)	22.7 (± 3.8)
Median	13.0	27.6

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Confidence interval.

TABLE 20. How current cigarette smokers* aged <18 years in middle school and high school usually obtained cigarettes (percentage distribution), by sex and race/ethnicity — National Youth Tobacco Survey, 2000

	Bought in a store	Bought from a vending machine	Bought by someone else	Borrowed from someone else	Took them from store or family member	Given by person aged ≥18 yrs	Obtained some other way	Total†
Middle school								
Sex								
Male	8.9	3.7	24.1	21.1	13.5	10.3	18.3	100.0
Female	5.5	1.3	24.8	23.8	8.3	14.5	21.7	100.0
Race/ethnicity								
White	6.0	1.7	26.6	24.3	9.5	12.7	19.0	100.0
Black	9.8	4.5	19.6	18.1	16.2	11.6	20.3	100.0
Hispanic	8.4	1.9	21.5	19.6	11.5	12.3	24.7	100.0
Asian	9.5	6.4	23.3	19.8	10.9	9.3	20.9	100.0
Total (middle school)	7.3	2.6	24.4	22.4	11.0	12.3	19.9	100.0
High school								
Sex								
Male	35.9	1.1	21.3	20.2	4.5	7.1	9.9	100.0
Female	28.2	0.6	29.3	21.7	1.1	10.7	8.4	100.0
Race/ethnicity								
White	32.4	0.5	27.1	21.1	2.4	8.3	8.2	100.0
Black	32.1	1.6	15.1	19.3	4.3	14.6	12.9	100.0
Hispanic	32.5	2.0	17.9	22.9	3.2	8.7	12.8	100.0
Asian	34.0	2.5	27.0	15.6	3.6	7.3	9.9	100.0
Total (high school)	32.2	0.9	25.1	20.9	2.9	8.8	9.2	100.0

* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

† Some totals might not add to 100% because of rounding.

TABLE 21. How current cigarette smokers* aged <18 years in middle school and high school usually obtained cigarettes (percentage distribution), by state — State Youth Tobacco Surveys, 2000

	Bought in a store	Bought from a vending machine	Bought by someone else	Borrowed from someone else	Took them from store or family member	Given by person aged ≥18 yrs	Obtained some other way	Total
Middle school								
Alabama	8.2	3.1	24.6	28.5	10.5	9.4	15.7	100.0
Arizona	9.5	6.1	15.2	23.5	16.8	8.5	20.3	100.0
Arkansas	7.2	2.8	26.2	22.6	12.5	10.7	18.0	100.0
California	9.8	2.2	21.2	19.8	9.0	6.8	31.2	100.0
Colorado	5.2	7.0	15.2	23.6	16.0	10.5	22.5	100.0
Connecticut	12.3	1.4	17.7	27.7	12.2	7.2	21.4	100.0
Delaware	9.0	2.7	23.6	27.4	12.0	11.4	13.9	100.0
District of Columbia	15.4	3.4	14.5	23.8	17.7	6.2	19.0	100.0
Florida	5.0	1.6	15.9	24.4	14.0	13.7	25.5	100.0
Indiana	9.6	2.6	25.8	30.8	7.4	10.1	13.6	100.0
Iowa	2.9	2.4	24.2	29.6	10.4	10.3	20.3	100.0
Kansas	7.5	1.5	23.3	21.5	11.2	6.4	28.7	100.0
Kentucky	5.3	3.3	21.7	33.0	10.8	12.2	13.7	100.0
Maine	3.3	1.0	26.5	30.6	15.5	4.2	19.0	100.0
Maryland	7.4	6.9	20.5	23.9	14.0	8.7	18.6	100.0
Minnesota	1.3	2.3	23.8	29.0	19.0	7.5	17.1	100.0
Mississippi — Private	5.3	1.3	33.7	28.3	11.8	6.6	13.0	100.0
Mississippi — Public	6.1	1.8	23.7	26.5	13.5	10.9	17.4	100.0
New Hampshire	NA [†]	NA	NA	NA	NA	NA	NA	NA
New York	11.9	1.2	22.8	21.3	9.0	8.0	25.7	100.0
Ohio	6.7	3.8	27.7	38.1	8.7	5.9	9.2	100.0
Tennessee	6.2	3.4	22.3	25.6	15.8	10.4	16.3	100.0
Texas	8.2	2.2	19.8	24.7	9.4	14.0	21.8	100.0
Vermont	1.9	0.7	20.4	27.4	15.6	6.5	27.5	100.0
West Virginia	5.0	3.9	27.8	22.0	11.9	9.0	20.4	100.0
Wisconsin	8.8	4.0	24.5	25.7	12.0	12.2	12.8	100.0
Wyoming	3.3	2.7	20.3	32.4	11.4	8.5	21.5	100.0
Median	7.0	2.6	23.0	26.1	12.0	8.8	19.0	
High school								
Alabama	27.6	1.1	25.9	27.5	4.7	3.7	9.5	100.0
Arkansas	14.7	0.5	40.4	19.9	3.4	8.5	12.7	100.0
California	16.2	1.0	29.4	25.3	3.0	9.1	15.9	100.0
Colorado	15.9	2.7	28.2	33.5	2.2	5.5	11.9	100.0
Connecticut	29.4	1.7	23.2	27.5	3.2	7.2	7.8	100.0
Delaware	30.6	1.0	27.1	24.1	3.3	6.6	7.4	100.0
District of Columbia	29.2	2.3	8.4	29.4	8.4	7.2	15.0	100.0
Florida	17.4	0.8	26.5	25.7	5.5	11.9	12.4	100.0
Hawaii	16.7	1.6	28.9	23.3	3.5	11.9	14.1	100.0
Indiana	19.1	0.5	35.4	23.9	4.4	9.3	7.4	100.0
Iowa	9.9	1.5	38.2	31.5	5.2	7.4	6.3	100.0
Kansas	15.8	1.0	37.7	25.9	2.2	8.5	8.9	100.0
Kentucky	21.7	1.5	32.3	24.9	2.0	9.4	8.2	100.0
Maryland	26.9	3.5	25.0	24.3	4.5	6.9	8.9	100.0
Minnesota	15.9	1.7	41.7	23.6	2.7	5.7	8.7	100.0
Mississippi — Private	26.0	2.1	32.6	26.8	2.2	5.1	5.2	100.0
Mississippi — Public	22.9	2.4	30.9	21.2	3.7	11.4	7.6	100.0
Nebraska	8.4	0.8	42.8	28.4	4.4	7.4	7.7	100.0
New York	36.2	0.6	19.8	23.3	3.3	7.8	9.0	100.0
Ohio	24.9	2.0	33.5	23.8	2.8	5.1	7.9	100.0
South Dakota	8.1	1.7	45.0	30.5	2.6	7.4	4.6	100.0
Tennessee	21.5	2.2	33.6	21.6	3.8	8.7	8.7	100.0
Texas	19.9	1.2	28.2	28.1	1.2	10.4	11.0	100.0
West Virginia	20.4	2.6	33.6	21.5	2.7	7.3	12.0	100.0
Wisconsin	25.2	0.3	38.2	21.2	3.0	5.8	6.3	100.0
Median	20.4	1.5	32.3	24.9	3.3	7.4	8.7	

* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

† Question was not asked.

TABLE 22. Where current cigarette smokers* aged <18 years in middle school and high school bought their last pack of cigarettes (percentage distribution), by sex and race/ethnicity—National Youth Tobacco Survey, 2000

	Gas station	Convenience store	Grocery store	Drug store	Vending machine	Internet	Restaurant	Total†
Middle school								
Sex								
Male	39.9	23.6	4.4	7.6	5.8	14.4	4.3	100.0
Female	39.1	23.4	5.8	14.0	5.4	10.5	1.9	100.0
Race/ethnicity								
White	40.5	23.0	5.2	10.4	5.5	13.5	2.0	100.0
Black	38.3	26.4	5.2	10.4	4.2	9.8	5.7	100.0
Hispanic	38.1	24.5	3.3	10.2	8.9	11.2	3.8	100.0
Asian	45.4	17.6	9.2	9.7	6.5	11.6	0.0	100.0
Total (middle school)	39.5	23.5	5.0	10.4	5.6	12.6	3.2	100.0
High school								
Sex								
Male	55.8	26.7	3.0	7.8	2.6	2.4	1.7	100.0
Female	58.5	25.9	2.8	7.5	2.3	1.9	1.1	100.0
Race/ethnicity								
White	60.6	24.7	2.6	7.3	2.0	1.5	1.2	100.0
Black	36.9	35.7	4.9	10.4	4.2	5.6	2.3	100.0
Hispanic	50.2	31.0	3.0	8.3	2.0	3.5	2.1	100.0
Asian	39.7	29.6	6.8	9.7	9.6	2.6	1.9	100.0
Total (high school)	57.1	26.3	2.9	7.7	2.5	2.2	1.4	100.0

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Some totals might not add to 100% because of rounding.

TABLE 23. Where current cigarette smokers* aged <18 years in middle school and high school bought their last pack of cigarettes (percentage distribution), by state — State Youth Tobacco Surveys, 2000

	Gas station	Convenience store	Grocery store	Drug store	Vending machine	Internet	Other	Total†
Middle school								
Alabama	26.0	11.8	5.2	0.0	2.9	1.6	52.4	100.0
Arizona	20.5	9.0	4.6	4.7	8.7	0.8	51.6	100.0
Arkansas	19.1	15.0	6.3	3.5	4.9	1.4	49.7	100.0
California	NA§	NA	NA	NA	NA	NA	NA	NA
Colorado	9.5	17.0	2.3	2.0	8.7	1.4	59.0	100.0
Connecticut	19.4	13.3	2.5	5.4	5.1	4.2	50.2	100.0
Delaware	27.7	15.8	4.1	1.3	5.4	1.0	44.8	100.0
District of Columbia	17.0	19.3	6.6	1.9	6.8	0.0	48.4	100.0
Florida	NA	NA	NA	NA	NA	NA	NA	NA
Indiana	33.6	8.5	6.4	4.2	6.8	0.0	40.4	100.0
Iowa	27.0	9.4	2.6	1.2	1.3	1.9	56.5	100.0
Kansas	24.9	10.6	5.3	0.0	7.9	0.0	51.3	100.0
Kentucky	18.2	12.3	6.8	2.8	7.1	1.1	51.8	100.0
Maine	34.2	31.3	16.9	10.8	4.2	0.0	2.7	100.0
Maryland	18.3	13.5	6.3	3.8	7.3	2.3	48.5	100.0
Minnesota	21.8	5.9	7.3	1.8	3.6	0.8	58.7	100.0
Mississippi — Private	NA	NA	NA	NA	NA	NA	NA	NA
Mississippi — Public	NA	NA	NA	NA	NA	NA	NA	NA
New Hampshire	NA	NA	NA	NA	NA	NA	NA	NA
New York	NA	NA	NA	NA	NA	NA	NA	NA
Ohio	20.1	15.7	6.4	2.1	4.2	0.0	51.5	100.0
Tennessee	23.1	7.2	6.0	1.6	5.0	0.6	56.5	100.0
Texas	NA	NA	NA	NA	NA	NA	NA	NA
Vermont	13.2	12.1	5.0	4.0	2.8	1.3	61.6	100.0
West Virginia	19.1	13.9	6.9	3.2	6.0	1.0	50.0	100.0
Wisconsin	30.1	7.8	3.0	4.0	4.7	2.6	47.9	100.0
Wyoming	21.1	14.1	6.5	1.1	2.9	1.2	53.0	100.0
Median	20.8	12.8	6.2	2.4	5.0	1.0	51.4	
High school								
Alabama	49.8	19.0	5.9	1.3	0.4	0.3	23.2	100.0
Arkansas	47.6	20.2	6.8	0.9	1.9	1.3	21.3	100.0
California	NA†	NA	NA	NA	NA	NA	NA	NA
Colorado	39.8	15.3	6.6	5.0	2.2	2.5	28.5	100.0
Connecticut	48.4	23.1	4.7	1.5	4.1	2.6	15.7	100.0
Delaware	41.8	28.5	4.2	3.4	1.3	1.1	19.7	100.0
District of Columbia	21.3	35.9	4.0	1.3	1.2	2.5	33.8	100.0
Florida	NA	NA	NA	NA	NA	NA	NA	NA
Hawaii	14.3	25.7	16.7	7.4	1.0	0.0	34.9	100.0
Indiana	57.0	13.6	6.8	2.3	0.0	0.8	19.5	100.0
Iowa	52.9	16.5	5.2	1.4	2.1	1.9	20.0	100.0
Kansas	45.8	17.1	8.7	1.0	2.8	0.0	24.6	100.0
Kentucky	42.1	24.3	8.3	0.5	1.7	1.7	21.5	100.0
Maryland	42.0	22.3	5.9	3.1	3.7	2.4	20.5	100.0
Minnesota	55.1	16.0	3.7	1.7	1.1	1.3	21.1	100.0
Mississippi — Private	47.2	27.3	4.7	1.5	0.7	3.1	15.5	100.0
Mississippi — Public	45.4	21.0	6.2	3.0	3.1	0.4	20.9	100.0
Nebraska	43.2	19.7	7.5	0.3	0.3	1.4	27.6	100.0
New York	NA	NA	NA	NA	NA	NA	NA	NA
Ohio	45.7	17.7	5.9	1.6	2.2	2.1	24.8	100.0
South Dakota	47.2	22.1	5.9	0.7	2.6	1.7	19.8	100.0
Tennessee	47.2	20.9	7.0	1.4	1.7	1.3	20.5	100.0
Texas	NA	NA	NA	NA	NA	NA	NA	NA
West Virginia	41.4	25.2	6.8	0.6	1.6	2.3	22.0	100.0
Wisconsin	66.8	11.0	4.8	1.6	2.1	0.3	13.5	100.0
Median	45.8	20.9	5.9	1.5	1.7	1.4	21.1	

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Some totals might not add to 100% because of rounding.

§ Question was not asked.

TABLE 24. Percentage of current cigarette smokers* aged <18 years in middle school and high school who purchased cigarettes in a store and were not asked to show proof of age or who were not refused purchase because of their age, by sex and race/ethnicity—National Youth Tobacco Survey, 2000

	<u>Were not asked to show proof of age when purchasing cigarettes</u> % (95% CI) [†]	<u>Were not refused purchase of cigarettes because of age</u> % (95% CI)
Middle school		
Sex		
Male	64.5 (± 6.9)	58.3 (± 5.7)
Female	76.8 (± 5.8)	68.4 (± 5.7)
Race/ethnicity		
White	69.9 (± 6.9)	63.0 (± 6.3)
Black	69.3 (±10.3)	58.7 (± 9.4)
Hispanic	72.3 (± 7.8)	58.8 (± 6.8)
Asian	[§]	[§]
Total (middle school)	69.4 (± 4.7)	62.4 (± 4.2)
High school		
Sex		
Male	54.5 (± 4.2)	54.4 (± 3.7)
Female	62.9 (± 4.4)	62.9 (± 3.7)
Race/ethnicity		
White	56.1 (± 4.3)	57.8 (± 3.4)
Black	64.9 (± 7.9)	58.8 (± 9.6)
Hispanic	67.5 (± 5.9)	60.5 (± 6.5)
Asian	36.4 (± 9.9)	45.9 (±13.0)
Total (high school)	58.1 (± 3.6)	58.1 (± 2.9)

* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

[†] Confidence interval.

[§] n<35.

TABLE 25. Percentage of current cigarette smokers* aged <18 years in middle school and high school who purchased cigarettes in a store and were not asked to show proof of age or who were not refused purchase because of their age, by state — State Youth Tobacco Surveys, 2000

	Were not asked to show proof of age when purchasing cigarettes % (95% CI) [†]	Were not refused purchase of cigarettes because of age % (95% CI)
Middle school		
Alabama	67.8 (±13.1)	61.1 (± 9.6)
Arizona	87.1 (± 9.7)	74.5 (±11.7)
Arkansas	65.1 (±11.7)	67.7 (±11.2)
California	66.4 (± 8.8)	52.1 (± 6.5)
Colorado	86.9 (± 8.6)	77.2 (±11.4)
Connecticut	73.6 (±12.3)	65.1 (±13.9)
Delaware	70.0 (± 5.5)	66.1 (±10.8)
District of Columbia	56.7 (±14.2)	55.5 (±16.0)
Florida	74.0 (± 5.0)	65.3 (± 4.3)
Indiana	76.3 (± 9.6)	66.1 (±12.6)
Iowa	81.0 (± 8.6)	70.5 (±12.3)
Kansas	82.8 (±15.4)	58.7 (±18.4)
Kentucky	74.7 (±11.2)	69.8 (± 8.5)
Maine	§	§
Maryland	72.0 (± 5.6)	64.4 (± 5.9)
Minnesota	78.5 (± 6.8)	78.8 (± 4.8)
Mississippi — Private	61.4 (±13.4)	¶
Mississippi — Public	70.3 (± 8.6)	¶
New Hampshire	90.5 (± 9.5)	69.3 (±12.2)
New York	57.3 (±23.8)	56.9 (±11.5)
Ohio	79.1 (± 7.0)	60.2 (± 9.2)
Tennessee	68.5 (± 6.9)	68.3 (± 3.4)
Texas	71.3 (± 8.0)	66.2 (± 8.3)
Vermont	87.2 (±10.6)	77.8 (±14.0)
West Virginia	81.3 (± 8.2)	65.1 (± 7.6)
Wisconsin	89.4 (± 7.4)	74.4 (± 8.6)
Wyoming	74.6 (±11.6)	73.4 (± 9.7)
Median	74.0	66.1
High school		
Alabama	60.7 (± 8.7)	57.4 (± 9.7)
Arkansas	58.3 (± 7.1)	60.3 (± 6.9)
California	51.6 (± 9.4)	57.1 (± 5.1)
Colorado	54.2 (± 8.7)	52.5 (±11.7)
Connecticut	50.9 (±11.5)	49.7 (±11.6)
Delaware	59.5 (±5.1)	57.7 (± 5.7)
District of Columbia	61.3 (±13.0)	51.2 (±10.9)
Florida	60.0 (± 4.0)	60.3 (± 3.1)
Hawaii	63.9 (±11.8)	56.7 (± 9.4)
Indiana	55.2 (± 7.0)	60.6 (± 8.4)
Iowa	66.1 (± 8.7)	69.2 (± 6.7)
Kansas	56.9 (±11.4)	62.0 (±12.7)
Kentucky	65.9 (± 8.5)	67.9 (±11.1)
Maryland	60.0 (± 2.2)	61.5 (± 2.1)
Minnesota	58.9 (± 4.0)	62.6 (± 3.4)
Mississippi — Private	46.9 (±10.3)	55.7 (± 6.4)
Mississippi — Public	52.2 (± 8.6)	62.4 (± 5.2)
Nebraska	68.3 (± 4.7)	70.6 (± 4.3)
New York	50.0 (± 5.8)	55.8 (± 7.2)
Ohio	65.5 (± 9.3)	53.5 (± 8.0)
South Dakota	56.5 (±10.4)	64.0 (±10.1)
Tennessee	60.1 (± 2.8)	64.1 (± 3.3)
Texas	56.9 (± 5.5)	53.4 (± 6.8)
West Virginia	63.6 (± 8.3)	63.0 (± 8.4)
Wisconsin	60.9 (± 8.7)	59.1 (± 7.5)
Median	59.5	60.3

* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

† Confidence interval.

§ n<35.

¶ Question was not asked.

TABLE 26. Percentage of all middle school and high school students who were influenced by media and advertising regarding tobacco, by sex and race/ethnicity — National Youth Tobacco Survey, 2000

	Saw antismoking commercials on TV or heard on radio in preceding 30 days % (95% CI)*	Saw actors using tobacco on TV or in movies % (95% CI)	Saw advertising for tobacco products on the Internet % (95% CI)
Middle school			
Sex			
Male	78.9 (±1.4)	82.8 (±1.2)	31.9 (±1.5)
Female	84.0 (±1.2)	83.8 (±1.2)	32.9 (±1.8)
Race/ethnicity			
White	83.4 (±1.3)	84.4 (±1.0)	31.5 (±1.9)
Black	76.5 (±2.2)	81.0 (±2.2)	35.0 (±2.0)
Hispanic	79.2 (±2.5)	81.5 (±1.8)	33.5 (±1.9)
Asian	82.1 (±4.9)	81.2 (±3.9)	28.6 (±4.0)
Total (middle school)	81.5 (±1.1)	83.3 (±0.8)	32.4 (±1.4)
High school			
Sex			
Male	82.9 (±1.5)	86.4 (±1.0)	23.5 (±1.2)
Female	85.6 (±1.1)	89.7 (±0.8)	25.3 (±1.5)
Race/ethnicity			
White	86.1 (±1.2)	89.2 (±0.8)	22.5 (±1.2)
Black	78.2 (±2.5)	84.6 (±2.0)	27.8 (±2.2)
Hispanic	81.5 (±2.0)	86.2 (±1.4)	30.6 (±2.6)
Asian	87.5 (±2.5)	87.4 (±2.9)	27.3 (±4.4)
Total (high school)	84.2 (±1.1)	88.0 (±0.8)	24.4 (±1.0)

* Confidence interval.

TABLE 27. Percentage of all middle school and high school students who were influenced by media and advertising regarding tobacco, by state — State Youth Tobacco Surveys, 2000

	Saw antismoking commercials on TV or heard on radio in preceding 30 days	Saw actors using tobacco on TV or in movies	Saw advertising for tobacco products on the Internet
	% (95% CI)*	% (95% CI)	% (95% CI)
Middle school			
Alabama	79.2 (±3.2)	85.3 (±2.9)	40.8 (±3.9)
Arizona	80.7 (±2.2)	82.6 (±2.3)	33.9 (±2.8)
Arkansas	79.0 (±3.4)	NA	37.0 (±3.6)
California	NA†	NA	29.7 (±2.7)
Colorado	76.7 (±4.3)	81.4 (±2.5)	30.2 (±2.9)
Connecticut	81.2 (±2.3)	79.3 (±3.9)	27.4 (±2.7)
Delaware	81.1 (±1.4)	85.2 (±1.7)	34.1 (±2.5)
District of Columbia	78.3 (±2.9)	83.4 (±2.5)	36.5 (±3.2)
Florida	79.8 (±1.1)	NA	NA
Indiana	76.2 (±3.0)	81.2 (±3.0)	33.0 (±3.2)
Iowa	79.8 (±3.0)	83.7 (±1.9)	35.0 (±3.7)
Kansas	78.7 (±2.7)	82.2 (±2.6)	31.9 (±3.4)
Kentucky	80.5 (±2.8)	89.7 (±2.0)	40.0 (±5.4)
Maine	75.7 (±3.8)	84.2 (±4.1)	37.9 (±5.6)
Maryland	72.2 (±1.4)	81.9 (±1.3)	NA
Minnesota	81.0 (±2.4)	84.0 (±1.5)	32.2 (±2.6)
Mississippi — Private	86.1 (±2.9)	NA	NA
Mississippi — Public	81.6 (±2.0)	NA	NA
New Hampshire	82.0 (±2.7)	86.3 (±1.9)	25.3 (±1.9)
New York	NA	NA	30.8 (±3.9)
Ohio	78.6 (±3.1)	87.5 (±2.1)	32.7 (±4.2)
Tennessee	78.7 (±1.3)	87.7 (±1.3)	34.4 (±2.2)
Texas	NA	NA	32.2 (±3.2)
Vermont	78.3 (±2.2)	84.0 (±2.7)	32.1 (±3.8)
West Virginia	77.9 (±3.5)	89.7 (±1.7)	38.1 (±2.8)
Wisconsin	83.5 (±2.3)	85.5 (±2.9)	35.1 (±3.6)
Wyoming	77.7 (±2.6)	88.8 (±1.3)	34.3 (±2.7)
Median	78.9	84.1	33.9
High school			
Alabama	86.7 (±1.9)	90.9 (±1.8)	30.5 (±2.8)
Arkansas	84.9 (±2.0)	NA	28.4 (±3.6)
California	NA†	NA	27.2 (±5.1)
Colorado	87.2 (±1.5)	88.5 (±1.7)	21.7 (±2.0)
Connecticut	82.6 (±2.9)	88.1 (±1.8)	20.0 (±2.5)
Delaware	85.5 (±1.7)	90.6 (±1.1)	25.8 (±2.6)
District of Columbia	78.3 (±2.6)	86.1 (±1.8)	29.0 (±3.5)
Florida	83.2 (±1.7)	NA	NA
Hawaii	78.7 (±2.4)	89.3 (±2.4)	23.2 (±3.9)
Indiana	86.9 (±1.9)	88.0 (±2.0)	24.1 (±2.9)
Iowa	86.6 (±2.3)	92.1 (±1.9)	25.6 (±2.8)
Kansas	83.9 (±2.6)	88.3 (±2.1)	28.6 (±2.9)
Kentucky	84.3 (±2.0)	91.2 (±2.1)	28.3 (±4.1)
Maryland	79.8 (±0.8)	88.3 (±0.5)	NA
Minnesota	86.0 (±1.4)	89.7 (±1.1)	28.2 (±1.7)
Mississippi — Private	87.3 (±2.4)	91.7 (±1.5)	26.3 (±2.5)
Mississippi — Public	83.5 (±2.0)	90.4 (±1.3)	34.9 (±3.7)
Nebraska	85.2 (±1.4)	90.8 (±0.9)	24.7 (±2.7)
New York	NA	NA	25.1 (±2.2)
Ohio	76.3 (±3.5)	89.4 (±1.6)	23.4 (±3.8)
South Dakota	86.5 (±1.9)	90.0 (±1.3)	24.1 (±2.5)
Tennessee	82.8 (±1.1)	91.5 (±1.1)	27.9 (±2.1)
Texas	NA	NA	26.3 (±2.6)
West Virginia	83.0 (±2.5)	89.2 (±1.7)	30.8 (±3.8)
Wisconsin	87.3 (±1.5)	91.8 (±1.5)	23.1 (±3.2)
Median	84.6	89.8	26.3

* Confidence interval.

† Question was not asked.

TABLE 28. Percentage of middle school and high school students who participated in any community event to discourage persons from using tobacco products, by tobacco use status, sex, and race/ethnicity — National Youth Tobacco Survey, 2000

	<u>Never used tobacco</u> % (95% CI) [†]	<u>Ever used tobacco*</u> % (95% CI)
Middle school		
Sex		
Male	14.2 (±1.6)	16.1 (±1.7)
Female	16.2 (±1.5)	15.5 (±1.5)
Race/ethnicity		
White	15.0 (±1.8)	14.3 (±1.8)
Black	18.5 (±2.3)	18.7 (±2.2)
Hispanic	13.3 (±2.1)	17.5 (±2.3)
Asian	12.2 (±3.5)	13.9 (±6.9)
Total (middle school)	15.3 (±1.4)	15.8 (±1.3)
High school		
Sex		
Male	8.8 (±1.2)	9.7 (±1.0)
Female	12.6 (±1.5)	8.8 (±0.9)
Race/ethnicity		
White	10.7 (±1.4)	7.6 (±0.8)
Black	13.0 (±2.8)	13.2 (±2.2)
Hispanic	8.5 (±1.8)	11.6 (±1.8)
Asian	10.1 (±2.5)	12.6 (±3.2)
Total (high school)	10.8 (±1.0)	9.3 (±0.8)

* Ever use of cigarettes *or* cigars *or* smokeless tobacco *or* pipes *or* bidis *or* kreteks.

[†] Confidence interval.

TABLE 29. Percentage of middle school and high school students who participated in any community event to discourage persons from using tobacco products, by tobacco use status and state — State Youth Tobacco Surveys, 2000

	Never used tobacco		Ever used tobacco*	
	% (95% CI)†		% (95% CI)	
Middle School				
Alabama	29.7	(±4.2)	23.8	(±3.6)
Arizona	18.2	(±3.1)	18.2	(±4.2)
Arkansas	40.2	(±6.7)	26.5	(±3.9)
California	13.8	(±2.1)	18.7	(±3.0)
Colorado	35.0	(±5.1)	22.7	(±4.2)
Connecticut	25.3	(±3.6)	22.4	(±4.1)
Delaware	26.4	(±2.8)	21.0	(±2.8)
District of Columbia	33.3	(±5.3)	32.3	(±5.3)
Florida	14.4	(±1.2)	14.4	(±1.0)
Indiana	18.5	(±5.0)	19.0	(±4.3)
Iowa	34.5	(±7.8)	26.6	(±4.7)
Kansas	28.4	(±4.7)	22.1	(±5.6)
Kentucky	28.0	(±7.3)	18.0	(±4.1)
Maine	8.9	(±3.6)	14.6	(±5.4)
Maryland	23.9	(±1.9)	27.3	(±2.3)
Minnesota	24.7	(±2.9)	24.0	(±4.0)
Mississippi — Private	26.9	(±9.4)	20.1	(±5.6)
Mississippi — Public	32.4	(±3.9)	27.1	(±4.1)
New Hampshire	10.3	(±2.8)	16.0	(±5.2)
New York	12.1	(±4.4)	11.9	(±2.6)
Ohio	19.4	(±4.7)	18.4	(±4.5)
Tennessee	23.9	(±2.9)	19.3	(±2.1)
Texas	14.2	(±3.5)	15.0	(±3.2)
Vermont	27.3	(±5.5)	20.9	(±6.4)
West Virginia	27.3	(±5.3)	24.3	(±4.5)
Wisconsin	29.1	(±8.3)	27.5	(±5.7)
Wyoming	34.5	(±4.4)	22.5	(±3.4)
Median	26.4		21.0	
High school				
Alabama	17.0	(±5.0)	14.4	(±3.1)
Arkansas	33.7	(±6.4)	15.2	(±4.2)
California	10.5	(±2.7)	11.7	(±2.1)
Colorado	17.1	(±4.7)	14.9	(±3.3)
Connecticut	12.5	(±3.7)	15.3	(±3.1)
Delaware	15.2	(±2.4)	12.2	(±2.4)
District of Columbia	22.0	(±5.8)	22.8	(±4.4)
Florida	10.0	(±0.8)	10.5	(±1.2)
Hawaii	25.1	(±6.2)	25.7	(±3.6)
Indiana	16.7	(±3.9)	8.9	(±2.2)
Iowa	19.2	(±6.0)	12.1	(±2.9)
Kansas	22.1	(±3.5)	13.6	(±2.9)
Kentucky	20.2	(±7.6)	11.8	(±1.9)
Maryland	15.2	(±1.1)	17.1	(±0.9)
Minnesota	19.2	(±2.7)	16.0	(±1.6)
Mississippi — Private	12.2	(±4.7)	8.9	(±4.2)
Mississippi — Public	28.0	(±6.5)	20.0	(±4.4)
Nebraska	24.1	(±3.6)	15.0	(±1.9)
New York	8.9	(±2.4)	11.7	(±3.3)
Ohio	19.9	(±5.3)	9.8	(±1.8)
South Dakota	21.3	(±6.3)	16.6	(±4.2)
Tennessee	19.0	(±3.7)	15.3	(±3.0)
Texas	9.0	(±2.6)	9.9	(±1.9)
West Virginia	23.9	(±7.6)	15.9	(±3.5)
Wisconsin	13.2	(±4.7)	14.5	(±3.6)
Median	19.0		14.5	

* Ever use of cigarettes or cigars or smokeless tobacco or bidis or kreteks.

† Confidence interval.

TABLE 30. Percentage of middle school and high school students receptive to tobacco company merchandise,* by tobacco use status, sex, and race/ethnicity—National Youth Tobacco Survey, 2000

	Bought or received anything with tobacco company name or picture on it		Would wear or use something with tobacco company name or picture on it	
	Never tobacco user % (95% CI) [§]	Current tobacco user [†] % (95% CI)	Never tobacco user % (95% CI)	Current tobacco user % (95% CI)
Middle school				
Sex				
Male	12.2 (±1.3)	48.4 (± 3.5)	15.3 (±1.6)	61.7 (± 3.3)
Female	10.7 (±1.5)	41.7 (± 3.7)	7.0 (±1.0)	51.3 (± 4.1)
Race/ethnicity				
White	11.6 (±1.4)	50.4 (± 3.1)	10.0 (±1.3)	62.6 (± 3.7)
Black	10.0 (±1.5)	29.0 (± 4.7)	11.4 (±1.6)	41.4 (± 6.5)
Hispanic	12.2 (±2.0)	45.1 (± 5.3)	14.1 (±2.0)	54.5 (± 5.5)
Asian	7.8 (±2.4)	36.6 (±12.5)	12.6 (±3.4)	55.2 (±14.5)
Total (middle school)	11.4 (±1.1)	45.5 (± 2.7)	10.8 (±1.0)	57.3 (± 3.1)
High school				
Sex				
Male	11.6 (±1.2)	37.3 (± 2.0)	20.6 (±1.7)	62.4 (± 2.4)
Female	10.6 (±1.2)	31.0 (± 2.6)	12.1 (±1.5)	54.2 (± 2.9)
Race/ethnicity				
White	11.1 (±1.2)	35.4 (± 2.2)	15.9 (±1.6)	63.4 (± 2.7)
Black	8.8 (±2.0)	29.2 (± 4.9)	15.1 (±2.7)	40.3 (± 4.3)
Hispanic	13.0 (±1.8)	35.2 (± 3.1)	19.0 (±2.7)	53.4 (± 3.9)
Asian	12.9 (±3.3)	33.7 (± 6.1)	12.7 (±3.3)	42.8 (± 7.2)
Total (high school)	11.1 (±0.9)	34.6 (± 1.8)	16.0 (±1.2)	58.9 (± 2.4)

* For example, a cigarette lighter or T-shirt.

[†] Use of cigarettes *or* cigars *or* smokeless tobacco *or* pipes *or* bidis *or* kreteks on ≥ 1 of the 30 days preceding the survey.

[§] Confidence interval.

TABLE 31. Percentage of middle school and high school students receptive to tobacco company merchandise,* by tobacco use status and state—State Youth Tobacco Surveys, 2000

	Bought or received anything with tobacco company name or picture on it		Would wear or use something with tobacco company name or picture on it	
	Never tobacco user	Current tobacco user†	Never tobacco user	Current tobacco user
	% (95% CI‡)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school				
Alabama	9.0 (±3.0)	44.2 (± 5.6)	10.8 (±3.2)	57.2 (± 4.9)
Arizona	15.0 (±2.8)	41.0 (± 5.6)	15.4 (±2.8)	57.8 (± 6.6)
Arkansas	9.9 (±2.4)	45.2 (± 5.1)	18.7 (±5.2)	63.0 (± 5.8)
California	9.5 (±1.4)	46.9 (± 8.2)	10.8 (±1.8)	46.7 (± 5.6)
Colorado	11.5 (±1.6)	37.1 (± 7.6)	8.1 (±2.7)	46.9 (± 6.7)
Connecticut	16.0 (±2.6)	48.3 (± 6.9)	13.6 (±2.2)	51.1 (± 7.0)
Delaware	10.6 (±1.2)	42.1 (± 3.8)	12.5 (±1.3)	54.1 (± 3.4)
District of Columbia	8.4 (±2.1)	29.4 (± 8.5)	17.4 (±3.0)	36.5 (± 8.2)
Florida	11.0 (±0.8)	38.2 (± 2.6)	14.6 (±0.9)	56.2 (± 2.2)
Indiana	12.7 (±3.1)	41.4 (± 6.2)	11.2 (±2.5)	53.5 (± 8.2)
Iowa	12.2 (±2.9)	50.1 (± 5.0)	14.7 (±2.5)	65.4 (± 7.6)
Kansas	12.6 (±2.8)	49.4 (± 9.1)	10.3 (±1.9)	60.8 (± 6.4)
Kentucky	15.1 (±3.9)	49.7 (± 4.9)	13.8 (±3.6)	68.1 (± 4.8)
Maine	20.5 (±4.2)	43.0 (±12.7)	16.2 (±6.8)	61.9 (±15.9)
Maryland	12.9 (±1.2)	42.6 (± 4.0)	11.1 (±1.2)	53.1 (± 3.4)
Minnesota	13.8 (±1.5)	50.5 (± 6.4)	8.7 (±1.7)	61.4 (± 4.2)
Mississippi — Private	13.7 (±3.0)	48.9 (± 6.3)	13.1 (±3.6)	62.3 (± 4.0)
Mississippi — Public	13.3 (±2.7)	45.1 (± 5.2)	17.5 (±2.9)	56.7 (± 5.8)
New Hampshire	14.0 (±2.4)	42.7 (± 5.7)	12.8 (±2.5)	59.2 (± 6.6)
New York	10.3 (±2.7)	44.7 (± 6.8)	11.6 (±2.4)	44.1 (± 8.3)
Ohio	11.7 (±2.4)	50.3 (± 5.4)	9.7 (±1.8)	58.7 (±10.3)
Tennessee	13.7 (±1.7)	46.7 (± 2.7)	13.8 (±1.4)	60.6 (± 2.8)
Texas	14.1 (±3.0)	45.2 (± 4.6)	13.6 (±3.2)	60.9 (± 5.6)
Vermont	9.6 (±2.4)	52.2 (± 6.6)	12.8 (±2.1)	64.8 (± 9.0)
West Virginia	16.0 (±2.8)	58.4 (± 4.8)	14.9 (±3.7)	71.1 (± 5.1)
Wisconsin	14.6 (±4.2)	47.1 (± 7.9)	12.8 (±2.8)	58.4 (± 8.2)
Wyoming	14.9 (±2.1)	48.2 (± 5.1)	16.5 (±2.7)	66.9 (± 5.0)
Median	12.9	45.2	13.1	58.7
High school				
Alabama	10.7 (±3.3)	36.1 (± 4.1)	21.8 (±4.3)	61.4 (± 5.6)
Arkansas	9.9 (±2.9)	38.3 (± 4.2)	22.3 (±5.9)	65.4 (± 4.7)
California	10.9 (±2.4)	34.6 (± 4.6)	16.8 (±3.1)	53.0 (± 3.7)
Colorado	10.9 (±3.3)	32.4 (± 3.3)	13.5 (±2.9)	55.6 (± 6.1)
Connecticut	16.4 (±3.8)	39.5 (± 7.3)	16.1 (±3.6)	54.1 (± 6.1)
Delaware	9.6 (±2.1)	30.7 (± 3.1)	14.4 (±2.5)	49.4 (± 3.8)
District of Columbia	10.5 (±3.1)	21.4 (± 5.9)	17.6 (±3.7)	39.5 (± 7.3)
Florida	9.9 (±1.0)	32.5 (± 1.8)	19.3 (±1.6)	58.9 (± 1.8)
Hawaii	12.3 (±3.6)	39.9 (± 5.6)	15.3 (±3.5)	54.7 (± 6.3)
Indiana	13.5 (±3.0)	36.4 (± 3.3)	15.6 (±3.9)	61.7 (± 4.5)
Iowa	9.7 (±3.2)	42.0 (± 4.7)	18.0 (±3.4)	67.2 (± 6.1)
Kansas	15.1 (±2.7)	40.2 (± 7.0)	17.4 (±4.6)	64.3 (± 4.8)
Kentucky	13.1 (±4.6)	46.6 (± 4.4)	18.8 (±4.0)	64.7 (± 4.9)
Maryland	12.7 (±1.0)	37.3 (± 1.5)	16.7 (±1.1)	54.6 (± 1.5)
Minnesota	14.0 (±1.5)	42.7 (± 1.8)	13.7 (±1.4)	63.8 (± 2.5)
Mississippi — Private	16.1 (±5.9)	46.9 (± 4.3)	24.7 (±5.1)	65.9 (± 5.2)
Mississippi — Public	13.1 (±4.3)	38.0 (± 3.4)	22.7 (±4.1)	57.8 (± 4.2)
Nebraska	12.5 (±1.7)	40.4 (± 3.0)	20.0 (±2.2)	69.9 (± 3.6)
New York	13.2 (±2.9)	33.9 (± 6.2)	18.0 (±4.0)	52.4 (± 4.1)
Ohio	10.6 (±3.6)	31.6 (± 6.2)	18.6 (±4.3)	55.0 (± 8.0)
South Dakota	10.7 (±3.3)	41.7 (± 4.3)	17.4 (±3.1)	65.3 (± 4.7)
Tennessee	13.2 (±2.2)	45.0 (± 2.0)	17.0 (±1.8)	60.7 (± 3.0)
Texas	9.7 (±1.6)	36.2 (± 6.0)	16.3 (±2.4)	59.7 (± 5.2)
West Virginia	17.9 (±5.2)	45.6 (± 4.8)	24.5 (±4.5)	67.0 (± 3.6)
Wisconsin	10.3 (±3.1)	42.5 (± 4.8)	16.6 (±3.4)	70.3 (± 4.2)
Median	12.3	38.3	17.4	60.7

* For example, a cigarette lighter or T-shirt.

† Use of cigarettes or cigars or smokeless tobacco or pipes or bidis or kreteks on ≥1 of the 30 days preceding the survey.

‡ Confidence interval.

TABLE 32. Percentage of current cigarette smokers* in middle school and high school who tried to quit and who want to stop smoking cigarettes, by sex and race/ethnicity—National Youth Tobacco Survey, 2000

	Tried to quit smoking cigarettes during preceding 12 months % (95% CI)[†]	Want to stop smoking cigarettes % (95% CI)
Middle school		
Sex		
Male	54.7 (±4.1)	51.5 (±5.4)
Female	65.6 (±4.3)	58.7 (±4.3)
Race/ethnicity		
White	60.1 (±3.5)	54.8 (±5.0)
Black	61.3 (±7.8)	58.6 (±7.4)
Hispanic	56.0 (±7.2)	53.1 (±7.2)
Asian	[§]	[§]
Total (middle school)	59.9 (±3.1)	55.0 (±3.9)
High school		
Sex		
Male	55.2 (±2.6)	59.2 (±2.3)
Female	63.6 (±2.4)	62.9 (±2.5)
Race/ethnicity		
White	59.4 (±2.2)	60.5 (±1.8)
Black	61.5 (±3.9)	66.1 (±4.6)
Hispanic	52.3 (±4.5)	57.5 (±4.2)
Asian	70.5 (±7.2)	73.5 (±7.6)
Total (high school)	59.3 (±2.0)	61.0 (±1.6)

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.[†] Confidence interval.[§] n<35.

TABLE 33. Percentage of current cigarette smokers* in middle school and high school who tried to quit and who want to stop smoking cigarettes, by state — State Youth Tobacco Surveys, 2000

	<u>Tried to quit smoking cigarettes during preceding 12 months</u> % (95% CI) [†]	<u>Want to stop smoking cigarettes</u> % (95% CI)
Middle school		
Alabama	58.2 (± 5.9)	54.9 (± 7.6)
Arizona	57.0 (± 8.3)	58.4 (± 9.7)
Arkansas	56.2 (± 7.2)	59.3 (± 9.7)
California	63.2 (± 9.5)	57.5 (± 9.2)
Colorado	50.4 (± 6.7)	62.0 (± 5.6)
Connecticut	50.3 (± 9.2)	53.2 (± 9.0)
Delaware	60.1 (± 6.1)	63.1 (± 4.7)
District of Columbia	56.0 (±11.1)	65.1 (± 9.6)
Florida	58.0 (± 3.3)	54.0 (± 3.2)
Indiana	51.8 (± 9.6)	55.2 (± 7.8)
Iowa	54.9 (± 7.9)	46.4 (± 8.1)
Kansas	48.3 (± 8.2)	45.0 (± 9.9)
Kentucky	55.3 (± 9.1)	57.7 (± 9.6)
Maine	47.6 (±12.1)	51.4 (± 4.9)
Maryland	63.7 (± 6.0)	55.3 (± 4.4)
Minnesota	61.2 (± 6.1)	55.7 (± 4.7)
Mississippi — Private	56.7 (± 8.0)	55.2 (± 7.2)
Mississippi — Public	57.7 (± 7.4)	58.4 (±10.7)
New Hampshire	51.7 (± 7.4)	46.9 (± 8.6)
New York	61.6 (±10.0)	51.1 (±15.3)
Ohio	55.6 (± 5.3)	56.8 (± 9.2)
Tennessee	61.0 (± 2.6)	56.0 (± 3.7)
Texas	59.3 (± 6.6)	50.6 (± 7.5)
Vermont	46.4 (±12.0)	41.1 (± 8.3)
West Virginia	60.0 (± 7.3)	53.3 (± 7.1)
Wisconsin	54.1 (± 9.6)	48.2 (± 6.4)
Wyoming	59.3 (± 5.6)	58.3 (± 7.9)
Median	56.7	55.2
High school		
Alabama	59.2 (± 7.3)	61.8 (± 7.8)
Arkansas	61.5 (± 5.0)	56.9 (± 5.5)
California	71.6 (± 4.2)	59.2 (± 6.8)
Colorado	58.7 (± 5.9)	60.6 (± 8.4)
Connecticut	53.0 (± 7.7)	56.4 (± 6.9)
Delaware	59.7 (± 4.6)	62.0 (± 4.4)
District of Columbia	55.9 (± 8.8)	61.7 (±10.1)
Florida	58.2 (± 2.5)	57.6 (± 2.1)
Hawaii	71.0 (± 6.2)	71.8 (± 6.4)
Indiana	53.4 (± 6.1)	56.5 (± 5.0)
Iowa	57.7 (± 6.5)	51.4 (± 5.8)
Kansas	56.2 (± 5.2)	49.9 (± 5.1)
Kentucky	59.7 (± 3.4)	57.1 (± 5.1)
Maryland	55.7 (± 1.6)	52.9 (± 1.9)
Minnesota	61.0 (± 2.7)	61.0 (± 2.6)
Mississippi — Private	51.3 (± 7.8)	52.4 (± 9.0)
Mississippi — Public	64.9 (± 3.7)	59.7 (± 5.8)
Nebraska	54.3 (± 4.1)	53.6 (± 4.3)
New York	76.3 (± 3.7)	63.8 (± 5.3)
Ohio	60.7 (± 5.9)	59.7 (± 4.8)
South Dakota	60.8 (± 6.2)	55.0 (± 5.4)
Tennessee	57.9 (± 2.3)	55.3 (± 2.5)
Texas	71.6 (± 5.1)	54.7 (± 5.1)
West Virginia	59.4 (± 6.6)	56.1 (± 5.1)
Wisconsin	57.3 (± 6.0)	60.8 (± 5.1)
Median	59.2	57.1

* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

† Confidence interval.

TABLE 34. Percentage of middle school and high school students who were in a room or who rode in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days and who think smoke from other persons' cigarettes is harmful, by smoking status, sex, and race/ethnicity — National Youth Tobacco Survey, 2000

	Were in the same room with someone who was smoking cigarettes ≥ 1 days of preceding 7 days				Rode in a car with someone who was smoking cigarettes ≥ 1 days of preceding 7 days				Think smoke from other persons' cigarettes is harmful to you			
	Never cigarette smokers		Current cigarette smokers*		Never cigarette smokers		Current cigarette smokers		Never cigarette smokers	Current cigarette smokers		
	% (95% CI) [†]		% (95% CI)		% (95% CI)		% (95% CI)		% (95% CI)	% (95% CI)		
Middle school												
Sex												
Male	46.1	(± 2.5)	86.0	(± 2.9)	29.5	(± 2.1)	77.0	(± 3.2)	91.9	(± 1.3)	77.9	(± 4.4)
Female	51.4	(± 2.3)	91.2	(± 2.5)	34.1	(± 2.4)	84.5	(± 3.1)	93.8	(± 1.0)	82.4	(± 3.2)
Race/ethnicity												
White	52.8	(± 2.7)	92.3	(± 1.7)	33.3	(± 2.5)	83.6	(± 3.1)	95.6	(± 0.7)	81.6	(± 3.3)
Black	45.8	(± 3.4)	80.2	(± 5.6)	33.3	(± 3.1)	74.5	(± 4.4)	85.3	(± 2.8)	79.6	(± 5.8)
Hispanic	36.8	(± 3.6)	84.6	(± 5.1)	25.4	(± 2.6)	74.0	(± 5.6)	87.3	(± 2.6)	76.0	(± 4.6)
Asian	33.5	(± 6.2)	§		19.3	(± 5.1)	§		94.6	(± 2.1)	§	
Total (middle school)	48.9	(± 2.1)	88.4	(± 2.0)	31.9	(± 2.0)	80.6	(± 2.4)	92.9	(± 0.9)	80.0	(± 2.6)
High school												
Sex												
Male	53.4	(± 2.7)	90.3	(± 1.4)	28.8	(± 2.5)	80.2	(± 2.0)	93.0	(± 1.5)	86.9	(± 1.7)
Female	58.7	(± 2.7)	92.9	(± 1.1)	30.3	(± 2.7)	85.8	(± 1.8)	96.2	(± 0.8)	92.9	(± 1.2)
Race/ethnicity												
White	59.2	(± 3.1)	93.4	(± 1.0)	29.6	(± 3.1)	84.9	(± 1.5)	96.9	(± 0.7)	91.9	(± 1.1)
Black	52.5	(± 4.2)	85.1	(± 3.8)	34.0	(± 4.0)	77.7	(± 4.2)	87.3	(± 3.5)	79.7	(± 4.9)
Hispanic	44.6	(± 4.0)	86.0	(± 3.2)	25.0	(± 3.5)	74.6	(± 4.3)	90.8	(± 2.5)	85.4	(± 3.1)
Asian	49.9	(± 4.8)	84.9	(± 6.4)	24.3	(± 4.9)	75.4	(± 6.4)	95.6	(± 2.0)	89.4	(± 4.7)
Total (high school)	56.2	(± 2.3)	91.6	(± 1.0)	29.6	(± 2.3)	82.9	(± 1.5)	94.7	(± 1.0)	89.8	(± 1.3)

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

[†] Confidence interval.

[§] n<35.

TABLE 35. Percentage of middle school and high school students who were in a room or who rode in a car with someone who was smoking cigarettes on ≥ 1 days of the preceding 7 days and who think smoke from other persons' cigarettes is harmful, by smoking status and state — State Youth Tobacco Surveys, 2000

	Were in the same room with someone who was smoking ≥ 1 days of preceding 7 days		Rode in a car with someone who was smoking ≥ 1 days of preceding 7 days		Think smoke from other persons' cigarettes is harmful to you	
	Never smokers	Current smokers*	Never smokers	Current smokers	Never smokers	Current smokers
	% (95% CI) [†]	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school						
Alabama	52.8 (±4.4)	86.6 (±4.1)	41.1 (±4.5)	77.5 (± 5.2)	91.4 (±3.3)	84.6 (± 4.5)
Arizona	44.0 (±4.1)	82.5 (±5.1)	27.0 (±3.7)	74.3 (± 7.4)	91.5 (±2.3)	84.3 (± 5.6)
Arkansas	53.3 (±4.9)	86.0 (±5.0)	41.9 (±4.0)	80.6 (± 5.0)	91.8 (±2.2)	85.4 (± 4.1)
California	36.4 (±4.0)	81.8 (±5.6)	22.3 (±3.4)	67.1 (± 6.0)	89.0 (±4.1)	75.6 (± 5.4)
Colorado	40.2 (±4.0)	85.1 (±6.3)	27.5 (±3.5)	65.1 (± 9.0)	90.6 (±2.4)	85.9 (± 4.7)
Connecticut	45.6 (±3.8)	88.2 (±4.8)	29.4 (±3.1)	74.5 (± 7.5)	93.8 (±2.2)	79.8 (± 4.7)
Delaware	49.1 (±2.5)	90.5 (±3.2)	37.3 (±3.4)	85.1 (± 3.5)	93.1 (±1.4)	84.6 (± 3.4)
District of Columbia	37.3 (±4.1)	77.1 (±8.0)	21.7 (±2.7)	69.4 (±10.0)	82.7 (±3.1)	67.6 (±10.2)
Florida	51.6 (±1.5)	87.2 (±2.0)	35.2 (±1.6)	77.8 (± 2.5)	91.0 (±0.8)	84.1 (± 2.2)
Indiana	50.5 (±5.3)	88.2 (±4.8)	37.2 (±4.5)	79.4 (± 4.8)	93.2 (±2.0)	79.1 (± 6.8)
Iowa	50.7 (±3.0)	90.5 (±3.4)	34.3 (±4.2)	86.8 (± 4.2)	92.1 (±2.6)	82.4 (± 5.1)
Kansas	43.1 (±4.2)	82.2 (±9.2)	30.5 (±3.8)	78.0 (± 8.9)	91.9 (±1.6)	81.9 (±10.1)
Kentucky	62.3 (±3.5)	91.3 (±3.1)	42.5 (±3.6)	81.1 (± 6.2)	91.4 (±3.1)	83.6 (± 4.6)
Maine	47.6 (±5.3)	93.4 (±5.8)	34.0 (±5.3)	87.1 (± 8.3)	97.5 (±1.2)	84.5 (± 6.0)
Maryland	40.9 (±2.5)	77.1 (±4.1)	27.9 (±2.0)	72.7 (± 4.0)	89.4 (±1.1)	76.3 (± 5.3)
Minnesota	43.2 (±3.2)	86.5 (±3.6)	27.8 (±2.4)	80.8 (± 5.0)	93.3 (±1.4)	87.8 (± 3.7)
Mississippi — Private	54.4 (±7.0)	87.1 (±5.2)	31.9 (±4.2)	79.3 (± 6.2)	95.3 (±1.5)	81.8 (± 4.4)
Mississippi — Public	52.2 (±4.7)	80.2 (±4.0)	42.3 (±4.6)	80.0 (± 5.0)	91.2 (±3.2)	83.7 (± 4.7)
New Hampshire	48.1 (±3.7)	89.8 (±3.8)	29.4 (±3.8)	78.9 (± 5.2)	95.3 (±1.4)	88.5 (± 3.9)
New York	50.1 (±4.3)	81.4 (±5.8)	28.8 (±4.1)	63.0 (±12.1)	92.3 (±3.4)	80.7 (± 9.4)
Ohio	54.6 (±4.7)	88.1 (±5.0)	37.7 (±4.5)	82.5 (± 5.0)	94.3 (±1.2)	81.2 (± 6.9)
Tennessee	54.7 (±2.9)	90.1 (±1.8)	39.9 (±2.8)	82.9 (± 2.2)	92.4 (±1.5)	82.7 (± 2.0)
Texas	46.6 (±3.4)	85.8 (±6.7)	35.6 (±4.7)	79.1 (± 5.5)	90.5 (±3.2)	75.1 (± 7.1)
Vermont	47.8 (±5.0)	83.9 (±4.9)	34.5 (±4.6)	77.0 (± 4.5)	91.5 (±1.9)	81.6 (± 9.7)
West Virginia	63.7 (±3.7)	94.0 (±2.9)	45.5 (±4.2)	84.3 (± 5.9)	91.7 (±2.4)	84.5 (± 6.6)
Wisconsin	51.5 (±4.5)	90.3 (±5.1)	34.8 (±4.1)	78.1 (± 6.9)	94.7 (±2.5)	81.7 (± 5.1)
Wyoming	44.3 (±3.8)	91.5 (±4.2)	28.6 (±2.9)	81.1 (± 5.2)	95.9 (±1.5)	86.0 (± 4.7)
Median	49.1	87.1	34.3	79.1	91.9	82.7
High school						
Alabama	62.2 (±5.6)	90.8 (±3.5)	35.8 (±4.4)	82.9 (± 3.6)	95.1 (±2.3)	91.7 (± 2.9)
Arkansas	62.1 (±4.1)	92.0 (±3.2)	40.4 (±5.8)	88.7 (± 4.5)	93.7 (±3.6)	91.4 (± 2.5)
California	44.1 (±2.7)	84.2 (±4.3)	23.3 (±3.8)	74.6 (± 7.0)	94.0 (±2.1)	88.8 (± 4.2)
Colorado	49.2 (±4.2)	91.3 (±2.9)	19.9 (±4.6)	83.2 (± 2.7)	96.0 (±1.8)	91.4 (± 3.1)
Connecticut	55.8 (±4.9)	90.4 (±4.5)	29.2 (±4.4)	86.2 (± 4.9)	93.8 (±2.8)	92.0 (± 2.8)
Delaware	60.6 (±3.2)	94.2 (±1.8)	36.1 (±3.3)	88.3 (± 2.2)	97.2 (±1.0)	91.0 (± 2.4)
District of Columbia	39.7 (±4.9)	81.2 (±7.2)	25.2 (±3.9)	70.0 (± 7.5)	87.1 (±3.0)	76.2 (± 6.2)
Florida	56.5 (±2.0)	90.1 (±1.4)	31.2 (±2.2)	81.0 (± 1.7)	90.3 (±1.6)	89.7 (± 1.5)
Hawaii	54.8 (±4.2)	86.7 (±4.2)	28.6 (±4.5)	77.3 (± 6.3)	94.1 (±3.1)	89.9 (± 4.9)
Indiana	57.0 (±5.8)	92.8 (±2.5)	36.1 (±4.0)	88.5 (± 2.4)	95.8 (±1.8)	89.9 (± 2.5)
Iowa	58.1 (±7.6)	94.6 (±2.6)	27.8 (±6.0)	87.1 (± 5.0)	97.2 (±1.6)	92.7 (± 2.4)
Kansas	51.8 (±5.6)	92.8 (±2.7)	25.3 (±4.5)	86.0 (± 4.9)	95.1 (±2.0)	87.6 (± 3.4)
Kentucky	71.5 (±5.6)	94.6 (±2.2)	43.3 (±6.1)	86.8 (± 2.9)	94.9 (±3.1)	88.8 (± 3.0)
Maryland	49.7 (±1.5)	84.8 (±1.3)	28.8 (±1.3)	81.2 (± 1.3)	92.1 (±0.9)	84.0 (± 1.2)
Minnesota	53.5 (±2.6)	93.8 (±1.0)	25.4 (±2.2)	86.9 (± 1.5)	95.9 (±0.8)	91.4 (± 1.1)
Mississippi — Private	67.1 (±4.5)	93.3 (±2.5)	39.7 (±5.9)	86.1 (± 5.2)	96.1 (±1.8)	90.3 (± 3.2)
Mississippi — Public	58.6 (±4.5)	89.7 (±2.9)	39.8 (±5.2)	83.8 (± 3.4)	94.0 (±1.8)	88.3 (± 3.7)
Nebraska	54.2 (±3.2)	91.3 (±1.5)	28.4 (±3.2)	87.7 (± 3.1)	95.6 (±1.6)	91.2 (± 2.8)
New York	59.1 (±5.4)	89.9 (±4.4)	28.0 (±5.1)	77.5 (± 4.5)	91.7 (±3.8)	85.5 (± 4.3)
Ohio	63.7 (±7.3)	94.5 (±2.4)	39.9 (±8.4)	86.1 (± 3.0)	96.1 (±2.4)	91.6 (± 4.2)
South Dakota	52.2 (±4.9)	91.5 (±2.3)	27.4 (±3.9)	90.1 (± 2.3)	92.4 (±2.2)	89.2 (± 3.8)
Tennessee	64.0 (±3.0)	90.4 (±2.1)	37.2 (±2.6)	86.4 (± 2.2)	95.3 (±1.7)	87.8 (± 1.7)
Texas	50.5 (±4.3)	89.7 (±3.8)	27.1 (±3.1)	79.0 (± 8.2)	94.0 (±2.0)	89.6 (± 3.5)
West Virginia	66.5 (±6.0)	93.7 (±2.5)	34.1 (±4.4)	87.0 (± 3.0)	94.6 (±3.1)	88.1 (± 3.4)
Wisconsin	53.7 (±4.6)	95.0 (±1.9)	34.2 (±6.4)	91.3 (± 3.4)	95.6 (±2.0)	93.8 (± 3.0)
Median	56.5	91.3	29.2	86.1	94.9	89.9

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Confidence interval.

TABLE 36. Percentage of middle school and high school students who were exposed to tobacco use at home from either cigarettes or smokeless tobacco, by tobacco use status, sex, and race/ethnicity — National Youth Tobacco Survey, 2000

	Anyone else in home smokes cigarettes		Anyone else in home uses smokeless tobacco	
	Never cigarette smoker	Current cigarette smoker*	Never smokeless tobacco user	Current smokeless tobacco user†
	% (95% CI‡)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school				
Sex				
Male	31.5 (±2.2)	67.1 (±3.3)	6.9 (±1.3)	47.1 (± 4.8)
Female	33.2 (±2.0)	73.4 (±3.8)	7.8 (±1.3)	50.5 (±10.1)
Race/ethnicity				
White	32.3 (±2.3)	73.8 (±3.7)	8.8 (±1.5)	49.3 (± 5.2)
Black	36.5 (±3.1)	63.7 (±5.7)	5.2 (±1.5)	44.4 (±14.7)
Hispanic	28.7 (±2.1)	62.0 (±5.0)	3.6 (±0.7)	41.9 (±11.8)
Asian	30.4 (±6.1)	¶	3.2 (±1.7)	¶
Total (middle school)	32.4 (±1.8)	70.1 (±2.6)	7.4 (±1.2)	47.7 (± 4.4)
High school				
Sex				
Male	27.1 (±2.5)	54.4 (±2.7)	5.2 (±0.7)	31.3 (± 3.8)
Female	27.9 (±2.1)	59.1 (±2.7)	7.8 (±1.1)	36.2 (± 9.0)
Race/ethnicity				
White	26.9 (±2.5)	56.7 (±2.5)	7.7 (±1.1)	31.2 (± 3.8)
Black	30.9 (±3.5)	59.4 (±5.0)	5.5 (±1.2)	33.5 (±13.7)
Hispanic	25.5 (±3.2)	54.8 (±4.5)	3.9 (±1.0)	38.0 (±10.3)
Asian	27.3 (±4.9)	46.1 (±8.3)	3.4 (±1.5)	¶
Total (high school)	27.5 (±2.0)	56.7 (±2.3)	6.7 (±0.8)	31.8 (± 3.5)

* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.† Used smokeless tobacco on ≥ 1 of the 30 days preceding the survey.

‡ Confidence interval.

¶ n<35.

TABLE 37. Percentage of middle school and high school students who were exposed to tobacco use at home from either cigarettes or smokeless tobacco, by tobacco use status and state — State Youth Tobacco Surveys, 2000

	Anyone else in home smokes cigarettes		Anyone else in home uses smokeless tobacco	
	Never cigarette smoker	Current cigarette smoker*	Never smokeless tobacco user	Current smokeless tobacco user†
	% (95% CI‡)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school				
Alabama	38.5 (±4.9)	64.6 (± 8.5)	13.3 (±2.8)	34.7 (±11.1)
Arizona	32.1 (±3.3)	58.1 (± 9.2)	10.4 (±1.8)	34.8 (±11.1)
Arkansas	43.8 (±4.1)	72.1 (± 8.8)	21.4 (±2.5)	61.3 (±11.2)
California	25.6 (±3.0)	57.1 (± 9.6)	3.7 (±1.1)	43.9 (±11.6)
Colorado	31.8 (±4.8)	66.7 (± 8.7)	10.5 (±2.3)	¶
Connecticut	32.8 (±3.3)	61.2 (±10.3)	5.4 (±1.3)	43.9 (±16.5)
Delaware	40.1 (±3.3)	72.3 (± 3.6)	3.9 (±0.7)	29.1 (±12.0)
District of Columbia	41.6 (±4.1)	59.8 (± 9.4)	6.2 (±1.5)	¶
Florida	35.4 (±1.6)	68.0 (± 2.8)	5.8 (±0.6)	37.8 (± 5.4)
Indiana	36.4 (±5.6)	76.4 (± 8.4)	9.5 (±2.1)	44.1 (±19.8)
Iowa	34.1 (±4.5)	70.2 (± 8.6)	10.9 (±1.3)	44.6 (± 8.8)
Kansas	29.8 (±3.8)	66.0 (±12.3)	14.0 (±2.0)	57.4 (±28.1)
Kentucky	41.7 (±4.6)	76.1 (± 7.3)	14.3 (±2.6)	51.4 (± 8.6)
Maine	36.9 (±6.0)	77.5 (±10.5)	5.7 (±2.1)	46.0 (±24.8)
Maryland	34.5 (±2.7)	67.2 (± 5.0)	7.3 (±0.9)	41.4 (± 9.5)
Minnesota	31.3 (±3.6)	69.5 (± 5.9)	8.2 (±1.1)	49.9 (±13.3)
Mississippi — Private	28.0 (±4.0)	59.9 (± 7.3)	17.1 (±2.0)	45.4 (±11.2)
Mississippi — Public	39.8 (±4.9)	70.5 (± 5.9)	19.7 (±2.4)	47.8 (± 7.6)
New Hampshire	31.0 (±4.1)	66.2 (± 6.0)	5.7 (±1.5)	31.0 (±12.9)
New York	32.8 (±3.8)	64.2 (± 9.2)	3.7 (±1.2)	56.7 (±16.4)
Ohio	36.8 (±4.4)	75.9 (± 6.1)	8.8 (±1.9)	46.0 (±11.3)
Tennessee	38.5 (±3.3)	71.9 (± 3.3)	15.3 (±1.5)	47.7 (± 5.6)
Texas	34.4 (±4.4)	70.1 (± 5.6)	9.8 (±3.2)	45.4 (±12.5)
Vermont	36.8 (±5.0)	68.9 (± 4.6)	7.8 (±1.8)	37.2 (±18.9)
West Virginia	39.3 (±3.8)	76.8 (± 4.3)	23.9 (±2.9)	57.7 (± 9.9)
Wisconsin	35.7 (±4.3)	64.7 (± 5.2)	9.9 (±2.3)	39.0 (±16.7)
Wyoming	25.2 (±2.8)	71.3 (± 5.4)	19.6 (±2.7)	50.2 (± 8.2)
Median	35.4	68.9	9.8	45.4
High school				
Alabama	34.2 (±4.7)	58.0 (± 7.2)	12.3 (±3.5)	38.8 (± 9.1)
Arkansas	33.0 (±5.8)	62.1 (± 5.4)	16.6 (±2.9)	35.1 (± 7.2)
California	28.2 (±3.9)	50.3 (± 8.4)	4.1 (±1.1)	45.4 (±14.7)
Colorado	20.8 (±3.6)	49.7 (± 5.8)	9.3 (±1.6)	29.3 (± 6.4)
Connecticut	27.9 (±4.9)	54.2 (± 6.6)	4.4 (±1.5)	25.8 (±12.0)
Delaware	30.8 (±2.9)	64.9 (± 4.3)	3.9 (±0.8)	37.8 (±11.1)
District of Columbia	36.0 (±4.4)	60.2 (±10.5)	6.1 (±1.6)	¶
Florida	28.7 (±2.1)	54.9 (± 2.8)	4.6 (±0.6)	40.6 (± 5.6)
Hawaii	38.4 (±4.4)	54.0 (± 7.1)	5.4 (±1.3)	21.5 (±14.7)
Indiana	30.4 (±5.0)	60.0 (± 4.9)	8.0 (±1.8)	34.8 (±11.3)
Iowa	25.2 (±5.4)	53.5 (± 5.8)	12.6 (±1.8)	34.3 (± 9.7)
Kansas	23.4 (±4.2)	54.0 (± 6.7)	12.6 (±2.5)	36.4 (±10.6)
Kentucky	40.3 (±5.7)	63.2 (± 4.6)	13.0 (±2.6)	41.9 (± 9.2)
Maryland	30.6 (±1.6)	56.7 (± 1.7)	8.2 (±0.7)	38.1 (± 3.9)
Minnesota	27.2 (±2.1)	52.9 (± 2.4)	9.7 (±1.2)	33.8 (± 5.0)
Mississippi — Private	24.8 (±5.1)	43.0 (± 4.3)	18.2 (±3.0)	50.5 (± 9.6)
Mississippi — Public	40.5 (±5.5)	59.9 (± 4.2)	19.4 (±3.3)	50.0 (± 9.0)
Nebraska	24.2 (±3.1)	57.5 (± 2.7)	11.3 (±1.4)	42.6 (± 4.1)
New York	25.2 (±3.8)	55.0 (± 4.5)	5.0 (±1.0)	27.5 (±14.0)
Ohio	36.8 (±7.5)	64.8 (± 6.9)	8.8 (±2.6)	33.4 (±14.1)
South Dakota	27.4 (±3.8)	57.4 (± 7.0)	13.8 (±1.9)	42.6 (±13.1)
Tennessee	34.4 (±2.2)	62.0 (± 2.2)	13.8 (±1.8)	38.6 (± 4.5)
Texas	27.0 (±3.4)	54.6 (± 8.0)	7.2 (±2.9)	40.7 (± 9.6)
West Virginia	30.7 (±5.4)	57.5 (± 6.3)	22.7 (±2.8)	49.2 (± 7.3)
Wisconsin	28.5 (±8.1)	55.1 (± 5.8)	6.0 (±1.4)	31.3 (±12.1)
Median	28.7	56.7	9.3	38.1

* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

† Used smokeless tobacco on ≥1 of the 30 days preceding the survey.

‡ Confidence interval.

¶ n<35.

TABLE 38. Percentage of all middle school and high school students who practiced ways to say “No” to tobacco as part of school curriculum and who smoked cigarettes or used smokeless tobacco on school property during the 30 days preceding the survey, by sex and race/ethnicity — National Youth Tobacco Survey, 2000

	<u>Practiced ways to say “No” to tobacco</u> % (95% CI*)	<u>Smoked cigarettes on school property</u> % (95% CI)	<u>Used smokeless tobacco on school property</u> % (95% CI)
Middle school			
Sex			
Male	40.5 (±3.0)	3.6 (±0.6)	3.3 (±1.3)
Female	43.2 (±3.5)	2.4 (±0.5)	0.8 (±0.2)
Race/ethnicity			
White	41.7 (±4.0)	2.6 (±0.6)	2.0 (±0.9)
Black	43.8 (±3.4)	3.2 (±0.7)	1.6 (±0.6)
Hispanic	41.0 (±4.2)	3.6 (±0.8)	2.1 (±0.6)
Asian	40.6 (±5.8)	2.6 (±1.4)	1.1 (±0.9)
Total (middle school)	41.9 (±3.1)	3.0 (±0.5)	2.0 (±0.6)
High school			
Sex			
Male	13.2 (±1.3)	11.1 (±1.1)	7.3 (±1.2)
Female	16.3 (±1.5)	7.7 (±1.1)	0.6 (±0.3)
Race/ethnicity			
White	12.4 (±1.5)	10.4 (±1.3)	4.5 (±0.8)
Black	21.0 (±2.2)	5.7 (±1.5)	2.6 (±0.8)
Hispanic	18.5 (±1.9)	8.0 (±1.3)	3.1 (±0.9)
Asian	18.6 (±3.3)	9.8 (±2.6)	1.8 (±0.9)
Total (high school)	14.7 (±1.3)	9.4 (±0.9)	4.0 (±0.6)

* Confidence interval.

TABLE 39. Percentage of all middle school and high school students who practiced ways to say "No" to tobacco as part of school curriculum and percentage who smoked cigarettes or used smokeless tobacco on school property during the 30 days preceding the survey, by state — State Youth Tobacco Surveys, 2000

	Practiced ways to say "No" to tobacco	Smoked cigarettes on school property	Used smokeless tobacco on school property
	% (95% CI)*	% (95% CI)	% (95% CI)
Middle school			
Alabama	45.6 (± 6.1)	4.7 (±1.4)	3.9 (±1.7)
Arizona	41.9 (± 4.2)	5.1 (±1.2)	3.0 (±2.2)
Arkansas	42.0 (± 5.4)	4.9 (±1.1)	3.8 (±1.4)
California	39.7 (± 7.6)	2.1 (±0.8)	1.6 (±0.5)
Colorado	38.2 (± 4.9)	3.8 (±1.3)	2.1 (±0.8)
Connecticut	43.4 (± 6.4)	3.2 (±1.1)	2.0 (±0.9)
Delaware	39.9 (± 4.7)	4.6 (±0.9)	1.7 (±0.5)
District of Columbia	35.8 (± 3.8)	3.9 (±1.4)	2.2 (±1.2)
Florida	34.8 (± 2.3)	3.2 (±0.4)	1.7 (±0.3)
Indiana	33.4 (± 5.7)	2.3 (±1.1)	2.3 (±1.0)
Iowa	45.3 (± 6.6)	3.8 (±0.8)	2.1 (±0.9)
Kansas	32.6 (± 5.9)	1.7 (±0.8)	1.4 (±0.7)
Kentucky	43.4 (± 4.4)	6.3 (±1.9)	4.7 (±2.1)
Maine	54.9 (± 9.0)	5.2 (±3.6)	1.9 (±0.9)
Maryland	NA [†]	3.1 (±0.5)	2.0 (±0.6)
Minnesota	44.1 (± 6.6)	2.7 (±0.6)	1.7 (±0.4)
Mississippi — Private	22.1 (± 5.8)	2.2 (±1.2)	4.0 (±1.5)
Mississippi — Public	40.6 (± 3.6)	4.7 (±1.2)	2.8 (±1.2)
New Hampshire	29.3 (± 5.9)	4.2 (±1.3)	1.1 (±0.5)
New York	42.8 (±13.5)	2.0 (±1.2)	1.3 (±0.9)
Ohio	43.9 (± 6.3)	3.9 (±1.1)	2.8 (±1.2)
Tennessee	42.0 (± 3.4)	3.8 (±0.6)	3.0 (±0.7)
Texas	36.4 (± 5.4)	3.1 (±1.3)	2.1 (±0.8)
Vermont	43.5 (± 8.6)	4.2 (±1.0)	1.6 (±0.8)
West Virginia	42.7 (± 6.7)	4.8 (±1.1)	3.8 (±1.3)
Wisconsin	45.5 (± 6.2)	4.2 (±1.7)	2.9 (±1.2)
Wyoming	44.1 (± 4.7)	4.4 (±1.1)	5.1 (±1.7)
Median	42.0	3.9	2.1
High school			
Alabama	18.6 (± 3.6)	9.2 (±1.9)	6.0 (±1.2)
Arkansas	14.8 (± 2.8)	9.7 (±1.9)	7.5 (±2.4)
California	19.1 (± 5.1)	6.4 (±1.4)	2.6 (±1.1)
Colorado	11.3 (± 2.3)	12.3 (±3.2)	5.7 (±2.0)
Connecticut	18.5 (± 3.4)	13.4 (±3.2)	4.2 (±1.3)
Delaware	16.1 (± 1.6)	12.5 (±1.8)	2.3 (±0.7)
District of Columbia	22.7 (± 2.7)	7.9 (±2.0)	3.6 (±1.8)
Florida	18.3 (± 1.8)	7.7 (±0.9)	3.3 (±0.5)
Hawaii	23.7 (± 4.1)	12.2 (±3.5)	3.3 (±2.0)
Indiana	10.9 (± 2.0)	10.5 (±2.0)	3.3 (±1.3)
Iowa	12.6 (± 3.2)	10.5 (±2.6)	5.1 (±1.5)
Kansas	12.5 (± 2.5)	8.2 (±2.0)	4.8 (±1.6)
Kentucky	14.5 (± 1.9)	18.6 (±3.2)	9.1 (±2.2)
Maryland	NA [†]	10.9 (±0.6)	4.8 (±0.4)
Minnesota	16.1 (± 1.6)	11.5 (±1.3)	6.7 (±1.0)
Mississippi — Private	9.3 (± 1.6)	10.5 (±2.4)	11.9 (±2.8)
Mississippi — Public	23.6 (± 4.5)	9.9 (±2.1)	7.8 (±1.9)
Nebraska	12.1 (± 1.5)	9.9 (±1.9)	5.4 (±1.2)
New York	15.0 (± 2.7)	12.0 (±4.5)	4.3 (±1.8)
Ohio	14.5 (± 3.5)	12.1 (±1.9)	5.5 (±1.9)
South Dakota	15.4 (± 3.3)	10.7 (±3.1)	7.1 (±2.7)
Tennessee	14.9 (± 2.6)	12.2 (±1.5)	8.0 (±1.3)
Texas	13.4 (± 2.0)	7.4 (±2.6)	5.1 (±2.5)
West Virginia	20.5 (± 4.9)	17.3 (±3.7)	11.1 (±2.6)
Wisconsin	13.5 (± 3.3)	13.8 (±2.7)	4.6 (±1.8)
Median	15.0	10.7	5.1

* Confidence interval.

† Question was not asked.

State and Territorial Epidemiologists and Laboratory Directors

State and Territorial Epidemiologists and Laboratory Directors are acknowledged for their contributions to *CDC Surveillance Summaries*. The epidemiologists and the laboratory directors listed below were in the positions shown as of October 2001.

State/Territory	Epidemiologist	Laboratory Director
Alabama	John P. Lofgren, MD	William J. Callan, PhD
Alaska	John P. Middaugh, MD	Bernard Jilly, PhD
Arizona	Victorio J. Vaz, DVM, PhD (Acting)	Wesley B. Press, MS
Arkansas	Thomas C. McChesney, DVM	Michael G. Foreman
California	Duc Vugia, MD, MPH	Paul Kimsey, PhD
Colorado	Lisa Miller, MD, SCPH (Acting)	Ronald L. Cada, DrPH
Connecticut	James L. Hadler, MD, MPH	Katherine Kelley, DrPH
Delaware	A. LeRoy Hathcock, PhD	Jane Getchall, DrPH
District of Columbia	Martin E. Levy, MD, MPH	Maurice Knuckles, PhD (Acting)
Florida	Steven T. Wiersma, MD, MPH	Ming S. Chan, PhD
Georgia	Paul Blake, MD, MPH	Elizabeth A. Franko, DrPH
Hawaii	Paul V. Effler, MD, MPH	Vernon K. Miyamoto, PhD
Idaho	Christine G. Hahn, MD	Richard H. Hudson, PhD
Illinois	Mark Dworkin, MD, MPH	David L. Maserang, PhD
Indiana	Robert Teclaw, DVM, PhD, MPH	David E. Nauth
Iowa	M. Patricia Quinlisk, MD, MPH	Mary J. R. Gilchrist, PhD
Kansas	Gianfranco Pezzino, MD, MPH	Duane Boline, PhD
Kentucky	Glyn G. Caldwell, MD	Samuel B. Gregorio, DrPH
Louisiana	Raoult Ratard, MD, MPH	Henry B. Bradford, Jr, PhD
Maine	Kathleen F. Gensheimer, MD, MPH	John A. Krueger
Maryland	Ross Brechner, MD, MPH	J. Mehsen Joseph, PhD
Massachusetts	Alfred DeMaria, Jr, MD	Ralph J. Timperi, MPH
Michigan	Matthew L. Boulton, MD, MPH	Frances Pouch Downes, DrPH
Minnesota	Harry Hull, MD	Norman Crouch, PhD
Mississippi	Mary Currier, MD, MPH	Joe O. Graves, PhD
Missouri	Eduardo J. Simoes, MD, MPH, MSPH	Eric C. Blank, DrPH
Montana	Todd A. Damrow, PhD, MPH	Mike Spence, MD
Nebraska	Thomas J. Safranek, MD	Steve Hinrichs, MD
Nevada	Randall L. Todd, DrPH	L. Dee Brown, MD, MPH
New Hampshire	Jesse Greenblatt, MD, MPH	Veronica C. Malmberg, MSN
New Jersey	Eddy A. Bresnitz, MD, MS	S. I. Shahied, PhD
New Mexico	C. Mack Sewell, DrPH, MS	David E. Mills, PhD
New York City	Benjamin A. Mojica, MD, MPH	Alex Ramon, MD, MPH
New York State	Perry F. Smith, MD	Lawrence S. Sturman, MD, PhD
North Carolina	J. Steve Cline, DDS, MPH	Lou F. Turner, DrPH, MPH
North Dakota	Larry A. Shireley, MPH, MS	Bonna R. Cunningham, MS
Ohio	Forrest W. Smith, MD	William Becker, DO
Oklahoma	J. Michael Crutcher, MD, MPH	John Hitz, DrPH
Oregon	Melvin Kohn, MD, MPH	Michael R. Skeels, PhD, MPH
Pennsylvania	James T. Rankin, Jr, DVM, PhD, MPH	Bruce Kleger, DrPH
Rhode Island	Utpala Bandyopadhyay, MD, MPH	Gregory Hayes, DrPH
South Carolina	James J. Gibson, MD, MPH	Harold Dowda, PhD
South Dakota	Sarah L. Patrick, PhD, MPH	Michael Smith
Tennessee	Allen S. Craig, MD	Michael W. Kimberly, DrPH
Texas	Dennis Perrotta, PhD, CIC	Susan Neill, PhD, MBA
Utah	Robert Rolfs, MD (Acting)	Charles D. Brokopp, DrPH
Vermont	Ann R. Fingar, MD, MPH	Burton W. Wilcke, Jr, PhD
Virginia	Robert B. Stroube, MD, MPH	James L. Pearson, DrPH
Washington	Juliet VanEenwyk, PhD	Romesh K. Gautam, PhD
West Virginia	Loretta E. Haddy, MS, MA	Andrea Labik, PhD
Wisconsin	Jeffrey P. Davis, MD	Ronald H. Laessig, PhD
Wyoming	Karl Musgrave, DVM, MPH	Richard Harris, PhD
American Samoa	Joseph Tufa, DSM, MPH	Joseph Tufa, DSM, MPH
Federated States of Micronesia	Jean-Paul Chaine	—
Guam	Robert L. Haddock, DVM, MPH	Aurelto S. Espinola, MD
Marshall Islands	Tom D. Kijiner	—
Northern Mariana Islands	Jose L. Chong, MD	Joseph K.P. Villagomez
Palau	—	—
Puerto Rico	Carmen C. Deseda, MD, MPH	José Luis Molinaris, MD
Virgin Islands	Jose Poblete, MD (Acting)	Norbert Mantor, PhD

MMWR

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/Publications/mmwr>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (888) 232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.