




MMWRTM

Morbidity and Mortality Weekly Report

Surveillance Summaries

July 25, 2003 / Vol. 52 / No. SS-6

Trichinellosis Surveillance — United States, 1997–2001



The *MMWR* series of publications is published by the Epidemiology Program Office, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

General: Centers for Disease Control and Prevention. *Surveillance Summaries*, July 25, 2003. MMWR 2003;52(No. SS-6).

Specific: [Author(s)]. [Trichinellosis Surveillance — United States, 1997–2001]. In: *Surveillance Summaries*, July 25, 2003. MMWR 2003;52 (No. SS-6):[inclusive page numbers].

Centers for Disease Control and Prevention

Julie L. Gerberding, M.D., M.P.H.
Director

Dixie E. Snider, Jr., M.D., M.P.H.
(Acting) Deputy Director for Public Health Science

Donna F. Stroup, Ph.D., M.Sc.
(Acting) Associate Director for Science

Epidemiology Program Office

Stephen B. Thacker, M.D., M.Sc.
Director

Division of Public Health Surveillance and Informatics

Daniel M. Sosin, M.D., M.P.H.
Director
Associate Editor, Surveillance Summaries

Office of Scientific and Health Communications

John W. Ward, M.D.
Director
Editor, MMWR Series

Suzanne M. Hewitt, M.P.A.
Managing Editor, MMWR Series

C. Kay Smith-Akin, M.Ed.
Lead Technical Writer/Editor

Teresa F. Rutledge
Project Editor

Beverly J. Holland
Lead Visual Information Specialist

Lynda G. Cupell
Malbea A. Heilman
Visual Information Specialists

Quang M. Doan
Erica R. Shaver
Information Technology Specialists

CONTENTS

Introduction	1
Methods	2
Results	3
Discussion	4
References	7

On the Cover: Number of reported trichinellosis cases, by year — United States, 1947–2001

Trichinellosis Surveillance — United States, 1997–2001

Sharon L. Roy, M.D.
Adriana S. Lopez, M.H.S.
Peter M. Schantz, V.M.D., Ph.D.
Division of Parasitic Diseases
National Center for Infectious Diseases

Abstract

Problem/Condition: Trichinellosis is a parasitic disease caused by tissue-dwelling roundworms of the species *Trichinella spiralis*. The organism is acquired by eating *Trichinella*-infected meat products. The disease has variable clinical manifestations, ranging from asymptomatic to fatal. In the United States, trichinellosis has caused hundreds of preventable cases of illness and occasional deaths. The national trichinellosis surveillance system has documented a steady decline in the reported incidence of this disease, as well as a change in its epidemiology.

Reporting Period Covered: This report summarizes surveillance data for trichinellosis in the United States for 1997–2001.

Description of System: Trichinellosis became a nationally reportable disease in 1966, but statistics have been kept on the disease since 1947. The national trichinellosis surveillance system is a passive system that relies on existing resources at the local, state, and federal levels. Cases are diagnosed based on clinical history with laboratory confirmation. Cases are reported weekly to CDC through the National Electronic Telecommunications System for Surveillance (NETSS). Detailed data regarding signs and symptoms, diagnostic tests, and food consumption are gathered by using a supplementary standardized surveillance form and are reported to CDC by fax or mail. This information is compared with NETSS data several times a year by CDC staff. Discrepancies are reviewed with the state health departments. The purpose of the surveillance system is to determine the incidence of trichinellosis, to maintain awareness of the disease, to monitor epidemiologic changes, to identify outbreaks, to guide prevention efforts, and to measure the effectiveness of those efforts.

Results: Although trichinellosis was associated historically with eating *Trichinella*-infected pork from domesticated sources, wild game meat was the most common source of infection during 1997–2001. During this 5-year period, 72 cases were reported to CDC. Of these, 31 (43%) cases were associated with eating wild game: 29 with bear meat, one with cougar meat, and one with wild boar meat. In comparison, only 12 (17%) cases were associated with eating commercial pork products, including four cases traced to a foreign source. Nine (13%) cases were associated with eating noncommercial pork from home-raised or direct-from-farm swine where U.S. commercial pork production industry standards and regulations do not apply.

Interpretations: The majority of the decline in reported trichinellosis cases is a result of improved observance of standards and regulations in the U.S. commercial pork industry, which has altered animal husbandry practices resulting in reduced *Trichinella* prevalence among swine.

Public Health Actions: Because of the change in epidemiology of trichinellosis and the continued occurrence of cases among consumers of wild game meat and noncommercial pork, more targeted public education is needed to further reduce the incidence of this disease.

Introduction

Trichinellosis is a parasitic disease caused by a roundworm of the *Trichinella* genus that infects carnivorous and omnivorous animals, including domestic swine and wild game (e.g., bears, cougars, and wild boars) (1). Humans become infected when they eat meat cooked inadequately from these animals containing larval cysts of *Trichinella* spp. Trichinellosis has a broad clinical presentation, ranging from asymptomatic to fatal. Initial symptoms of abdominal pain, vomiting, diarrhea,

constipation, and low-grade fever might occur within 1 week of ingestion of infected meat, as the larvae invade the bowel wall. These symptoms might be followed in approximately 1–8 weeks by facial edema, peripheral edema, headaches, intermittent fever, conjunctivitis, rashes, pruritus, cough, myalgia, and eosinophilia, as the larvae migrate through the tissues and encyst in the muscles (2,3). Fatal cases might result from neurologic and myocardial complications (3).

In the United States, the national trichinellosis surveillance system has documented a steady decline in the reported

"Learning is like rowing upstream; not to advance is to fall back."

Chinese Proverb

MMWR Continuing Education provides timely courses on public health and clinical topics that help you advance your clinical skills.

Review course descriptions, take exams, track results, and receive course certificates – all from your own computer, when and where your schedule allows.

MMWR CE
Log on. Sit ready. Advance.

cdc.gov/mmwr



incidence of this disease. During 1947–1951, a median of 393 cases (range: 327–487) was reported annually, including 57 trichinellosis-related deaths (4). During 1997–2001, the incidence decreased to a median of 12 cases annually (range: 11–23) and no reported deaths. No apparent change in the trichinellosis surveillance system can account for this decline. However, because the surveillance system is not designed to detect asymptomatic cases, the number of reported cases probably represents only a portion of the total number of infections.

Historically, pork products were the most commonly identified source of *Trichinella* infection in the United States. However, the number of annual reported cases attributed to pork has declined for many years, largely because of changes implemented by the U.S. pork industry that have resulted in reduced prevalence of *Trichinella* among domestic swine. This report presents data for 1997–2001 on trichinellosis and describes the changing epidemiology of this disease.

Methods

State health departments report new cases of trichinellosis by week to CDC through the National Electronic Telecommunications System for Surveillance (NETSS) and through trichinellosis surveillance case report forms (CDC 54.7 Rev. 02-90) containing additional epidemiologic information. Additional cases might be identified through laboratory testing for *Trichinella* performed at CDC or through investigations conducted by CDC. NETSS data are reviewed several times a year and matched to case report forms or laboratory reports.

A confirmed case of trichinellosis is defined as one that is laboratory confirmed and is clinically compatible with the common signs and symptoms of trichinellosis among symptomatic persons. These signs and symptoms include eosinophilia, fever, myalgia, and periorbital edema. Laboratory confirmation is made either by identification of *Trichinella* larvae through muscle biopsy or by a positive serologic test. An outbreak of trichinellosis is defined as ≥ 2 cases, of which at least one must be laboratory confirmed. If associated patients share an epidemiologically implicated meal or meat product and have either positive serologic tests or clinically compatible illnesses, they should also be reported as confirmed (5).

Cases reported by states through NETSS that were unaccompanied by case report forms or that did not meet the case definition for a confirmed case were excluded from this report. During 1997–2001, only two cases were unaccompanied by case report forms and excluded. For data analysis, commercial pork products were defined as pork purchased at supermarkets, butcher shops, or public eating places. This category

includes pork products that were eaten or obtained in other countries. Noncommercial pork products were defined as pork purchased directly from a farm, obtained from home-raised swine, or obtained from wild boars (6).

Results

During 1997–2001, a total of 18 states reported 72 cases of trichinellosis to CDC, including nine outbreaks involving ≥ 2 persons. States in the Midwest and Mountain West accounted for 63 (88%) cases, of which 50 (69%) were reported from five states: California (21), Ohio (nine), Illinois (seven), Montana (seven), and Alaska (six) (Table 1). The nine common-source outbreaks occurred in these five states and accounted for 33 (46%) cases (Table 2). Of the 72 cases, 52 (72%) occurred among males. The median age of patients was 45 years (range: 3–81 years) (Figure 1).

During 1997–2001, a total of 55 (76%) of the 72 patients had at least one common sign or symptom of trichinellosis; 46 (64%) had myalgia, 39 (54%) had eosinophilia, 33 (46%) had fever, and 22 (31%) had periorbital edema. A total of 40 (56%) patients reported more than one sign or symptom. Of the 17 patients who did not report signs or symptoms, one was a girl aged 3 years with a positive *Trichinella* antibody titre but no eosinophilia, fever, or periorbital edema; myalgia could not be assessed. The signs and symptoms of the remaining 16 patients were unknown. For the 24 patients for whom the dates of consumption of the implicated meat and the onset of symptoms were available, the median incubation period was 13 days (range: 1–50 days).


TABLE 1. Number of trichinellosis cases and outbreak cases, by reporting states — United States, 1997–2001

State	1997	1998	1999	2000	2001	Total	Outbreak cases
Alaska	0	0	0	5	1	6	4
California	1	3	2	0	15	21	12
Connecticut	0	0	1	0	0	1	0
Florida	0	0	1	0	0	1	0
Hawaii	0	0	0	1	0	1	0
Iowa	0	0	0	0	2	2	0
Illinois	0	1	3	2	1	7	4
Indiana	1	0	0	0	0	1	0
Kansas	0	0	1	0	0	1	0
Michigan	1	0	0	2	0	3	0
Montana	5	2	0	0	0	7	5
New Jersey	2	0	0	0	0	2	0
New York	0	0	2	0	2	4	0
Ohio	1	8	0	0	0	9	8
Vermont	0	1	0	0	0	1	0
Washington	0	0	0	1	0	1	0
Wisconsin	1	0	1	0	1	3	0
Wyoming	0	0	0	0	1	1	0
Total	12	15	11	11	23	72	33

TABLE 2. Outbreaks of trichinellosis among patients, by year, state, number of cases, month of illness onset, and implicated meat — United States, 1997–2001

Year	State	No. of cases	Month of illness onset	Implicated meat
1997	Montana	5	December	Bear jerky
1998	Ohio	8	October–November	Bear roast and ground bear meat
1999	Illinois	2	March–May	Pork sausage and pork jerky
2000	Illinois	2	January	Pork sausage and smoked pork
2000	Alaska	4	August–September	Bear steak (fried)
2001	California	2	May	Home-raised pork
2001	California	6	May–June	Home-raised pork (raw)
2001	California	2	August	Bear
2001	California	2	November	Bear
Total		33		

FIGURE 1. Cumulative number* of patients with trichinellosis, by sex and age group — United States, 1997–2001



* N = 66. Age was unknown for six patients.

Of the 72 patients, 54 (75%) had serologic testing performed; 50 (93%) were seropositive for *Trichinella* antibodies. Muscle biopsies were performed on four (6%) patients; all were positive.

The source of infection was known or suspected for 57 (79%) patients. Pork products were associated with 22 (39%) cases: 12 with commercial pork, nine with home-raised or direct-from-farm swine, and one with a wild boar. Of the 12 cases associated with commercial pork, eight were linked to U.S. commercial pork, and four were linked to pork obtained in Egypt, Vietnam, and Yugoslavia. Nonpork products were associated with 30 (53%) cases: 29 with bear meat and one with cougar meat. The source of infection was unknown for 20 (35%) cases; these included five cases that were associated with both pork and nonpork products. Because a single type of meat could not be implicated, each patient was classified as having an unknown source of infection. None of the meats consumed by these five patients were examined for larvae (Table 3).

TABLE 3. Number and percentage of trichinellosis cases, by type and source of implicated meat product — United States, 1997–2001

Meat type/Source	No.	(%)
Pork products	22	(31)
Commercial	12	(17)
Sausage	4	(6)
Chops	1	(1)
Ham	1	(1)
Pickled pigs feet	1	(1)
Sausage and chops	1	(1)
Sausage and smoked pork	2	(3)
Unspecified	2	(3)
Direct from farm	9	(13)
Hunted (wild boar)	1	(1)
Nonpork products	30	(42)
Bear meat	29	(40)
Cougar meat	1	(1)
Unknown	20	(28)
No information	15	(21)
Both pork and nonpork	5	(7)
Commercial pork chops, bacon, hamburger	1	(1)
Commercial pork roast, ham, bacon, moose/quail/pheasant meat	1	(1)
Commercial pork roast, hamburger	1	(1)
Commercial pork sausage, ham, hamburger	1	(1)
Commercial pork sausage, sliced beef	1	(1)
Total	72	(100)*

* Might not total 100% because of rounding.

During 1997–2001, implicated meat was examined for *Trichinella* larvae in 34 cases, which included cases from seven of the nine common-source outbreaks. Larvae were present in meat associated with 32 of the 34 cases, including all seven of the outbreaks for which meat was examined. These outbreaks accounted for 29 of the 32 cases. Five outbreaks were linked to larvae-positive bear meat (Montana 1997, five cases; Ohio 1998, eight cases; Alaska 2000, four cases; California 2001, two cases; and California 2001, two cases). Two outbreaks were linked to larvae-positive meat from home-raised swine (California 2001, six cases; California 2001, two cases). Three isolated cases also were associated with larvae-positive meat: two with bear and one with cougar.

The method of cooking the meat was identified in 38 (53%) of 72 cases: uncooked, 20 cases; open fire roasted or barbecued, 14 cases; fried, 13 cases; stewed, six cases; baked, two cases; and microwaved, one case. Certain cases were associated with multiple cooking methods.

The source of the meat was identified in 55 (76%) of 72 cases. Meat was purchased commercially in 12 cases: supermarket, nine cases; butcher shop, one case; restaurant, one case; and a county fair, one case. Meat was obtained non-commercially in 41 cases: directly from a farm or from a home-raised animal, nine cases (all pork); and hunted or trapped, 32 cases (29 bear meat, one cougar meat, one wild boar meat, and one moose/quail/pheasant meat). The meat was consumed

in another country in four cases: two in Yugoslavia (pork) and one each in Egypt (pork) and Vietnam (pork).


Of the nine trichinellosis outbreaks, five were associated with bear meat, two with home-raised swine, and two with commercial pork. In December 1997, an outbreak of five cases in Montana was associated with bear jerky. In October–December 1998, an outbreak of eight cases in Ohio was associated with meat from a bear killed in northern Ontario, Canada, in March 1998 and transported back to the United States. The meat was prepared as a roast, a spaghetti dish, and hamburgers and eaten on October 13, 1998, at a church barbeque. The hunter tested positive for *Trichinella* by muscle biopsy. Another member of the outbreak tested positive by enzyme immunoassay and bentonite flocculation antibody tests. The remaining six cases were linked epidemiologically through the church barbeque. In August–September 2000, an outbreak of four cases in Alaska was associated with steaks from a bear killed in Alaska in August 2000 and fried on a camp stove. In August 2001, an outbreak of two cases in California was associated with “medium-rare” barbecued meat from a bear killed in Alaska in June 2001. In November 2001, an outbreak of two cases in California was associated with uncooked bear meat. Of the two outbreaks associated with home-raised swine, one was associated with uncooked pork (California, May 2001, two cases), and the other was associated with a combination of fried, barbecued, stewed, and uncooked pork (California, May–June 2001, six cases). Of the two outbreaks associated with commercial pork, one was associated with homemade pork sausage and jerky (Illinois, March–May 1999, two cases), and the other was associated with sausage and smoked pork consumed in Yugoslavia (Illinois, January 2000, two cases).

In previous years, seasonal peaks in trichinellosis in the United States were observed in December and January, corresponding to consumption of homemade pork sausage during the Christmas holidays (6). However, during 1997–2001, no such pattern was observed for patients with known dates of illness onset in which commercial pork products were the only implicated meats (Figure 2). For cases in which noncommercial pork products were the only implicated meats, seasonal peaks occurred in May and June and were associated with home-raised swine (Figure 3). For cases in which nonpork products were the only implicated meats, seasonal peaks occurred during August–September and November–December and were associated with bear meat (Figure 4).

Discussion


The reported incidence of trichinellosis has declined substantially since the U.S. Public Health Service began keeping

FIGURE 2. Cumulative number* of patients with trichinellosis associated with eating commercial pork products, by month of illness onset — United States, 1997–2001




* N = 11.

FIGURE 3. Cumulative number* of patients with trichinellosis associated with eating noncommercial pork products, by month of illness onset — United States, 1997–2001



* N = 10.

FIGURE 4. Cumulative number* of patients with trichinellosis associated with eating nonpork products, by month of illness onset — United States, 1997–2001




* N = 20.

records on the disease in 1947 (Figure 5). The decrease in cases has mirrored the decline in prevalence of *Trichinella* in commercial pork products. In 1900, the trichinellosis prevalence in U.S. swine was estimated to be 1.41%. The estimated prevalence decreased to 0.125% during 1966–1970 and to 0.013% in 1995 (7). This decline reflects changes in the U.S. pork industry that have reduced exposure of domestic swine to *Trichinella*. Historically, the major sources of *Trichinella*-infected pork were swine fed garbage containing animal waste products. Trichinellosis prevalence among swine decreased after garbage cooking laws were passed to control vesicular exanthema (1953–1954) and hog cholera (1962) (8,9). In 1980, Congress passed the Federal Swine Health Protection Act (Public Law 96-468) that prohibits feeding potentially contaminated garbage to swine. Other risk factors for *Trichinella*-infected pork include consumption by swine of small mammals infected with *Trichinella*, including rodents, raccoons, skunks, and opossums, and cannibalism among swine within an infected herd (10). These risk factors are being addressed through industry-wide emphasis on 1) maintaining effective rodent-control programs; 2) controlling access of wildlife and pets to swine, facilities, and feed stuffs; 3) maintaining good hygiene in facilities; and 4) removing dead swine immediately when they are found (10,11). The United States Department of Agriculture (USDA), the National Pork Producers Council, and the pork processing industry have developed the *Trichinae* Herd Certification Program. This is a voluntary pre-harvest pork safety program that provides documentation of swine management practices to minimize *Trichinella* exposure. The goal of the program is to establish a system under which pork production facilities that follow good production practices might be certified as *Trichinella*-safe (12).

In addition to the reduction in *Trichinella* prevalence in commercial pork, processing methods also have contributed to the dramatic decline in human trichinellosis associated with pork products. Through the U.S. Code of Federal

FIGURE 5. Number of reported trichinellosis cases, by year — United States, 1947–2001




Regulations, USDA has stipulated specific cooking temperatures and times, freezing temperatures and times, and curing methods for processed pork products to control post-harvest human exposure to *Trichinella* (13). Increased public awareness regarding safe cooking and freezing of fresh pork also has reduced post-harvest exposure.

Although the number of reported human trichinellosis cases related to eating pork has decreased, since 1982, the number of reported cases related to eating nonpork products has remained constant. During 1997–2001, the number of cases associated with eating nonpork products surpassed that of pork for the first time in the United States (Figure 6).

Home-raised and direct-from-farm swine and pork from foreign sources are unaffected by regulations and standards followed by the U.S. commercial pork production industry that have helped to reduce *Trichinella* prevalence in commercial pork. Outbreaks of trichinellosis associated with noncommercial sources of pork have been reported previously (14,15) and continue to be reported. Trichinellosis cases associated

FIGURE 6. Number of trichinellosis cases, by source of infection, as reported in surveillance summaries — United States, 1975–2001



* Bailey TM, Schantz PM. Trends in the incidence and transmission patterns of trichinosis in humans in the United States: comparisons of the periods 1975–1981 and 1982–1986. *Rev Infect Dis* 1990;12:5–11.

† McAuley JB, Michelson MK, Schantz PM. Trichinellosis surveillance, United States, 1987–1990. In: *CDC Surveillance Summaries* (December). *MMWR* 1991;40(No. SS-3):35–42.

‡ Moorhead A, Grunenwald PE, Dietz VJ, Schantz PM. Trichinellosis in the United States, 1991–1996: declining but not gone. *Am J Trop Med Hyg* 1999;60:66–9.

e xplore.

Trekking through medical and scientific literature on the Web can sometimes be a long journey – but not at MMWR Online.

Whether you're researching past CDC reports or looking for the latest updates on important medical topics, MMWR Online quickly guides you to the information you need, free of charge.

Log on to cdc.gov/mmwr and boldly discover more.

know what matters.



with noncommercial pork now outnumber those cases associated with U.S. commercial pork, reflecting a changing risk pattern in pork consumption.

The emergence of wild game meat as the most common source of trichinellosis and the continued occurrence of trichinellosis among consumers of pork obtained from small farms or other countries suggest that educational messages concerning the risks for eating meat cooked improperly, especially from noncommercial sources, are not reaching persons at risk for trichinellosis. USDA recommends that consumers of fresh pork cook the meat to an internal temperature of 160°F (71°C) (16). *T. spiralis* larvae in pork are killed at lower temperatures (e.g., 140°F [60°C] for 2 minutes or 131°F [55°C]) for 6 minutes (17); however, USDA has recommended a higher temperature to allow for different cooking methods that might result in uneven temperature distributions throughout the meat (e.g., microwave cooking) (13). *Escherichia coli*, *Salmonella*, *Staphylococcus aureus*, and *Listeria monocytogenes*, other foodborne microorganisms that also can be found in pork, are destroyed by thorough cooking to an internal temperature of 160°F (71°C) (16).

Freezing kills *T. spiralis* larvae in pork. Pork <6 inches thick can be made safe if frozen to -20°F (-29°C) for 6 days, -10°F (-23°C) for 10 days, or 5°F (-17°C) for 20 days (18). However, freezing might not kill other species and types of *Trichinella* found in wild game. Infective *Trichinella* spp. larvae have been found in frozen bear meat: grizzly bear meat frozen at -4°–20.3°F (-20°– -6.5°C) for 27 months (19) and polar bear meat frozen at -0.4°F (-18°C) for approximately 24 months (20,21). Infective *Trichinella* spp. larvae also have been found in other wild carnivore meat frozen at 5°F (-15°C): marten frozen for 5 months, wolverine frozen for 6 months, and arctic fox frozen for 14 months (22). In addition, viable *Trichinella* spp. larvae were found in black bear meat that had been processed into ham and jerky by dry curing with a commercial salt mixture at a USDA-licensed establishment by using procedures similar to those used to prepare pork for human consumption. However, no viable *Trichinella* spp. larvae were found in ground bear meat preparations (e.g., sausage, pepperoni, or salami) that had been processed according to standards mandated by USDA for processing pork (19).

The dramatic decline of trichinellosis in the United States reflects changes in industrial practices, increased government regulations, and increased public awareness. As a result, the epidemiology of the disease has changed. Because of the successful reduction in *Trichinella* prevalence among swine in the U.S. commercial pork industry, the majority of cases of human trichinellosis are now associated with wild game meat, noncommercial pork, and foreign pork. Persons at risk need a

greater understanding of the changing risks for trichinellosis, of the methods for safe meat preparation, and of the limitations of those methods in certain circumstances, if further reduction in the incidence of this disease is to be achieved in the United States.

References

1. Gamble HR, Brady RC, Bulaga LL, et al. Prevalence and risk association for *Trichinella* infection in domestic pigs in northeastern United States. *Vet Parasitol* 1999;82:59–69.
2. Schantz PM, Michelson MK. Trichinosis. In: Gorbach SL, Bartlett JG, Blacklow NR, eds. *Infectious Diseases*, 2nd ed. Philadelphia, PA: W.B. Saunders Company, 1998:616–20.
3. Murrell KD. Trichinosis. In: Strickland GT, ed. *Hunter's Tropical Medicine and Emerging Infectious Diseases*, 8th ed. Philadelphia, PA: W.B. Saunders Company, 2000:780–7.
4. Schantz PM. Trichinellosis in the United States—1947–1981. *Food Technol* 1983;37:83–6.
5. CDC. Case definitions for infectious conditions under public health surveillance. *MMWR* 1997;46(No. RR-10).
6. Bailey TM, Schantz PM. Trichinosis surveillance, United States, 1986. In: CDC Surveillance Summaries (December 1). *MMWR* 1988;37(No. SS-5):1–8.
7. Gamble HR, Bush E. Seroprevalence of *Trichinella* infection in domestic swine based on the National Animal Health Monitoring System's 1990 and 1995 swine surveys. *Vet Parasitol* 1999;80:303–10.
8. Zimmermann WJ, Zinter DE. The prevalence of trichiniasis in swine in the United States, 1966–70. *HSMHA Health Rep* 1971;86:937–45.
9. Zimmermann WJ, Steele JH, Kagan IG. Trichiniasis in the U.S. population, 1966–70: Prevalence and epidemiologic factors. *Health Serv Rep* 1973;88:606–23.
10. Gamble HR. *Trichinae*: pork facts—food quality and safety. Beltsville, MD: United States Department of Agriculture Animal and Plant Health Inspection Service, 2001. Available at http://www.aphis.usda.gov/vs/Trichinella/docs/fact_sheet.htm.
11. National Pork Board. *Quality Assurance: A Program of America's Pork Producers, Level III*. Des Moines, Iowa: National Pork Board, 1997: 47–52.
12. National Pork Board. *Trichinae Herd Certification*. Des Moines, Iowa: National Pork Producers Council, 2000. Available at <http://www.aphis.usda.gov/vs/Trichinella/>.
13. Gamble HR. Parasites associated with pork and pork products. *Rev Sci Tech* 1997;16:496–506.
14. McAuley JB, Michelson MK, Schantz PM. Trichinellosis surveillance, United States, 1987–1990. In: CDC Surveillance Summaries (December). *MMWR* 1991;40(No. SS-3):35–42.
15. McAuley JB, Michelson MK, Hightower AW, Engeran S, Wintermeyer LA, Schantz PM. A trichinosis outbreak among Southeast Asian refugees. *Am J Epidemiol* 1992;135:1404–10.
16. Food Safety and Inspection Service, United States Department of Agriculture. *Safety of fresh pork...from farm to table*. Washington, DC: Food Safety and Inspection Service, United States Department of Agriculture, September 2002. Available at <http://www.fsis.usda.gov/OA/pubs/pork.htm>.
17. Kotula AW, Murrell KD, Acosta-Stein L, Lamb L, Douglass L. *Trichinella spiralis*: effect of high temperature on infectivity in pork. *Exp Parasitol* 1983;56:15–9.

18. Moorhead A, Grunenwald PE, Dietz VJ, Schantz PM. Trichinellosis in the United States, 1991–1996: declining but not gone. *Am J Trop Med Hyg* 1999;60:66–9.
 19. Worley DE, Seese FM, Espinosa RH, Sterner MC. Survival of sylvatic *Trichinella spiralis* isolates in frozen tissue and processed meat products. *J Am Vet Med Assoc* 1986;189:1047–9.
 20. Kjos-Hanssen B. Freeze-resistance of *Trichinella* cysts in polar bears from the high-arctic region of Norway (Svalbard). *Acta Vet Scand* 1983;24:244–6.
 21. Kjos-Hanssen B. *Trichinella* isolates from polar bears in Svalbard. Freeze resistance and infectivity in rats and swine. *Nord Vet Med* 1984;36:57–61.
 22. Chadee K, Dick TA. Designation and freezing resistance of isolates of *Trichinella spiralis* from wild carnivores. *J Wildl Dis* 1982;18:169–73.
-

trust·wor·thy: *adj*

('trəst-"wər-thē) 1 : worthy of belief

2 : capable of being depended upon;

see also *MMWR*.



know what matters.



The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/publications/mmwr>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone 888-232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

All *MMWR* references are available on the Internet at <http://www.cdc.gov/mmwr>. Use the search function to find specific articles.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.