



---

# MMWR<sup>TM</sup>

## Morbidity and Mortality Weekly Report

---

Surveillance Summaries

August 29, 2003 / Vol. 52 / No. SS-9

---

### **Assisted Reproductive Technology Surveillance — United States, 2000**

The *MMWR* series of publications is published by the Epidemiology Program Office, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

**SUGGESTED CITATION**

General: Centers for Disease Control and Prevention. *Surveillance Summaries*, August 29, 2003. MMWR 2003;52(No. SS-9).

Specific: [Author(s)]. [Title of particular article]. In: *Surveillance Summaries*, August 29, 2003. MMWR 2003;52(No. SS-9):[inclusive page numbers].

**Centers for Disease Control and Prevention**

Julie L. Gerberding, M.D., M.P.H.  
*Director*

Dixie E. Snider, Jr., M.D., M.P.H.  
*(Acting) Deputy Director for Public Health Science*

Donna F. Stroup, Ph.D., M.Sc.  
*(Acting) Associate Director for Science*

**Epidemiology Program Office**

Stephen B. Thacker, M.D., M.Sc.  
*Director*

**Division of Public Health Surveillance and Informatics**

Daniel M. Sosin, M.D., M.P.H.  
*Director*  
*Associate Editor, Surveillance Summaries*

**Office of Scientific and Health Communications**

John W. Ward, M.D.  
*Director*  
*Editor, MMWR Series*

Suzanne M. Hewitt, M.P.A.  
*Managing Editor, MMWR Series*

C. Kay Smith-Akin, M.Ed.  
*Lead Technical Writer/Editor*  
*Project Editor*

Beverly J. Holland  
*Lead Visual Information Specialist*

Lynda G. Cupell  
Malbea A. Heilman  
*Visual Information Specialists*

Quang M. Doan  
Erica R. Shaver  
*Information Technology Specialists*

**CONTENTS**

Introduction ..... 2  
 Methods ..... 2  
 Results ..... 4  
 Discussion ..... 6  
 Acknowledgments ..... 8  
 References ..... 8

## Assisted Reproductive Technology Surveillance — United States, 2000

Victoria C. Wright, M.P.H.  
Laura A. Schieve, Ph.D.  
Meredith A. Reynolds, Ph.D.  
Gary Jeng, Ph.D.

*Division of Reproductive Health  
National Center for Chronic Disease Prevention and Health Promotion*

### Abstract

**Problem/Condition:** In 1996, CDC initiated data collection regarding assisted reproductive technology (ART) procedures performed in the United States to determine medical center-specific pregnancy success rates, as mandated by the Fertility Clinic Success Rate and Certification Act (FCSRCA) (Public Law 102-493, October 24, 1992). ART includes fertility treatments in which both eggs and sperm are handled in the laboratory (i.e., in vitro fertilization and related procedures). Patients who undergo ART treatments are more likely to deliver multiple-birth infants than women who conceive naturally. Multiple births are associated with increased risk for mothers and infants (e.g., pregnancy complications, premature delivery, low-birthweight infants, and long-term disability among infants).

**Reporting Period Covered:** 2000.

**Description of System:** CDC contracts with a professional society, the Society for Assisted Reproductive Technology (SART), to obtain data from fertility medical centers located in the United States. Since 1997, CDC has compiled data related to ART procedures. The Assisted Reproductive Technology Surveillance System was initiated by CDC in collaboration with the American Society for Reproductive Medicine, the Society for Assisted Reproductive Technology, and RESOLVE: The National Infertility Association.

**Results:** In 2000, a total of 25,228 live-birth deliveries and 35,025 infants resulting from 99,629 ART procedures were reported to CDC from 383 medical centers that performed ART in the United States and U.S. territories. Nationally, 75,516 (76%) of ART treatments were freshly fertilized embryos using the patient's eggs; 13,312 (13%) were thawed embryos using the patient's eggs; 7,919 (8%) were freshly fertilized embryos from donor eggs; and 2,882 (3%) were thawed embryos from donor eggs. The national live-birth delivery per transfer rate was 30.8%. The five states that reported the highest number of ART procedures were California (13,194), New York (11,239), Massachusetts (8,041), Illinois (7,323), and New Jersey (5,506). These five states also reported the highest number of live-birth deliveries and infants born as a result of ART. Overall, 47% of women undergoing ART-transfer procedures using freshly fertilized embryos from their own eggs were aged <35 years; 23% were aged 35–37 years; 19% were aged 38–40 years; 7% were aged 41–42 years; and 4% were aged >42 years. Among ART treatments in which freshly fertilized embryos from the patient's eggs were used, substantial variation in patient age, infertility diagnoses, history of past infertility treatment, and past births was observed. Nationally, live-birth rates were highest for women aged <35 years (38%). The risk for a multiple-birth delivery was highest for women who underwent ART-transfer procedures using freshly fertilized embryos from either donor eggs (40%) or from their own eggs (35%). Among women who underwent ART-transfer procedures using freshly fertilized embryos from their own eggs, further variation by patient age and number of embryos transferred was observed. Of the 35,025 infants born, 44% were twins, and 9% were triplet and higher order multiples, for a total multiple-infant birth rate of 53%. Patient's residing in states with the highest number of live-birth deliveries also reported the highest number of infants born in multiple-birth deliveries.

**Interpretation:** Whether an ART procedure was successful (defined as resulting in a pregnancy and live-birth delivery) varied according to different patient and treatment factors. Patient factors included the age of the woman undergoing ART, whether she had previously given birth, whether she had previously undergone ART, and the infertility diagnosis of both the female and male partners. Treatment factors included whether eggs were from the patient or a woman serving as an egg donor, whether the embryos were freshly fertilized or previously frozen and thawed, how long the embryos were kept in culture, how many embryos were transferred, and whether various specialized treatment procedures were used in conjunction with ART. ART poses a major risk for multiple births. This risk varied according to the

patient's age, the type of ART procedure performed, and the number of embryos transferred. In addition, the increased risk for multiple births has a notable population impact in certain states.

**Public Health Actions:** As use of ART and ART success rates continue to increase, ART-related multiple births are an increasingly important public health problem nationally and in many states. The proportion of infants born through ART in 2000 that were multiple births (53%) was substantially higher than in the general U.S. population during the same period. Data in this report indicate a need to reduce multiple births associated with ART. Efforts should be made to limit the number of embryos transferred for patients undergoing ART. In addition, continued research and surveillance is key to understanding the effect of ART on maternal and child health.

## Introduction

For >2 decades, assisted reproductive technologies (ARTs) have been used by couples to overcome infertility. ARTs include those infertility treatments in which both eggs and sperm are handled in the laboratory (i.e., in vitro fertilization and related procedures). Since the birth of the first U.S. infant conceived with ART in 1981, use of these treatments has increased dramatically. Each year, both the number of medical centers providing ART services and the total number of procedures performed have increased notably (1).

In 1992, Congress passed the Fertility Clinic Success Rate and Certification Act (FCSRCA),\* which requires each medical center in the United States that performs ART to report data to CDC annually and to include every ART procedure initiated. CDC uses the data to report medical center-specific pregnancy success rates. In 1997, CDC published the first surveillance report under this mandate (2). That report was based on ART procedures performed in 1995. Since then, CDC has continued to publish a surveillance report annually that details each medical center's success rates. CDC has also used this surveillance data file to perform more in-depth analyses of infant outcomes (e.g., multiple births) (3,4). Multiple-infant births are associated with greater health problems for both mothers and infants, including higher rates of caesarean deliveries, prematurity, low birthweight, and infant death and disability. This report is based on ART surveillance data provided to CDC's National Center for Chronic Disease Prevention and Health Promotion (NCCDPHP), Division of Reproductive Health, regarding procedures performed in 2000. A report of these data according to the medical center in which the procedure was performed was published separately (1). In this report, emphasis is on presenting state-specific data and presenting more detailed data regarding multiple-birth risk for 2000.

## Methods

Each year, the Society for Assisted Reproductive Technology (SART), an organization of ART providers affiliated with the American Society for Reproductive Medicine, collects data regarding ART procedures from medical centers performing ART in the United States and its territories and provides these data to CDC by contract. Data collected include patient demographics, medical history and infertility diagnoses, clinical information pertaining to the ART procedure, and information regarding resultant pregnancies and births. The data file is organized with one record per ART procedure performed. Multiple procedures from a single patient are not linked. Despite the federal mandate, certain centers (6%–7%/year) have not reported their data; the majority of these are believed to be smaller than average practices. For this report, data pertaining to ART procedures initiated January 1–December 31, 2000, are presented.

ART data and outcomes from ART are presented by state of residence. In cases of missing residency data (10%), the state of residency was assigned as the state in which the ART procedure was performed. Additionally, data regarding number of procedures are presented by treatment type and stage of treatment. ART procedures are usually classified into four groups according to whether a woman used her own eggs or received eggs from a donor and whether or not the embryos transferred were freshly fertilized or previously frozen and thawed. Because both success rates and multiple-birth risk vary substantially among these four treatments groups, data are presented separately for each type.

In addition to treatment types, within a given treatment procedure, different stages exist. A typical ART procedure begins when a woman starts taking drugs to stimulate egg production or begins having her ovaries monitored with the intent of having embryos transferred. If eggs are produced, the procedure progresses to the egg retrieval stage. After the eggs are retrieved, they are combined with sperm in the laboratory, and if fertilization is successful, the resulting embryos are selected for transfer. If the embryo implants in the uterus, the cycle progresses to a clinical pregnancy (i.e., the presence

\* Fertility Clinic Success Rate and Certification Act of 1992 (FCSRCA), Public Law 102-493, October 24, 1992.

of a gestational sac detectable by ultrasound). The resulting pregnancy might progress to a live-birth delivery. A live-birth delivery is defined as the delivery of  $\geq 1$  live-born infant. Only ART procedures involving freshly fertilized eggs include an egg retrieval stage; ART procedures using thawed eggs do not include egg retrieval because eggs were fertilized during a previous procedure and the resulting embryos were frozen until the current procedure. An ART procedure can be discontinued at any step for medical reasons or by the patient's choice.

Variations in a typical ART procedure are noteworthy. Although a typical ART procedure includes in vitro fertilization (IVF) of gametes, culture for  $\geq 2$  days and embryo transfer into the uterus (i.e., transcervical embryo transfer), in certain cases, unfertilized gametes (eggs and sperm) or zygotes (early embryos [i.e., a cell that results from fertilization of the egg by a sperm]) are transferred into the fallopian tubes within a day or two of retrieval. These are known as gamete and zygote intrafallopian transfer (GIFT and ZIFT). Another adaptation is intracytoplasmic sperm injection (ICSI) in which fertilization is still in vitro but is accomplished by selection of a single sperm that is injected directly into the egg. This technique was originally developed for couples with male factor infertility but is now commonly used for an array of diagnostic groups.

Detailed data are presented in this report for the most common treatment type, those using freshly fertilized embryos from the patient's eggs. These procedures account for  $>70\%$  of the total number of ART procedures performed each year. For those procedures that progressed to the embryo-transfer stage, percentage distribution of selected patient and treatment factors were calculated. In addition, success rates, defined as live-birth deliveries per ART-transfer procedure, were calculated according to the same patient and treatment characteristics.

Patient factors included the age of the woman undergoing ART, whether she had previously given birth, the number of past ART attempts, and the infertility diagnosis of both the female and male partners. The patient's age at the time of the ART procedure were grouped into five categories: aged  $<35$  years, 35–37 years, 38–40 years, 41–42 years, and  $>42$  years. Diagnoses ranged from one factor in one partner to multiple factors in one or both partners and were categorized as

- tubal factor — the woman's fallopian tubes are blocked or damaged, causing difficulty for the egg to be fertilized or for an embryo to travel to the uterus;
- ovulatory dysfunction — the ovaries are not producing eggs normally; such dysfunctions include polycystic ovarian syndrome and multiple ovarian cysts;
- diminished ovarian reserve — the ability of the ovary to produce eggs is reduced; reasons include congenital, medical, or surgical causes or advanced age ( $>40$  years);

- endometriosis — involves the presence of tissue similar to the uterine lining in abnormal locations; this condition can affect both fertilization of the egg and embryo implantation;
- uterine factor — a structural or functional disorder of the uterus that results in reduced fertility;
- male factor — a low sperm count or problems with sperm function that cause difficulty for a sperm to fertilize an egg under normal conditions;
- other causes of infertility — immunological problems or chromosomal abnormalities, cancer chemotherapy, or serious illnesses;
- unexplained cause — no cause of infertility was detected in either partner;
- multiple factors, female — diagnosis of  $>1$  female cause; or
- multiple factors, male and female — diagnosis of  $\geq 1$  female cause and male factor infertility.

Treatment factors included

- the number of days the embryo was cultured;
- the number of embryos that were transferred;
- whether the procedure was IVF-transfer only, IVF with ICSI, GIFT, ZIFT, or a combination of IVF with or without ICSI and either GIFT or ZIFT; and
- whether a woman other than the patient (a surrogate) gestated the pregnancy (i.e., a gestational carrier).

The number of embryos transferred in an ART procedure was categorized as 1, 2, 3, 4, or  $\geq 5$ . The number of days of embryo culture was calculated by using dates of egg retrieval and embryo transfer and was categorized as 1–6. However, because of limited sample sizes, live-birth rates are presented only for the two most common days, 3 and 5. Likewise, live-birth rates are presented for IVF with and without ICSI only and not for GIFT and ZIFT.

Multiple birth was assessed in two ways. First, each multiple-birth delivery was defined as a single event. A multiple-birth delivery was defined as the delivery of  $\geq 2$  infants in which at least one was live-born. The multiple-birth risk was thus calculated as the proportion of multiple-birth deliveries among total live-birth deliveries. Multiple birth was also assessed according to the proportion of infants from multiple deliveries among total infants (i.e., each infant was considered separately in this calculation). The proportion of live-born infants who were multiples (twins and triplets or more) was then calculated. Each of these measures represents a different focus. The multiple-birth risk, based on number of deliveries (or infant sets), provides an estimate of the individual risk posed by ART to the woman for multiple birth. The proportion of infants born in a multiple-birth delivery provides a measure of the



effect of ART treatments on children in the population. Both measures are presented by type of ART treatment and by maternal age for births conceived with the patient's eggs. Multiple-birth risk is further presented by number of embryos transferred. Proportion of infants born in a multiple-birth delivery is presented separately by state of residency. All analyses were performed by using SAS software system (5).

## Results

A total of 99,629 ART procedures performed in 2000 were reported to CDC (Table 1). The largest number of ART procedures occurred among patients who used their own freshly fertilized embryos (75,516; 76%). Of the 99,629 procedures started, 81,915 (82%) progressed to embryo transfer. Overall, 37% of ART procedures that progressed to the transfer stage resulted in a pregnancy, and 31% resulted in a live-birth delivery. Both pregnancy and live-birth rates varied according to type of ART. The highest rates were observed among ART procedures using donor eggs and freshly fertilized embryos (51% pregnancy rate and 44% live-birth rate). The lowest rates were observed among procedures using the patient's eggs and thawed embryos (26% pregnancy rate and 20% live-birth rate).

In all, the 25,228 live-birth deliveries from ART procedures resulted in 35,025 infants (Table 1); the number of infants born was higher than the number of live-birth deliveries because of multiple-infant births. The largest proportion of infants born (76.5%;  $n = 26,800$ ) were from ART procedures in which patients used freshly fertilized embryos from their own eggs.

Of 408 medical centers in the United States and surrounding territories that performed ART in 2000, a total of 383 provided data to CDC (Figure 1). The majority of medical centers that provided ART services were located in the eastern United States, in or near major cities. Within states, the number of medical centers performing ART was variable. Four states had no ART medical centers (Alaska, Maine, Montana, and Wyoming). States with the largest number of ART centers were California (56), Florida (28), New York (28), Illinois (25), and Texas (24).

The number of ART procedures performed among residents of each state approximately paralleled the data by medical center location (Table 2). The greatest numbers of ART procedures reported in 2000 were performed among residents of California (13,194), New York (11,239), Massachusetts (8,041), Illinois (7,323), and New Jersey (5,506). The five states with the largest number of ART procedures performed also reported the majority of live-birth deliveries and infants born. ART was used by residents of certain states and territo-

ries without an ART medical center (e.g., Alaska, Guam, Maine, Montana, Northern Mariana Islands, Virgin Islands, and Wyoming); however, each accounted for a limited percentage of total ART usage in the United States. Non-U.S. residents accounted for an estimated 1% of ART procedures, live-birth deliveries, and infants born.

Forty-seven percent of ART-transfer procedures using freshly fertilized embryos from the patient's eggs were performed on women aged <35 years; 23% on women aged 35–37 years; 19% on women aged 38–40 years; 7% on women aged 41–42; and 4% on women aged >42 years. Patient and treatment characteristics of these women varied by age (Table 3). The most common diagnoses reported for infertility among couples were male factor and tubal factor; however, diagnoses varied overall. Tubal factor, male factor, endometriosis, and ovulatory dysfunction were more commonly reported among younger women than women in older age categories. In contrast, diminished ovarian reserve was reported for only 1% of women aged <35 years; it was reported for 12% of women aged 41–42 years, and 22% of women aged >42 years. Among all women, 10%–12% were reported as having unexplained infertility; 11%–16% were reported as having multiple female factors; and 17%–18% were reported as having both male and female factors.

Approximately 60% of women aged <35 years were undergoing their first ART procedure. The percentage of women who had undergone at least one previous ART procedure increased with increasing age: only 40% of women aged >42 years were undergoing their first ART procedure. The percentage of women who had had a previous birth followed similar patterns. Although 19% of women aged <35 years reported at least one previous birth, this increased steadily with age: >35% of women in the oldest age group had had a previous birth.<sup>†</sup>

The majority of ART procedures used IVF with or without ICSI. Only a limited proportion of ART procedures used GIFT or ZIFT. Use of ICSI demonstrated a slight inverse relationship with patient age; these findings are consistent with higher rates of male factor infertility among younger age groups. In all age groups, the majority of procedures included embryo culture for 3 days; the next most common procedure involved embryo culture to day 5. Culture to day 5 coincides with development of the embryo to the blastocyst stage, which was used more frequently among younger women.

Although limited variation existed by age, the majority of ART procedures involved transfer of >1 embryo. Among women aged <35 years, 96% of procedures involved transfer

<sup>†</sup> Data were not available to distinguish whether previous births were conceived naturally or conceived with ART or other infertility treatments.

of  $\geq 2$  embryos, and 63% involved transfer of  $\geq 3$  embryos. For women aged  $>42$  years, 86% involved transfer of  $\geq 2$  embryos, and 68% involved transfer of  $\geq 3$  embryos. Use of a gestational carrier or surrogate was  $\leq 1\%$  for all age groups.

Live-birth rates for women who underwent ART procedures using freshly fertilized embryos from their own eggs also varied by patient age and selected patient and treatment factors (Table 4). Although the average live-birth rate for ART-transfer procedures performed among women who used their own freshly fertilized eggs was 32%, live-birth rates ranged from 38% among women aged  $<35$  years to 6% among women aged  $>42$  years (Table 4). Women aged  $\leq 40$  years who had an infertility diagnosis of ovulatory dysfunction or endometriosis or had unexplained infertility tended to have higher live-birth rates. Women aged  $>40$  years with a diagnosis of uterine factor had above average live-birth rates (22% for women aged 41–42 years and 15% for women aged  $>42$  years). Across all age groups, a diagnosis of diminished ovarian reserve was related to lower live-birth rates. Also, live-birth rates were lower for couples with a diagnosis of multiple factors, particularly male and female factors. Women aged  $\leq 42$  years who had undergone a previous ART procedure had lower live-birth rates than women undergoing their first ART procedure. However, the number of previous ART procedures cannot be subdivided by whether they were successful or not. Women aged  $\leq 42$  years who had had  $\geq 1$  previous birth had higher live-birth rates than those with no previous births. Women aged  $>42$  years had low live-birth rates overall (6%), regardless of whether they had had a previous ART or a previous birth.

ART procedures in which IVF was used with ICSI had lower live-birth rates in all age groups (Table 4). Live-birth rates were higher for women who had had extended embryo culture to day 5 than for women who had embryos cultured to day 3; this pattern was also observed for all age groups. In addition, among all age groups, the highest live-birth rates were observed among women who had transferred  $\geq 2$  embryos and among women who had used a gestation carrier. However, all of the results for treatment factors need to be considered cautiously because treatment was not randomized but rather based on medical center assessment and patient choice.

Of 8,806 multiple-birth deliveries, 6,784 were from pregnancies conceived with freshly fertilized embryos from the patient's eggs; 609 were from thawed embryos from the patient's eggs; 1,236 were from freshly fertilized embryos from a donor's eggs; and 177 were from thawed embryos from a donor's eggs (Table 5). In comparison with ART procedures using the patient's eggs and freshly fertilized embryos, the risks for multiple-birth delivery were increased when eggs from a donor were used and decreased when thawed embryos were

used. Among ART procedures in which the patient's own eggs were used, a strong inverse relation existed between multiple-birth risk and patient age. The average multiple-birth risk (i.e., multiple-birth delivery rate) for ART procedure in which freshly fertilized embryos from the patient's eggs were used was 35%. This rate varied from 39% among women aged  $<35$  years to 12% among women aged  $>42$  years.

Of 35,025 infants born through ART, 53% (18,603) were born in multiple-birth deliveries (Table 5). The proportion of infants born in a multiple-birth delivery also varied by type of ART procedure and patient age.

A more detailed examination of multiple-birth risk for women who underwent ART procedures in which freshly fertilized embryos from their own eggs were used revealed that number of embryos transferred was a risk factor for multiple-birth delivery, but the magnitude of the risk varied according to patient age (Figures 2–6). Among all age groups, transfer of  $\geq 2$  embryos resulted in increased live-birth delivery rates. However, the multiple-birth risk was also substantially increased. Among women aged  $<35$  years (Figure 2) and 35–37 years (Figure 3), the percentage of twin deliveries increased steadily with increasing number of embryos transferred from two to five or more. The percentage of triplet and higher order multiple-birth deliveries demonstrated an even more marked increase as the number of embryos transferred increased. As a result, the percentage of singleton deliveries for both of these two youngest age groups was actually lower when  $\geq 3$  embryos were transferred in comparison with two embryos being transferred. For women aged 38–40 years (Figure 4), transfer of  $\geq 3$  embryos offered a certain advantage in terms of live-birth delivery rates. However, as among younger age groups, the percentage of twin deliveries and triplet or higher order multiple-birth deliveries were substantially increased with  $\geq 3$  embryos having been transferred compared with two. For women aged 41–42 years (Figure 5), both the live-birth delivery rate and the multiple-birth risk increased steadily with an increased number of embryos having been transferred. The percentage of triplet or higher order multiple-birth deliveries did not demonstrate a trend. For women aged  $>42$  years (Figure 6), the percentage of twin deliveries (11%–13%) and triplet or higher order multiple-birth deliveries (0%–2%) did not vary substantially by number of embryos ( $\geq 2$ ) having been transferred.<sup>§</sup>

The total number and percentage of infants born in multiple-birth deliveries by maternal state of residency is presented

<sup>§</sup> Results are based on total multiple-birth risk and thus do not provide an indication of pregnancies that began as twins, triplets, or more, but reduced (either spontaneously or through medical intervention) to singletons or twins (see Figures 2–6).

(Table 6). Among states and territories with one or more total infants, conceived through ART, the numbers of infants born in multiple-birth deliveries ranged from 0 for residents of the Virgin Islands to 2,315 for California residents. The states with the highest number of ART-associated live-birth deliveries also had the highest number of infants born in multiple-birth deliveries. These include California (2,315), New York (1,946), Massachusetts (1,268), New Jersey (1,205), and Illinois (1,099). Nationally, the percentage of infants born in multiple-birth deliveries after ART was used was 53%; the percentage of twins and triplets or more were 44% and 9%, respectively. The percentage of infants born in multiple-birth deliveries was >50% in the majority of states. The states with the highest proportion of infants born in multiple-birth deliveries were Wyoming (69%), Nevada (62%), North Carolina (61%), Idaho (61%), and Oregon (60%).

## Discussion

According to the latest estimates of infertility in the United States from the 1995 National Survey of Family Growth, 15% of women of reproductive age reported a past infertility-associated health-care visit, and 2% reported a visit in the past year (6). Among married couples in which the woman was of reproductive age, 7% reported they had not conceived after 12 months of unprotected intercourse. With advances in ART, couples are increasingly turning to these treatments to overcome their infertility.

Since the birth of the first infant through ART in the United States in 1981, use of ART has grown substantially. Since 1997, CDC has been monitoring ART procedures performed in the United States. During that time, a notable and consistent increase in the use of ART has occurred. The increased use of ART coupled with increases in ART success rates has resulted in dramatic increases in the number of children conceived through ART each year. From 1996 (i.e., the first full year for which CDC collected data) through 2000, the number of ART procedures performed increased 54%, from 64,724 to 99,629 (7). Additionally, live-birth rates for all types of ART procedures increased from 11% to 22% during 1996–2000. The number of infants conceived through ART increased 67%, from 20,921 infants conceived through ART procedures performed in 1996 to 35,025 infants conceived through ART procedures performed in 2000.

This report documents that in 2000, ART use varied according to state of residency. Residents of California, New York, Massachusetts, Illinois, and New Jersey reported the highest number of ART procedures. These states also reported the highest number of infants conceived through ART. In

2000, ART use by state of residency was partly in line with expectations based on the total population size of women of reproductive age (i.e., women aged 15–44 years) within states (7). For example, California, New York, and Illinois were in the top five states for both ART usage and total number of women of reproductive age. However, ART use did not completely parallel population size. Whereas Massachusetts had the third highest number of ART procedures performed, it ranked thirteenth in terms of women of reproductive age. Likewise, New Jersey ranked fifth according to ART use, but ranked tenth according to population size. This divergence is not completely surprising because both Massachusetts and New Jersey had statewide mandates for insurance coverage for ART procedures in 2000. The state variation might also be related to availability of ART services within each state. However, we cannot disentangle the relation between demand for services and availability (i.e., increased availability in certain states might reflect the increased demand for ART among state residents).

Patients with an array of characteristics used ART services. Among ART treatments in which freshly fertilized embryos from the patient's eggs were used (i.e., the most frequent type of ART treatment), substantial variation was observed in patient age, infertility diagnoses, history of past infertility treatment, and past births.

Success rates from ART use are affected by numerous patient and treatment factors; hence, considering one single measure of success in evaluating ART efficacy is not informative. At a minimum, ART treatments need to be subdivided into general categories on the basis of the source of the egg (patient or donor) and the status of the embryos (freshly fertilized or thawed) because success rates vary substantially across these types. Within the type of ART treatment, further variation exists in success rates by patient and treatment factors, most notably patient age. Other factors to consider when assessing success rates are infertility diagnosis, number of previous ART procedures, number of previous births, type of ART procedure, number of days of embryo culture, number of embryos transferred, and use of a gestational carrier (surrogate). Variation exists in success rates according to each of these factors.

CDC's primary focus in collecting ART data has been live-birth deliveries as an indicator of success, because ART surveillance activities were developed in response to a federal mandate to report ART success rate data. Thus, a key role for CDC has been to publish standardized data related to ART success rates, including information regarding factors that affect these rates. With these data, couples can make informed decisions regarding whether to undergo this time-consuming



and expensive treatment (8,9).<sup>‡</sup> However, success-rate data must also be balanced with consideration of effects on maternal and infant health. Thus, CDC also closely monitors multiple births conceived through ART.

Multiple births are associated with an increased health risk for both mothers and infants (10–12). Women with multiple-gestation pregnancies are at increased risk for maternal complications (e.g., hemorrhage and hypertension). Infants born in a multiple-birth delivery are at increased risk for prematurity, low birthweight, infant mortality, and long-term disability. The health risks associated with multiple births have also contributed to rising health-care costs. In 2000, the estimated costs (per family) of delivering multiple-gestation pregnancies resulting from ART procedures ranged from \$58,865 for twins to \$281,698 for quadruplets (12). Additionally, a study regarding costs per delivery resulting from IVF pregnancies estimated \$39,000 for singleton or twin pregnancies and \$340,000 for triplet and quadruplet pregnancies (13).

In the United States, multiple births have increased dramatically during the last 2 decades (14,15). The rise in multiple births has been attributed to an increased use of ART and delayed childbearing (4,16,17). A recent study concerning trends in multiple births reported that 14% of multiple births in the United States in 2000 were attributed to ART; the proportions of twins and triplets or more attributed to ART were 12% and 43%, respectively (4). Other non-ART infertility treatments were estimated to also have contributed substantially to multiple births in the United States. Thus, even with shifts toward delayed child-bearing and the concomitant increase in multiple births known to be associated with advanced maternal age, by 2000, only 67% of twin births and 18% of triplet or higher order multiple births were estimated to have been naturally conceived.

In this report, 53% of infants born through ART in 2000 were multiple births; this compares with 3% in the general U.S. population during the same period (14,18). The twin rate was 44%, 22 times higher than in the general U.S. population (2%); the triplet and higher order multiples rate was 9%, 50 times higher than the general U.S. population (0.18%). Regarding the specific type of ART treatment, the rates are even higher for women who underwent ART procedures using freshly fertilized embryos from their own eggs (54% total multiple births) or from donor eggs (59% total multiple births). In the majority of states, >50% of infants conceived through ART were born in multiple-birth deliveries. North Carolina, Nevada, Idaho, Oregon, and Wyoming reported ART-associated multiple-birth rates >60%. Multiple births resulting from ART are an increasing public health problem,

nationally and for the majority of states. The findings in this report confirm the need to reduce the occurrence of multiple births resulting from ART.

This analysis was subject to certain limitations. First, ART surveillance data are reported for each ART procedure performed rather than for each patient who used ART. Linking procedures among patients who underwent >1 ART procedure in a given year is not possible. Because patients undergoing >1 procedure in a given year are most likely to be those who failed  $\geq 1$  treatment, the success rates reported here might underestimate the true per-patient success rate. Second, these data represent couples who sought ART services in 2000; therefore, success rates do not represent all couples with infertility who were potential ART users in 2000. Third, 6% of medical centers that performed ART in 2000 did not report their data to CDC as required. Fourth, comparisons among states are based on absolute numbers that are not adjusted for state population size.

ART data are reported to CDC by the ART medical center where the procedure was performed rather than by the state where the patient resided. In this report, we present ART data by the female patient's state of residence. In previous reports (16), we were unable to present ART data by state of residence because of incompleteness of residency data. In 2000, residency data were missing for 10% of all live-birth deliveries reported to CDC. The range of missing residency data varied by medical center. Medical centers located in 35 states had <5% missing residency data; medical centers located in six states had 5%–9% missing residency data; medical centers located in one state had 10%–15% missing residency data, and medical centers located in six states had >15% missing residency data. These states were Arizona, Georgia, Massachusetts, Minnesota, New York, and Pennsylvania. In cases of missing residency data, we assigned residency as the state in which the ART procedure was performed. Thus, the number of procedures performed among state residents, number of infants, and number of multiple-birth infants might have been overestimated for these states. Concurrently, the numbers might be underestimated in states bordering states with missing data, particularly states in the Northeast region of the United States. Nonetheless, the effects of missing residency data were not substantial. We separately evaluated statistics according to the state in which the ART medical center was located rather than the patient's state of residency. The rankings of the states in terms of total number of infants and multiple-birth infants were similar to the rankings based on state of residency (data not shown).

Despite these limitations, findings from national surveillance of ART procedures performed in the United States provide useful information for patients contemplating ART, ART

<sup>‡</sup> Estimated costs for one cycle of IVF range from \$7,000 to \$11,000 (8,9).

providers, and health-care policy makers. First, ART surveillance data can be used to monitor trends in ART use and outcomes from ART procedures. Second, data from ART surveillance can be used to assess patient and treatment factors that contribute to higher success rates. Third, ongoing surveillance data can be used to assess the risk of multiple births. Fourth, surveillance data provide information to assess changes in clinical practice related to ART treatment.

Multiple births are one of the most important public health concerns associated with using ART. Increased use of ART treatments and the widespread practice of transferring multiple embryos during ART treatments has led to a substantial increase in multiple-birth rates in the United States (4,15). Although balancing the chance of success with ART against the risk of multiple births is difficult in certain cases, efforts should be made to limit the number of embryos transferred for patients undergoing ART. Such efforts will ultimately require ART patients and providers to view treatment success in terms of singleton pregnancies and births. Additionally, continued research is critical to understanding the effect of ART on maternal and child health. CDC will continue to provide updates of ART use in the United States as data become available.

### Acknowledgments

The data used for this study were collected by the Society for Assisted Reproductive Technology (SART). The SART system is jointly supported by CDC, Atlanta, Georgia; SART, Birmingham, Alabama; and the American Society for Reproductive Medicine (ASRM), Birmingham, Alabama. The authors thank SART and ASRM, without whose contributions this work would not have been possible.

### References

1. CDC, American Society for Reproductive Medicine, Society for Assisted Reproductive Technology, and RESOLVE. 2000 assisted reproductive technology success rates. Atlanta, GA: US Department of Health and Human Services, CDC, 2002.
2. CDC, American Society for Reproductive Medicine, Society for Assisted Reproductive Technology, and RESOLVE. 1995 assisted reproductive technology success rates. Atlanta, GA: US Department of Health and Human Services, CDC, 1997.
3. Schieve LA, Peterson HB, Meikle SF, et al. Live-birth rates and multiple-birth risk using in vitro fertilization. *JAMA* 1999;282:1832–8.
4. Reynolds MA, Schieve LA, Martin JA, Jeng G, Macaluso M. Trends in multiple births conceived using assisted reproductive technology, United States, 1997–2000. *Pediatrics* 2003;111:1159–62.
5. SAS Institute, Inc. SAS/STAT® user's guide. Version 8. Cary, NC: SAS Institute Inc, 1999.
6. Abma JC, Chandra A, Mosher WD, Peterson LS, Piccinino LJ. Fertility, family planning, and women's health: new data from the 1995 National Survey of Family Growth. Hyattsville, MD: US Department of Health and Human Services, CDC, National Center for Health Statistics, 1997. (Vital and Health Statistics, vol 23).
7. US Census Bureau. Census 2000 summary file 1 (SF 1) 100-Percent Data. Sex by Age [49]—universe: total population. Washington, DC: US Census Bureau, 2000. Available at <http://factfinder.census.gov>.
8. Neumann PJ, Gharib SD, Weinstein MC. Cost of a successful delivery with in vitro fertilization. *N Engl J Med* 1994;331:239–43.
9. Collins J. Cost-effectiveness of in vitro fertilization. *Semin Reprod Med* 2001;19:279–89.
10. Schieve LA, Meikle SF, Ferre C, Peterson HB, Jeng G, Wilcox LS. Low and very low birth weight in infants conceived with use of assisted reproductive technology. *N Engl J Med* 2002;346:731–7.
11. Senat MV, Ancel PY, Bouvier-Colle MH, Breart G. How does multiple pregnancy affect maternal mortality and morbidity? *Clin Obstet Gynecol* 1998;41:78–83.
12. ESHRE Capri Workshop Group. Multiple gestation pregnancy. *Hum Reprod* 2000;15:1856–64.
13. Goldfarb JM, Austin C, Lisbona H, Peskin B, Clapp M. Cost-effectiveness of in vitro fertilization. *Obstet Gynecol* 1996; 87:18–21.
14. Martin JA, Hamilton BE, Ventura SJ, Menacker F, Park MM. Births: final data for 2000. *National Vital Statistics Reports* 2002;50(5):1–101.
15. Martin JA, Park MM. Trends in twin and triplet births: 1980–1997. *National Vital Statistics Reports* 1999;47(24):1–16.
16. CDC. Use of assisted reproductive technology—United States, 1996 and 1998. *MMWR* 2002;51:97–101.
17. Kiely JL, Kleinman JC, Kiely M. Triplets and higher-order multiple births: time trends and infant mortality. *Am J Dis Child* 1992;146:862–8.
18. MacDorman MF, Minino AM, Strobino DM, Guyer B. Annual summary of vital statistics—2001. *Pediatrics* 2002;110:1037–52.

**TABLE 1. Outcomes of assisted reproductive technology (ART), by procedure type — United States, 2000**

ART procedure type	No. of ART procedures started	No. of procedures progressing to retrievals	No. of procedures progressing to transfers	No. of pregnancies	Pregnancies per transfer procedure (%)	No. of live-birth deliveries	Live-birth deliveries per transfer procedure (%)	No. of infants born
Patient's eggs used								
Freshly fertilized embryos	75,516	64,784	60,780	23,258	38.3	19,219	31.6	26,800
Thawed embryos	13,312	N/A*	11,602	2,975	25.6	2,360	20.3	3,048
Donor eggs used								
Freshly fertilized embryos	7,919	7,243	6,989	3,566	51.0	3,041	43.5	4,382
Thawed embryos	2,882	N/A	2,544	758	29.8	608	23.9	795
<b>Total</b>	<b>99,629</b>	<b>—</b>	<b>81,915</b>	<b>30,557</b>	<b>37.3</b>	<b>25,228</b>	<b>30.8</b>	<b>35,025</b>

\* Data not available.

**TABLE 2. Number of reported assisted reproductive technology (ART) procedures performed, number of pregnancies, and number of live-birth deliveries, by patient's state/territory of residence\* at time of treatment — United States, 2000**

Patient's state/ territory of residence	No. of ART procedures started	No. of transfer procedures	No. of pregnancies	No. of live- birth deliveries	No. of infants born
Alabama	578	476	174	144	216
Alaska	73	66	34	31	44
Arizona	1,111	907	359	299	444
Arkansas	371	293	116	105	142
California	13,194	11,352	4,009	3,245	4,453
Colorado	1,588	1,424	689	599	863
Connecticut	1,947	1,582	591	486	679
Delaware	438	290	107	86	117
District of Columbia	380	320	128	109	157
Florida	4,168	3,318	1,302	1,100	1,534
Georgia	2,308	1,794	677	570	774
Guam	1	1	0	0	0
Hawaii	457	358	105	93	128
Idaho	293	252	123	107	162
Illinois	7,323	5,820	1,883	1,551	2,137
Indiana	1,940	1,543	527	427	597
Iowa	800	613	243	202	275
Kansas	707	549	218	187	255
Kentucky	722	615	221	182	247
Louisiana	722	516	176	132	189
Maine	90	79	35	29	37
Maryland	3,202	2,581	994	825	1,153
Massachusetts	8,041	6,817	2,425	1,950	2,609
Michigan	3,420	2,765	938	801	1,122
Minnesota	1,943	1,705	789	669	908
Mississippi	268	217	71	61	84
Missouri	1,271	1,071	407	345	491
Montana	94	81	35	28	39
Nebraska	468	391	125	104	138
Nevada	735	633	251	211	313
New Hampshire	428	364	133	103	145
New Jersey	5,506	4,277	1,821	1,507	2,139
New Mexico	228	199	91	76	107
New York	11,239	9,238	3,432	2,753	3,771
North Carolina	1,551	1,300	510	429	638
North Dakota	174	131	50	40	51
Northern Mariana Islands	1	1	0	0	0
Ohio	2,349	1,921	698	593	843
Oklahoma	527	438	187	158	220
Oregon	748	571	241	219	321
Pennsylvania	3,511	2,770	917	742	997
Puerto Rico	344	301	113	81	112
Rhode Island	720	644	147	123	174
South Carolina	657	562	228	196	283
South Dakota	105	80	28	24	32
Tennessee	603	490	244	211	287
Texas	5,249	4,322	1,711	1,414	1,978
Utah	446	371	145	126	171
Vermont	127	111	39	31	40
Virgin Islands	8	8	1	1	1
Virginia	2,406	1,988	813	665	922
Washington	1,846	1,481	541	460	639
West Virginia	210	172	61	49	68
Wisconsin	879	765	265	224	305
Wyoming	46	41	20	17	26
Non U.S. resident	1,068	940	369	308	448
<b>Total</b>	<b>99,629</b>	<b>81,915</b>	<b>30,557</b>	<b>25,228</b>	<b>35,025</b>

\* In cases of missing residency data, the patient's state/territory of residence was assigned as the state in which the ART procedure was performed. Medical centers in all but six states had missing residency data for <15% of infants born through ART. Medical centers located in Arizona, Georgia, Massachusetts, Minnesota, New York, and Pennsylvania had >15% missing residency data.

**TABLE 3. Percentage distribution of selected patient and treatment factors for assisted reproductive technology (ART) transfer procedures among patients who used freshly fertilized embryos from their own eggs, by patient age — United States, 2000**

	Patient age (yrs)				
	<35 (n = 28,778) (%)	35–37 (n = 14,146) (%)	38–40 (n = 11,301) (%)	41–42 (n = 4,365) (%)	>42 (n = 2,190) (%)
<b>Patient factors</b>					
Diagnosis					
Tubal factor	16.8	18.3	16.3	12.9	9.2
Ovulatory dysfunction	7.0	4.5	3.6	3.4	2.8
Diminished ovarian reserve	1.2	1.9	5.0	12.0	22.1
Endometriosis	9.2	8.1	6.5	4.5	3.1
Uterine factor	0.8	1.4	1.5	2.1	1.8
Male factor	23.3	19.9	16.7	10.5	8.7
Other causes	4.6	5.3	6.6	9.1	10.6
Unexplained cause	9.7	11.4	11.9	11.2	10.8
Multiple factors, female only	10.5	11.6	13.6	16.3	13.5
Multiple factors, female and male	17.0	17.6	18.3	18.1	17.5
Number of previous ART procedures					
0	59.8	50.0	45.9	43.8	39.8
≥1	40.2	50.0	54.1	56.2	60.2
Number of previous births					
0	81.0	71.8	68.0	64.9	64.5
≥1	19.0	28.2	32.0	35.1	35.5
<b>Treatment factors</b>					
Method of embryo fertilization and transfer*					
IVF-ET without ICSI	42.5	44.9	45.9	49.2	48.5
IVF-ET with ICSI	55.7	53.3	52.1	49.0	47.8
GIFT	0.7	0.7	0.8	1.1	2.6
ZIFT	1.1	0.9	1.1	0.6	1.0
Combination	0.1	0.2	0.2	0.2	0.2
No. of days of embryo culture†					
1	0.5	0.6	0.5	0.4	0.3
2	5.9	5.1	5.4	4.7	5.7
3	67.4	72.3	76.9	80.2	78.6
4	2.9	2.9	3.3	3.9	4.8
5	18.9	15.3	10.5	7.6	6.1
6	2.8	2.3	1.9	1.4	1.0
Number of embryos transferred					
1	4.0	5.7	6.9	9.2	13.9
2	33.3	21.8	17.2	15.7	17.9
3	40.5	35.8	27.2	22.1	19.2
4	15.7	26.5	29.8	24.3	19.3
≥5	6.5	10.1	18.8	28.7	29.7
Use of gestational carrier					
Yes	0.7	0.8	1.0	0.9	1.0
No	99.3	99.3	99.0	99.1	99.0

\* IVF-ET: in vitro fertilization with transcervical embryo transfer; ICSI: intracytoplasmic sperm injection; GIFT: gamete intrafallopian transfer; ZIFT: zygote intrafallopian transfer; and Combination: combination of IVF with or without ICSI and either GIFT or ZIFT.

† In cases of GIFT, gametes not cultured but transferred on day one.

**TABLE 4. Live-birth rates for assisted reproductive technology (ART) transfer procedures performed among patients who used freshly fertilized embryos from their own eggs, by patient age and selected patient and treatment factors — United States, 2000**

	Patient age (yrs)				
	<35 Live births per transfer procedure (%)	35–37 Live births per transfer procedure (%)	38–40 Live births per transfer procedure (%)	41–42 Live births per transfer procedure (%)	>42 Live births per transfer procedure (%)
<b>Total</b>	<b>38.4</b>	<b>33.0</b>	<b>24.3</b>	<b>14.3</b>	<b>6.0</b>
<b>Patient factors</b>					
Diagnosis					
Tubal factor	37.6	33.0	25.1	13.3	5.4
Ovulatory dysfunction	40.2	35.8	25.6	17.7	6.6
Diminished ovarian reserve	31.2	29.6	19.5	13.9	4.8
Endometriosis	39.7	33.3	26.0	17.9	3.0
Uterine factor	34.8	33.2	20.8	22.2	15.0
Male factor	38.9	34.4	25.8	14.2	8.9
Other causes	37.8	30.6	25.2	16.1	6.5
Unexplained cause	41.4	36.9	27.5	15.1	5.1
Multiple factors, female only	37.4	32.0	23.2	12.4	6.8
Multiple factors, female and male	36.8	29.5	21.6	13.3	5.5
Number of previous ART procedures					
0	40.6	35.0	25.8	14.8	6.0
≥1	35.2	31.0	23.1	14.0	6.0
Number of previous births					
0	37.3	31.7	23.1	13.4	6.0
≥1	42.9	36.3	26.9	16.0	6.0
<b>Treatment factors</b>					
Method of embryo fertilization and transfer*					
IVF-ET without ICSI	40.7	35.5	26.2	15.5	7.0
IVF-ET with ICSI	36.7	31.0	22.5	13.0	4.9
Number of days of embryo culture†					
3	37.7	32.8	24.4	14.4	5.8
5	44.4	37.7	29.6	18.2	7.5
Number of embryos transferred					
1	14.7	11.2	8.0	2.7	2.0
2	42.1	31.6	19.0	10.1	2.6
3	39.7	36.1	25.4	11.5	7.4
4	35.4	34.1	28.4	16.3	6.4
≥5	33.4	34.3	27.3	20.8	8.8
Use of gestational carrier					
Yes	44.7	41.5	29.4	15.8	13.6
No	38.4	32.9	24.3	14.3	5.9

\*IVF-ET: in vitro fertilization with transcervical embryo transfer, and ICSI: intracytoplasmic sperm injection.

†Limited to 3 and 5 days to embryo culture. ART procedures including 1, 2, 4, and 6 days to embryo culture were not included because each of these accounted for a limited proportion of procedures.



TABLE 5. Multiple-birth risk, by type of assisted reproductive technology transfer procedure performed — United States, 2000

	Patient age (yrs)	No. of live-birth deliveries	No. of multiple-birth deliveries	Multiple-birth deliveries (%)*	No. of infants born	No. of infants born in multiple-birth deliveries	Infants born in multiple-birth deliveries (%)
<b>Patient's eggs used</b>							
Freshly fertilized embryos	All ages	19,219	6,784	35.3	26,800	14,365	53.6
	<35	11,050	4,263	38.6	15,866	9,079	57.2
	35–37	4,663	1,648	35.3	6,489	3,474	53.5
	38–40	2,750	747	27.2	3,558	1,555	43.7
	41–42	625	110	17.6	739	224	30.3
	>42	131	16	12.2	148	33	22.3
Thawed embryos	All ages	2,360	609	25.8	3,048	1,297	42.6
	<35	1,378	382	27.7	1,806	810	44.9
	35–37	576	141	24.5	736	301	40.9
	38–40	283	62	21.9	357	136	38.1
	41–42	82	17	20.7	100	35	35.0
	>42	41	7	17.1	49	15	30.6
<b>Donor's eggs used†</b>							
Freshly fertilized embryos	All ages	3,041	1,236	40.6	4,382	2,577	58.8
Thawed embryos	All ages	608	177	29.1	795	364	45.8

\* Multiple-birth risk.

† Age-specific statistics are not presented for procedures that used donor eggs because only limited variation occurred by age among these procedures.



Need the latest CDC guidance on a crucial public health topic?

No problem—log on to [cdc.gov/mmwr](http://cdc.gov/mmwr) and quickly find the information you need. Browse the latest reports, research important health topics—even download ready-to-print copies—all free of charge.

Save time, get more. MMWR Online.

know what matters.



**TABLE 6. Number and percentage of infants born in multiple-birth deliveries, by patient's state/territory of residence\* at time of assisted reproductive technology (ART) treatment — United States, 2000**

Patient's state/ territory of residence	No. of infants born	No. of infants born in multiple-birth deliveries	Infants born in multiple-birth deliveries (%) <sup>†</sup>	Infants born in twin deliveries (%)	Infants born in triplet or higher order deliveries (%)
Alabama	216	128	59.3	35.2	24.1
Alaska	44	25	56.8	50.0	6.8
Arizona	444	264	59.5	43.2	16.2
Arkansas	142	70	49.3	40.8	8.5
California	4,453	2,315	52.0	45.2	6.8
Colorado	863	501	58.1	48.7	9.4
Connecticut	679	364	53.6	44.2	9.4
Delaware	117	59	50.4	42.7	7.7
District of Columbia	157	94	59.9	56.1	3.8
Florida	1,534	822	53.6	44.4	9.2
Georgia	774	385	49.7	39.7	10.1
Guam	0	0	0	0	0
Hawaii	128	69	53.9	46.9	7.0
Idaho	162	98	60.5	38.3	22.2
Illinois	2,137	1,099	51.4	40.2	11.2
Indiana	597	315	52.8	39.2	13.6
Iowa	275	141	51.3	44.7	6.5
Kansas	255	129	50.6	42.4	8.2
Kentucky	247	127	51.4	45.7	5.7
Louisiana	189	110	58.2	48.7	9.5
Maine	37	16	43.2	43.2	0
Maryland	1,153	630	54.6	46.3	8.3
Massachusetts	2,609	1,268	48.6	42.2	6.4
Michigan	1,122	603	53.7	42.6	11.1
Minnesota	908	449	49.4	39.3	10.1
Mississippi	84	41	48.8	31.0	17.9
Missouri	491	272	55.4	43.2	12.2
Montana	39	20	51.3	35.9	15.4
Nebraska	138	62	44.9	33.3	11.6
Nevada	313	195	62.3	53.7	8.6
New Hampshire	145	82	56.6	51.7	4.8
New Jersey	2,139	1,205	56.3	47.3	9.0
New Mexico	107	59	55.1	46.7	8.4
New York	3,771	1,946	51.6	44.0	7.6
North Carolina	638	390	61.1	48.6	12.5
North Dakota	51	22	43.1	43.1	0
Northern Mariana Islands	0	0	0	0	0
Ohio	843	464	55.0	42.2	12.8
Oklahoma	220	117	53.2	43.6	9.5
Oregon	321	193	60.1	49.8	10.3
Pennsylvania	997	483	48.4	40.5	7.9
Puerto Rico	112	56	50.0	33.9	16.1
Rhode Island	174	94	54.0	41.4	12.6
South Carolina	283	159	56.2	41.0	15.2
South Dakota	32	16	50.0	50.0	0
Tennessee	287	144	50.2	41.8	8.4
Texas	1,978	1,087	55.0	48.0	6.9
Utah	171	87	50.9	45.6	5.3
Vermont	40	18	45.0	45.0	0
Virgin Islands	1	0	0.0	0	0
Virginia	922	496	53.8	47.8	6.0
Washington	639	340	53.2	44.3	8.9
West Virginia	68	36	52.9	44.1	8.8
Wisconsin	305	157	51.5	45.2	6.2
Wyoming	26	18	69.2	69.2	0
Non U.S. resident	448	263	58.7	47.3	11.4
<b>Total</b>	<b>35,025</b>	<b>18,603</b>	<b>53.1</b>	<b>44.2</b>	<b>8.9</b>

\* In cases of missing residency data, the patient's state/territory of residence was assigned as the state in which the ART procedure was performed. Medical centers in all but six states had missing residency data for <15% of infants born through ART. Medical centers located in Arizona, Georgia, Massachusetts, Minnesota, New York, and Pennsylvania had >15% missing residency data.

<sup>†</sup> Numbers might not sum to total because of rounding.

FIGURE 1. Location of assisted reproductive technology (ART) medical centers — United States and Puerto Rico, 2000






FIGURE 2. Live births per transfer and percentages of multiple-infant births for assisted reproductive technology procedures performed among women aged <35 years who used freshly fertilized embryos from their own eggs, by number of embryos transferred — United States, 2000



\* Percentages of live births that were singletons, twins, and triplets or higher order are in parentheses.


**FIGURE 3. Live births per transfer and percentages of multiple-infant births for assisted reproductive technology procedures performed among women aged 35–37 years who used freshly fertilized embryos from their own eggs, by number of embryos transferred — United States, 2000**



\* Numbers might not add to 100% because of rounding.


† Percentages of live births that were singletons, twins, and triplets or higher order are in parentheses.

**FIGURE 4. Live births per transfer and percentages of multiple-infant births for assisted reproductive technology procedures performed among women aged 38–40 years who used freshly fertilized embryos from their own eggs, by number of embryos transferred — United States, 2000**




\* Percentages of live births that were singletons, twins, and triplets or higher order are in parentheses.

**FIGURE 5. Live births per transfer and percentages of multiple-infant births for assisted reproductive technology procedures performed among women aged 41–42 years who used freshly fertilized embryos from their own eggs, by number of embryos transferred — United States, 2000**



\* Numbers might not add to 100% because of rounding.  
 † Percentages of live births that were singletons, twins, and triplets or higher order are in parentheses.

**FIGURE 6. Live births per transfer and percentages of multiple-infant births for assisted reproductive technology procedures performed among women aged >42 years who used freshly fertilized embryos from their own eggs, by number of embryos transferred — United States, 2000**




\* Percentages of live births that were singletons, twins, and triplets or higher order are in parentheses.



up-to-the-minute: *adj*

1 : extending up to the immediate present, including the very latest information; see also *MMWR*.



know what matters.



The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy each week, send an e-mail message to [listserv@listserv.cdc.gov](mailto:listserv@listserv.cdc.gov). The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/publications/mmwr>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone 888-232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

All *MMWR* references are available on the Internet at <http://www.cdc.gov/mmwr>. Use the search function to find specific articles.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.