

MMWRTM
**MORBIDITY AND MORTALITY
WEEKLY REPORT**

- 1117 Hypertrophic Pyloric Stenosis in Infants Following Pertussis Prophylaxis with Erythromycin — Knoxville, Tennessee, 1999
- 1121 Carbon Monoxide Poisoning Associated with Use of LPG-Powered (Propane) Forklifts in Industrial Settings — Iowa, 1998
- 1124 Global Measles Control and Regional Elimination, 1998–1999
- 1130 Notices to Readers

**Hypertrophic Pyloric Stenosis in Infants
Following Pertussis Prophylaxis with Erythromycin —
Knoxville, Tennessee, 1999**

In February 1999, pertussis was diagnosed in six neonates born at hospital A in Knoxville, Tennessee. Because a health-care worker at hospital A was most likely the source of exposure, the local health department recommended on February 25, 1999, that erythromycin be prescribed as postexposure prophylaxis for the approximately 200 infants born at hospital A during February 1–24, 1999. In March 1999, local pediatric surgeons noticed an increased number of cases of infantile hypertrophic pyloric stenosis (IHPS) in the area, with seven cases occurring during a 2-week period. All seven IHPS cases were in infants born in hospital A during February who were given erythromycin orally for prophylaxis following possible exposure to pertussis, although none had pertussis diagnosed. The Tennessee Department of Health and CDC investigated the cluster of IHPS cases and its possible association with use of erythromycin. This report summarizes the results of the investigation, which suggest a causal role of erythromycin in this cluster of IHPS cases (1).


Case Review

IHPS cases occurring during 1997–1999 were ascertained by reviewing medical records in the two area hospitals that provide IHPS treatment. IHPS was defined as a hospital diagnosis of pyloric stenosis (*International Classification of Diseases, Ninth Revision, Clinical Modification*, code 750.5) that required pyloromyotomy in an infant born in one of the six birthing facilities in the region during 1997–1999. The rate of IHPS cases per 1000 live-born infants for each month was calculated using the number of live-born infants at the six birthing facilities as the denominator. The incidence of IHPS among infants born at hospital A peaked during February 1999 with seven IHPS cases among 217 live-born infants (rate: 32.3 cases per 1000 live-born infants) (Figure 1), a rate that was nearly seven times higher than during 1997–1998 (relative risk=6.8; 95% confidence interval [CI]=3.0–15.7). No additional IHPS cases were reported among infants born during March–May 1999 at hospital A, and the risk for IHPS in the region returned to the background rates following the peak in February 1999.

To compare the clinical characteristics of the seven index IHPS cases with those of historical IHPS cases, a detailed chart review of IHPS cases from January 1998

Stenosis — Continued

FIGURE 1. Incidence* of hypertrophic pyloric stenosis among infants born in hospital A and in all other birthing facilities — Knoxville, Tennessee, 1997–May 1999



*Per 1000 live-born infants.

through March 1999 was conducted at the two hospitals in the region that had pediatric surgery services. The diagnostic features of the seven index cases were similar to 40 historical cases. Compared with historical cases, index case-patients were younger at the time of admission for IHPS (mean age=25.6 days versus 35.4 days) and were less likely to have a family history of IHPS (0% versus 17.5%). The mean pyloric thickness and length as measured on ultrasound were similar in the two groups. All index case-patients had received oral erythromycin, compared with none of the historical case-patients.

To validate the IHPS diagnoses, a pediatric radiologist, who was blinded to the original readings, reviewed ultrasound films for the seven index case-patients and seven infants without IHPS. The ultrasound review showed perfect agreement with the original readings (Kappa=1.0; 95% CI=0.48–1.0).

Cohort Study

A retrospective cohort study of 282 infants born during January–February 1999 at hospital A was conducted to assess a possible association between erythromycin use, gastrointestinal symptoms, and IHPS. In the cohort, 157 infants (55.7%) had a history of oral erythromycin use. The prevalence of erythromycin use was 8.6% among 116 infants born during January 1999 and 88.6% among 166 infants born during

Stenosis — Continued

February 1999. The erythromycin preparations administered to the infants included ethyl succinate (n=83), estolate (n=59), both ethyl succinate and estolate (n=one), and unknown (n=14). No differences were observed in gastrointestinal symptoms or risk for IHPS in relation to the type of erythromycin preparation.

The infants who were given erythromycin but who did not develop IHPS were aged 1–53 days when they began erythromycin (median age=13 days; mean=14.1 days), and the duration of erythromycin exposure ranged from 1 to 21 days (median duration=14 days; mean=12.2 days). The seven index IHPS case-patients were aged 2–17 days when they began erythromycin (median=5 days; mean=9.3 days), and the duration of their erythromycin exposure ranged from 10–18 days (median duration=14 days; mean=13.3 days). Seven IHPS cases occurred among infants who were exposed to erythromycin and none among infants not exposed to erythromycin (relative risk=infinity, lower bound of exact 95% CI=1.7).

Reported by: L Patterson, MD, J Peeden, MD, S Sirlin, MD, East Tennessee Children's Hospital; S Hall, MD, IM Himelright, MD, Knox County Health Dept, Knoxville; AS Craig, MD, WL Moore, MD, State Epidemiologist, Tennessee Dept of Health. B Lee, MD, Johns Hopkins School of Medicine, Baltimore, Maryland. Child Vaccine Preventable Diseases Br, Epidemiology and Surveillance Div, National Immunization Program; Birth Defects and Pediatric Genetics Br, Div of Birth Defects, Child Development, and Disability and Health (proposed), National Center for Environmental Health, CDC.

Editorial Note: IHPS is a hypertrophy of the pyloric muscle that usually results in non-bilious, projectile vomiting that begins at about 3.5 weeks of age (2). IHPS affects approximately one to three infants per 1000 live-born infants and affects about four to five times as many male as female infants (3,4). Evidence suggests that the pyloric muscle hypertrophy of IHPS develops postnatally (5). The first case reports of a possible association between IHPS and erythromycin in five neonates were published in 1976 (6), but the association was considered improbable and had remained unconfirmed. The only subsequent report of this association was a single case report of IHPS in a breastfed infant whose mother had taken erythromycin (7). The findings in this report provide further evidence that erythromycin has a causal role in the etiology of IHPS and raise concerns about the use of erythromycin in neonates.

The peak in IHPS incidence in this region corresponded temporally with the use of erythromycin following the county health department recommendation. All index IHPS case-patients began having symptoms of either vomiting or excessive irritability while taking erythromycin.

The study described in this report is not population-based but includes all live-born infants at facilities in the Knoxville metropolitan area. Local clinicians and public health workers considered it unlikely that an infant born at one of these facilities would be referred outside the region for pediatric surgery, but this possibility cannot be completely eliminated. No evidence indicated a change in case definition, in referral patterns, or in pediatric surgeons or pediatric radiologists that could account for this increase in IHPS incidence. It is unlikely that children with severely hypertrophied pylori would not exhibit symptoms, and evaluation of the pyloric muscle of normal children versus those with IHPS has not demonstrated the existence of severe hypertrophy among asymptomatic children (8). Therefore, it is unlikely that IHPS cases were missed.

Previous epidemiologic studies of IHPS have not identified erythromycin as a risk factor, possibly because few neonates included in such studies were exposed to

Stenosis — Continued

erythromycin. In most mass prophylaxis situations, the number of neonates treated may be small, possibly explaining why an increased risk for IHPS with erythromycin had not been established.

The prevention of pertussis in infants is important; most hospitalizations for and deaths from pertussis occur in children aged <1 year (9). Although no data exist to confirm a safe and effective alternative to erythromycin for prophylaxis of neonates exposed to pertussis, these findings indicate a need for further examination of recommendations for erythromycin prophylaxis (10). The high case-fatality ratio of pertussis in neonates demonstrates the need to prevent pertussis in this age group, as was done successfully in Tennessee. However, public health officials should continue to use caution in defining risk groups to minimize unnecessary prophylaxis. Physicians who prescribe erythromycin to newborns should inform parents about the possible risks for IHPS and counsel them about signs of developing IHPS.

Cases of pyloric stenosis following use of oral erythromycin should be reported to the Food and Drug Administration (FDA) MedWatch, telephone (800) 332-1088, or through the World-Wide Web, <http://www.fda.gov/medwatch>.^{*} Additional information on use of erythromycin for treatment of ophthalmia neonatorum and infant pneumonia caused by *Chlamydia trachomatis* in newborns is available at <http://www.cdc.gov/nchstp/dstd/eryth.htm> or by fax, (800) 332-0178.

References

1. Honein MA, Paulozzi LJ, Himelright IM, et al. Infantile hypertrophic pyloric stenosis after pertussis prophylaxis with erythromycin: a case review and cohort study. *Lancet* 1999;354:2101-5.
2. Spicer RD. Infantile hypertrophic pyloric stenosis: a review. *Br J Surg* 1982;69:128-35.
3. Rasmussen L, Green A, Hansen LP. The epidemiology of infantile hypertrophic pyloric stenosis in a Danish population, 1950-84. *Int J Epidemiol* 1989;18:413-7.
4. Schechter R, Torfs CP, Bateson TF. The epidemiology of infantile hypertrophic pyloric stenosis. *Paediatr Perinat Epidemiol* 1997;11:407-27.
5. Rollins MD, Shields MD, Quinn RJM, Wooldridge MAW. Pyloric stenosis: congenital or acquired? *Arch Dis Child* 1989;64:138-47.
6. San Filippo JA. Infantile hypertrophic pyloric stenosis related to ingestion of erythromycin estolate: a report of five cases. *J Pediatr Surg* 1976;11:177-80.
7. Stang H. Pyloric stenosis associated with erythromycin ingested through breastmilk. *Minn Med* 1986;69:669-70,682.
8. Rohrschneider WK, Mittnacht H, Darge K, Troger J. Pyloric muscle in asymptomatic infants: sonographic evaluation and discrimination from idiopathic hypertrophic pyloric stenosis. *Pediatr Radiol* 1998;28:429-34.
9. Sutter RW, Cochi SL. Pertussis hospitalizations and mortality in the United States, 1985-1988. *JAMA* 1992;267:386-91.
10. American Academy of Pediatrics. Pertussis. In: Peter G, ed. 1997 Red book: report of the Committee on Infectious Diseases. 24th ed. Elk Grove Village, Illinois: American Academy of Pediatrics 1997:397.

^{*}References to sites of non-CDC organizations on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC. CDC is not responsible for the content of pages found at these sites.

Carbon Monoxide Poisoning Associated with Use of LPG-Powered (Propane) Forklifts in Industrial Settings — Iowa, 1998

In 1998, the Iowa Department of Public Health (IDPH) and Iowa State University (ISU) Extension Department, with the assistance of local health departments, investigated a series of carbon monoxide (CO) poisonings associated with the use of liquified petroleum gas (LPG)-powered forklifts in light industry. In each episode, forklifts emitting high CO concentration levels were operated in inadequately ventilated warehouse and production facilities, which resulted in high CO accumulations. Employees at each site developed symptoms of CO poisoning, and some employees received inadequate or inappropriate medical care. This report summarizes the investigations and provides recommendations to prevent such incidents.

Incident 1

On August 17 and 18, 1998, during three consecutive 8-hour shifts, 34 (45%) of 75 plastic manufacturing plant employees experienced symptoms of CO poisoning (primarily headaches) while at work. Ten ill employees were evaluated at three local emergency departments (EDs). Of five employees seen at one ED, possible CO poisoning initially was diagnosed in three workers. However, because of high pulse oximeter readings, this diagnosis was dismissed erroneously, and the three employees were discharged and returned to work. The other two employees had "possible poly vinyl chloride inhalation" and "syncopal episode" diagnosed, respectively; one was admitted to the hospital, and one was discharged home. Of four employees seen at a second ED, the first two had "migraine headache" and "torticollis" diagnosed, and the second two were suspected to be CO poisoned and had carboxyhemoglobin (COHb) levels of 3.8% (1 hour after leaving work) and 10.7% (2 hours after leaving work), respectively.* One employee was seen at a third ED, and a headache of undetermined cause was diagnosed.

A local physician notified IDPH when several plant employees sought follow-up treatment the next day. Overall, 25 (38%) of 65 plant employees interviewed by IDPH had illnesses that met the case definition of CO poisoning (i.e., headache and at least one of the following: weakness, dizziness, or nausea). Illness rates increased with each shift, and no substantial associations were found between illness and age, sex, recent illness such as cold or influenza, illness in family members, hay fever, asthma, or smoking.

When measured by investigators, the plant's two forklifts each emitted concentrations of CO in excess of 40,000 ppm (recommended guidelines range from 2000 to 10,000 ppm [1-3]). On August 17, the plant's air-conditioning system had been shut down for servicing, and an exhaust fan had malfunctioned, reducing the effective ventilation rate. However, the forklifts emitted such excessive amounts of CO that no practical level of ventilation could have maintained CO concentrations below recommended exposure limits.† Neither employees nor managers were aware that the

*Normal COHb concentrations are <2% in nonsmokers and 5%–9% in smokers.

†CDC's National Institute for Occupational Safety and Health recommends that CO exposure not exceed 35 ppm as an 8-hour time-weighted average and that point exposure should never exceed 200 ppm.

Carbon Monoxide Poisoning — Continued

symptoms they experienced were related to CO poisoning, which delayed recognition and response.

Incident 2

In November 1998, after experiencing headaches, nausea, and dizziness over several days, employees of a warehouse brought conventional residential CO detectors to work; these detectors registered CO concentrations of 30–136 ppm. In the adjacent office area, concentrations as high as 76 ppm were recorded before employees inactivated the detectors to silence the continuous alarms. Employing industrial CO detectors, the investigation by IDPH determined that the facility's LPG-powered forklifts (producing from 40,000 to 70,000 ppm of CO) and inadequate plant ventilation allowed accumulations of CO up to 267 ppm in the warehouse. No employees reported seeking medical treatment.

Incident 3

From December 1998 through January 5, 1999, employees of an embroidery company experienced headaches and fatigue, and an employee's puppy became somnolent when brought to work. A local energy company was called to investigate. The company measured CO concentrations of 100–200 ppm in the embroidery offices. While attempting to find the source of CO, investigators found levels of 200–450 ppm in a wooden pallet manufacturer located in the same building one floor below the embroidery offices.

One symptomatic office employee, a pregnant woman, consulted her obstetrician and reportedly was told that no postexposure treatment existed. Approximately 24 hours after her last exposure to CO and after seeking medical advice from experts in CO poisoning, she and another symptomatic employee were treated with hyperbaric oxygen (4). At the time of treatment, their COHb levels were within the normal range but both were still having symptoms. Both employees demonstrated substantial subjective improvement after treatment. The since-delivered child is being monitored for CO-related complications such as neurologic conditions and growth abnormalities.

In the subsequent investigation, 23 workers were interviewed; two (29%) of seven embroidery employees and four (25%) of 16 pallet company employees had illnesses that met the case definition for CO poisoning. Investigators found an association between illness and proximity of the person's work station to areas where the forklifts were operated. The pallet manufacturer's forklifts emitted up to 75,000 ppm of CO into the inadequately ventilated warehouse. The embroidery office's furnace was vented properly with satisfactory combustion. However, the furnace was in the warehouse of the pallet company and pulled high CO-content ambient air from the warehouse into the heating system and distributed it to the embroidery office.

Reported by: RD Comstock, MS, RW Currier, DVM, KV Markiewicz, PhD, RL Welke, MP Quinlisk, MD, State Epidemiologist, Iowa Dept of Public Health; TH Greiner, PhD, Iowa State Univ Extension Dept, Ames, Iowa. Denver Field Office, Div of Surveillance, Hazard Evaluation, and Field Studies, National Institute for Occupational Safety and Health, CDC.

Editorial Note: CO poisoning associated with indoor combustion sources has long been recognized but continues to be a problem in the United States. The events described in this report illustrate factors that result in failure to adequately prevent CO poisoning and to promptly recognize such incidents when they occur. Timely and

Carbon Monoxide Poisoning — Continued

correct clinical diagnosis of acute CO poisoning remains elusive because of the non-specific and protean nature of its signs and symptoms (i.e., headache, nausea, lethargy, weakness, abdominal discomfort/pain, confusion, dizziness, visual disturbances [including blurred vision], numbness and tingling, ataxia, irritability, agitation, chest pain, dyspnea on exertion, palpitations, seizures, and loss of consciousness). In incident 1, failure to diagnose illness correctly in the first employees evaluated resulted in some CO-intoxicated employees being sent back to work and further exposure and in continued exposures to other workers at the plant. Correct diagnosis can be achieved by determining COHb levels in the patient. However, screening can be performed by breath analyzer instruments. Pulse oximeter testing does not reflect tissue hypoxia and cannot be used to screen or diagnose (5). Correct identification of the CO source requires specific resources (i.e., proper monitoring equipment; time for thorough investigation; and knowledge about potential CO sources, such as LPG-powered forklifts); these resources often may be unavailable on site, particularly in small business or light industrial settings but are frequently available through local utility companies.

Treatment for acute CO poisoning varies. The Undersea and Hyperbaric Medical Society provides guidelines to physicians for treating CO poisoning (6). These guidelines recommend that patients who manifest signs and symptoms of intoxication (e.g., altered mental status or neurologic signs, cardiovascular dysfunction, pulmonary edema, or severe acidosis) be referred for hyperbaric therapy regardless of their COHb levels (4).

In June 1998, the Council of State and Territorial Epidemiologists (CSTE) adopted a surveillance case definition for acute CO poisoning (7) that delineates criteria for categorizing reported acute CO poisonings. However, no commonly accepted clinical case definition nor consistent constellation of signs or symptoms exists that would unequivocally identify a case. All cases described in this report met the CSTE surveillance criteria for classification as confirmed cases.

Circumstances surrounding the continuing occurrence of CO poisonings and related confusion about identification of disease symptoms and appropriate treatment of cases illustrate the need for 1) improved education for ED and primary-care physicians about symptoms of CO poisoning, appropriate testing, and treatment (4,6); 2) improved education for employers, employees, and forklift maintenance providers about the hazards of using improperly or poorly maintained LPG-powered forklifts indoors, CO poisoning symptoms, and the appropriate response to CO symptoms; and 3) improved forklift maintenance, ventilation, and CO-monitoring procedures when LPG-powered forklifts are used in enclosed settings.

References

1. McCammon JB, McKenzie LE, Heinzman M. Carbon monoxide poisoning related to the indoor use of LPG-fueled forklifts in Colorado workplaces. *Appl Occ Envir Hyg* 1996;11:192-8.
2. Michigan Department of Public Health. Industrial lift trucks: maintaining acceptable air quality in the workplace. *Michigan's Occupational Health* 1996;29:1-8.
3. American Council of Industrial Hygienists. *Industrial ventilation: a manual of recommended practice*. 22nd ed. Cincinnati, Ohio: American Council of Industrial Hygienists, 1995.
4. Ilano AL, Raffin TA. Management of carbon monoxide poisoning. *Chest* 1990;97:165-9.
5. Reisdorf EJ, Shah SM, Nelson R. Carbon monoxide poisoning: from crib death to pickup trucks. *Emergency Medical Report* 1993;14:181-90.
6. Hampson NB, ed. *Hyperbaric oxygen therapy: 1999 committee report*. Kensington, Maryland: Undersea and Hyperbaric Medical Society, 1999:9-12.

Carbon Monoxide Poisoning — Continued

7. Council of State and Territorial Epidemiologists. CSTE: position statement EH-1. Surveillance case definition for acute carbon monoxide poisoning. Atlanta, Georgia: Council of State and Territorial Epidemiologists, June 1998.

Global Measles Control and Regional Elimination, 1998–1999

In 1989, the World Health Assembly adopted the goal of reducing measles morbidity and mortality by 90% and 95%, respectively, by 1995, compared with estimates of the disease burden in the prevaccine era (1). In 1990, the World Summit for Children adopted a goal of vaccinating 90% of children against measles by 2000. Three regions of the World Health Organization (WHO) have targeted elimination: in 1994, the American Region (AMR) targeted elimination by 2000; in 1997, the Eastern Mediterranean Region (EMR) targeted elimination by 2010; and in 1998, the European Region (EUR) targeted elimination by 2007. This report updates progress since 1997 (2) toward global measles control and regional elimination of measles, and includes vaccination coverage and disease surveillance data received by WHO as of August 14, 1999. Data for 1998 suggest that routine measles vaccination coverage has declined in some regions, the number of countries reporting cases and coverage to WHO has decreased, and measles continues to be an important cause of morbidity and mortality.

Reported Routine Measles Vaccination Coverage

Global reported coverage with one dose of measles vaccine declined from 79% in 1997 to 72% in 1998 (Table 1). In 1998, 14 countries reported measles coverage below 50%: 10 in the African Region (AFR) (Burundi, Cameroon, Central African Republic, Chad, Democratic Republic of Congo, Ethiopia, Liberia, Nigeria, Togo, and Uganda), one in AMR (Haiti), two in EMR (Afghanistan and Somalia), and one in the South-East Asia Region (SEAR) (Democratic People's Republic of Korea).

Among regions focusing on measles control, AFR and SEAR reported the lowest routine vaccination coverage rates, 49% and 67%, respectively (Table 1). These regions reported the greatest decrease in coverage during 1997–1998. The Western Pacific Region (WPR) continued to report the highest routine vaccination coverage (93%).

Among regions with an elimination target, AMR reported the highest coverage rate (86%) (Table 1). In EMR, regional measles vaccination coverage was 78%, and 14 polio-free countries that began implementing measles elimination strategies reported routine coverage rates >85% (3). EUR reported a routine first dose coverage rate of 71% in 1998; 21 (41%) of 51 EUR countries* did not report vaccination coverage data to WHO.

Supplementary Vaccination Campaigns

Supplemental vaccination campaigns have been conducted in several countries targeting either measles morbidity and mortality reduction or elimination. In 1998 and 1999, 31 countries in AFR[†] and three countries in EMR (Djibouti, Egypt, and Sudan)

*Andorra, Austria, Belgium, Denmark, Finland, France, Germany, Greece, Iceland, Ireland, Luxembourg, Monaco, Norway, Poland, San Marino, Spain, Sweden, Switzerland, the former Yugoslav Republic of Macedonia, Turkey, and Yugoslavia.

[†]Angola, Benin, Burkina Faso, Burundi, Cameroon, Central African Republic, Chad, Congo, Côte d'Ivoire, Democratic Republic of Congo, Equatorial Guinea, Eritrea, Ethiopia, Gabon, Ghana, Guinea, Kenya, Liberia, Madagascar, Mali, Mauritania, Mozambique, Niger, Nigeria, Rwanda, Senegal, Sierra Leone, Togo, Uganda, United Republic of Tanzania, and Zambia.

TABLE 1. Reported routine measles vaccination* coverage among children aged 1 year, by World Health Organization (WHO) region — worldwide, 1997 and 1998†

| Region | Reported coverage | | | Completeness of reporting from countries | | | | |
|---------------------------------|-------------------|--------------|----------------------------|------------------------------------------|-----------------------------------|------------|------------------------------------------|------------|
| | 1997 | 1998 | % change from 1997 to 1998 | Total | Countries and areas | | % completeness of reporting [§] | |
| | | | | | No. countries and areas reporting | | 1997 | 1998 |
| Measles elimination goal | | | | | | | | |
| American¶ | 88% | 86% | -2 | 46 | 40 | 38 | 98% | 99% |
| Eastern Mediterranean | 80%** | 78%** | -2 | 24 | 23 | 20 | 97% | 94% |
| European | 76%** | 71%** | -5 | 51 | 35 | 30 | 64% | 57% |
| Measles control goal | | | | | | | | |
| African | 56%** | 49%** | -7 | 48 | 41 | 36 | 92% | 89% |
| South East Asian | 84%** | 67%** | -17 | 10 | 8 | 9 | 96% | 97% |
| Western Pacific | 93%** | 93%** | 0 | 36 | 35 | 31 | 100% | 95% |
| Total | 79%** | 72%** | -7 | 215 | 182 | 164 | 94% | 91% |

* One dose of measles-containing vaccine (MCV).

† Reported to WHO as of August 14, 1999.

§ Numerator=total number of surviving infants in countries reporting MCV coverage to WHO; denominator=1998 estimates of surviving infants in region (Source: United Nations. World population prospects: 1998 revision, Population Division, Department of Economic and Social Affairs, New York: United Nations, 1999).

¶ Data provided by the Pan American Health Organization, excluding the United States. In the United States, one dose MCV coverage among children aged 19–35 months was 91% in 1997 and 92% in 1998.

** Model-based imputation used to account for missing data.

Global Measles Control — Continued

conducted mass vaccination campaigns in high-risk areas to reduce morbidity and mortality among those children who were not vaccinated through routine vaccination services. During 1998–1999, two countries (Marshall Islands and Palau) in WPR conducted vaccination campaigns targeting children who had not been vaccinated through routine vaccination services, two countries (Lao People's Democratic Republic and Viet Nam) delivered measles vaccination to remote populations during polio subnational immunization days, and one country (Viet Nam) conducted a pilot campaign in one province.

WHO's measles elimination strategy comprises a three-part vaccination strategy (i.e., "catch-up," "keep-up," and "follow-up"[§]); two parts are supplemental vaccination (4). All countries in AMR, except the United States and the French and Dutch Antilles, completed catch-up campaigns by 1996. Since then, most countries in AMR have been conducting follow-up campaigns.

In nine of 15 EMR countries where measles elimination activities are ongoing, 13 million children have been vaccinated during catch-up measles vaccination campaigns conducted since 1994 (3). In EUR, Romania implemented a catch-up campaign during 1998–1999 targeting all children aged 7–18 years (girls aged 15–18 years received measles and rubella vaccine). Approximately 2 million children were vaccinated and 93% coverage was reported (WHO, unpublished data, 1999). During 1998–1999, staff from 23 (45%) of 51 countries[¶] in EUR attended workshops at which they evaluated their age-specific susceptibility to measles and determined strategies to reduce susceptibility to <15% for ages 0–4 years, <10% for ages 5–9 years, and <5% for ages ≥10 years (5).

Since 1995, 23 million children have been vaccinated during catch-up campaigns in the six southern African nations where measles-elimination initiatives have been launched (6). In addition, United Kingdom (1994), Bhutan (1995), the Maldives (1995), Mongolia (1996), Papua New Guinea (1997), New Zealand (1997), Australia (1998), parts of China (1997–1998), the Philippines (1998), and 13 Pacific island countries and areas (since 1997) conducted catch-up campaigns.

Reported and Estimated Measles Morbidity and Mortality

Among regions with measles elimination goals, the AMR reported the lowest incidence (1.6 per 100,000) in 1998 (Table 2). The measles outbreak that began in Brazil in 1997 affecting unvaccinated adults continued in 1998 and 1999 among unvaccinated young children in Argentina, Bolivia, Colombia, the Dominican Republic, and Paraguay. As of November 27, 1999, 2698 measles cases have been confirmed in the region compared with 10,067 cases for the same period in 1998. During 1997–1998 in EMR, the number of cases reported increased by 58%; outbreaks were reported in Iran, Syria, Morocco, and Saudi Arabia. In EUR, the number of cases reported declined 59%, but the number of countries reporting measles cases declined from 45 in 1997 to 31 in 1998. Among all regions, AFR reported the highest number of measles cases and

[§] "Catch-up" is a one-time, nationwide vaccination campaign targeting usually all children aged 9 months–14 years, regardless of history of measles disease or vaccination status; "keep-up" is routine services aimed at vaccinating 95% of each successive birth cohort; and "follow-up" is subsequent nationwide vaccination campaigns conducted every 2–5 years targeting usually all children born after the catch-up campaign.

[¶] Andorra, Bulgaria, Croatia, Czech Republic, Denmark, Germany, Greece, Hungary, Italy, Kazakhstan, Kyrgyzstan, Malta, Netherlands, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Switzerland, Tajikistan, Turkmenistan, and Uzbekistan.

TABLE 2. Reported measles cases and a comparison of measles surveillance, by World Health Organization (WHO) region — worldwide, 1997 and 1998*

| Region | Reported cases | | % change from 1997 to 1998 | Incidence [†] | | Completeness of reporting from countries | | | | |
|---------------------------------|----------------|----------------|----------------------------|------------------------|-------------|------------------------------------------|--------------------------------------|-------------------------------------------------------------------------|------------|------------|
| | 1997 | 1998 | | 1997 | 1998 | Countries and areas | | % region's population [§] living in countries reporting to WHO | | |
| | | | | | | Total | No. of countries and areas reporting | 1997 | 1998 | |
| Measles elimination goal | | | | | | | | | | |
| American | 51,926 | 12,941 | -75% | 6.5 | 1.6 | 47 | 44 | 43 | 100% | 100% |
| Eastern Mediterranean | 33,342 | 52,666 | 58% | 8.0 | 11.1 | 24 | 20 | 23 | 90% | 100% |
| European | 103,129 | 42,768 | -59% | 14.4 | 8.2 | 51 | 45 | 31 | 82% | 60% |
| Measles control goal | | | | | | | | | | |
| African | 299,623 | 349,814 | 17% | 49.2 | 61.7 | 48 | 45 | 34 | 100% | 91% |
| South East Asian | 114,331 | 62,722 | -45% | 7.8 | 4.2 | 10 | 9 | 10 | 100% | 100% |
| Western Pacific | 142,115 | 76,037 | -46% | 8.7 | 5.0 | 36 | 36 | 32 | 100% | 92% |
| Total | 744,466 | 596,948 | -16% | 13.2 | 11.1 | 216 | 199 | 173 | 97% | 91% |

* Reported to WHO as of August 14, 1999.

[†] Reported cases per 100,000 total population of the countries reporting in the region.

[§] 1998 total population estimates by country (Source: United Nations. World population prospects: 1998 revision, Population Division, Department of Economic and Social Affairs, New York: United Nations, 1999).

Global Measles Control — Continued

incidence. Of all the cases reported, more than half were reported from countries in AFR.

Each year, WHO estimates actual measles morbidity and mortality; because measles is not a notifiable disease in some countries, substantial underreporting of measles occurs, and measles deaths are not reported to WHO. For 1998, WHO estimated that approximately 30 million measles cases and 888,000 measles-related deaths occurred worldwide; an estimated 85% of the measles-related deaths occurred in AFR and SEAR (7).

Global Measles Laboratory Network

Efforts are under way to establish a Global Measles Laboratory Network. Measles laboratories of CDC and the Central Public Laboratory Services in the United Kingdom have been selected as the Global Measles Strain Banks. Activities to strengthen laboratory capacity to support measles surveillance include assessment of country laboratory needs, training of laboratory staff, provision of diagnostic kits, and collection of specimens for diagnosis and virus isolation. During 1998–1999, eight measles laboratory workshops were conducted, and 105 laboratory staff from 42 countries in five regions were trained in basic measles diagnostic methods.

Reported by: Vaccines and Biologicals Dept, World Health Organization, Geneva, Switzerland. Respiratory and Enteric Viruses Br, Div of Viral and Rickettsial Diseases, National Center for Infectious Diseases; Epidemiology and Surveillance Div; Vaccine Preventable Disease Eradication Div, National Immunization Program; and an EIS Officer, CDC.

Editorial Note: With approximately 1 million deaths attributed to measles in 1998, measles remains an important cause of vaccine-preventable illness and death. Failure to deliver at least one dose of measles vaccine to all infants remains the primary reason, despite widespread availability of an effective and safe vaccine. Morbidity and mortality decrease with increasing vaccination coverage levels; those regions with the lowest coverage levels have the highest burden, with AFR continuing to report both the lowest coverage and highest incidence.

Global and regional (except AMR) routine vaccination coverage rates in 1997 and 1998 were calculated using model-based estimates to account for missing data (8). Nationwide surveys indicated that in some countries actual coverage may be lower than reported coverage (9). For this reason, some countries in SEAR (Bangladesh, India, and Indonesia) have begun reporting coverage based on surveys rather than the administrative method. In part, this change in reporting accounts for the decline in reported coverage in SEAR in 1998. Although some regions (e.g., WPR) may have achieved the World Summit for Children goal, coverage in some WPR countries and in the remaining five regions is <90%. Reported regional routine vaccination coverage rates in the three regions with measles elimination goals are <90%, thus increasing the speed at which susceptible children accumulate and the need for more frequent follow-up campaigns to prevent re-emergence of measles (10). Further improvements in routine vaccination coverage and methods used to monitor it are needed to decrease the morbidity and mortality associated with measles.

During 1997–1998, the number of countries reporting vaccination coverage or measles cases decreased in some regions. EUR had the highest proportion of regional population from which data were not reported. Strengthening of measles surveillance is required in both developed and developing countries to monitor progress toward achieving morbidity and mortality reduction or regional elimination

Global Measles Control — Continued

goals. All countries should improve routine reporting of measles cases by month of occurrence and geopolitical unit. Countries should use outbreak investigations to obtain data on age and vaccination status of persons with measles and to estimate population-based case-fatality ratios. Case-based surveillance with laboratory confirmation of suspected measles cases and virus isolation from all outbreaks are needed when incidence of measles decreases to low levels following implementation of measles elimination measures. The global measles laboratory network needs to be strengthened by WHO, especially in those countries with elimination goals, by recruiting additional laboratories and compiling standard procedures for testing of samples.

Reduced measles incidence under conditions of improved surveillance suggests substantial progress in AMR toward achieving the regional measles elimination goal. Recent resurgence of measles in this region emphasizes the importance of full and timely implementation of elimination strategies. In EMR, routine vaccination coverage and surveillance need to be further strengthened throughout the region. Appropriate vaccination strategies for elimination need to be implemented to reduce susceptibility to measles in countries of EUR. Lack of reporting from some of the western European countries impairs assessment of disease burden and coverage in the region and suggests an urgent need to improve measles surveillance and to monitor vaccination coverage.

The priorities for countries pursuing accelerated measles control include improving routine vaccination coverage levels to at least 80% in all districts of every country, achieving at least 90% coverage nationwide, conducting supplementary vaccination campaigns together with administration of vitamin A in high-risk areas, and improving completeness and timeliness of reporting of measles cases at district level. Priorities for countries and regions with a measles elimination goal include improving routine vaccination coverage levels to at least 90% in all districts of every country (resulting in nationwide coverage $\geq 95\%$); achieving coverage $>90\%$ in catch-up and follow-up campaigns or achieving nationwide coverage $\geq 95\%$ with a routine second dose of measles vaccine, and establishing case-based surveillance with laboratory confirmation of suspected cases and virus isolation from all chains of transmission. Adherence to these priorities will ensure that the measles morbidity and mortality burden will decrease and that the measles disease reduction targets can be reached.

References

1. World Health Assembly. Executive summary. Geneva, Switzerland: World Health Organization, 1989; resolution no. WHA 42.32.
2. CDC. Progress toward global measles control and regional elimination, 1990–1997. *MMWR* 1998;47:1049–54.
3. CDC. Progress toward measles elimination—Eastern Mediterranean region, 1980–1998. *MMWR* 1999;48:1081–6.
4. De Quadros CA, Olive JM, Hersh BS, et al. Measles elimination in Americas: evolving strategies. *JAMA* 1996;275:224–9.
5. World Health Organization. Strategic plan for the elimination of measles in the European region. Expanded program on immunization, seventh meeting of national program managers. Berlin, Germany, November 10–12, 1997.
6. CDC. Progress toward measles elimination—southern Africa, 1996–1998. *MMWR* 1999;48:585–9.
7. World Health Organization. The world health report 1999—making a difference. Geneva, Switzerland: World Health Organization, 1999:98,104.

Global Measles — Continued

8. World Health Organization. WHO Vaccine Preventable Diseases Monitoring System 1999 global summary. Geneva, Switzerland: World Health Organization, Department of Vaccines and Other Biologicals, 1999 (WHO/V&B/99.17).
9. CDC. Measles control—South-East Asia Region, 1990–1997. *MMWR* 1999;48:541–5.
10. Nokes DJ, Swinton J. Vaccination in pulses: a strategy for global eradication of measles and polio? *J Math Appl Med Biol* 1995;12:29–53.

Notice to Readers**Publication of the Updated Inventory
of Managed-Care-Related Projects, 1998**

CDC supports extramural projects in various managed-care settings and periodically inventories them to inform public and private prevention communities of relevant findings, products and ongoing efforts; and to provide benchmarks for new project development. In 1996, CDC published its first *Inventory of Managed Care-Related Projects: Fiscal Year 1995–1996*, which catalogued 83 activities. This latest release, the *Inventory of Managed Care-Related Projects: 1998 (1)*, describes 107 projects covering a wide range of activities—from studies of behavior interventions to analyses of vaccine effectiveness to comparisons of health-care delivery systems, and including examples of successful collaborations between the public health and managed-care communities.

The *Inventory* can be viewed on CDC's World-Wide Web site at <http://www.cdc.gov/epo/dpram/managedcare/intro.htm>. Paper copies can be obtained from the Office of HealthCare Partnerships, CDC, 4770 Buford Highway, Mailstop K73, Atlanta, GA 30341; or telephone (770) 488-8186.

Reference

1. CDC. Inventory of managed care-related projects: 1998. Atlanta, Georgia: US Department of Health and Human Services, CDC, 1999.


Notice to Readers**Epidemiology in Action: Intermediate Methods**

CDC and Emory University's Rollins School of Public Health will co-sponsor a course, "Epidemiology in Action: Intermediate Methods" on February 7–11, 2000, in Atlanta. The course is designed for state and local public health professionals.

The course will review the fundamentals of descriptive epidemiology and biostatistics, analytic epidemiology, and Epi Info 6 but will focus on mid-level epidemiologic methods directed at strengthening participants' quantitative skills, with an emphasis on up-to-date data analysis. Topics include advanced measures of association, normal and binomial distributions, logistic regression, field investigations, and summary of statistical methods. Prerequisite is an introductory course in epidemiology (e.g., such as *Epidemiology in Action* or *International Course in Applied Epidemiology*) or any other introductory class. There is a tuition charge.

(Continued on page 1139)

FIGURE I. Selected notifiable disease reports, comparison of provisional 4-week totals ending December 11, 1999, with historical data — United States



*Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

TABLE I. Summary — provisional cases of selected notifiable diseases, United States, cumulative, week ending December 11, 1999 (49th Week)

| | Cum. 1999 | | Cum. 1999 |
|---------------------------------------------|-----------|-----------------------------------------|-----------|
| Anthrax | - | HIV infection, pediatric* ⁵ | 137 |
| Brucellosis* | 48 | Plague | 8 |
| Cholera | 3 | Poliomyelitis, paralytic | - |
| Congenital rubella syndrome | 6 | Psittacosis* | 16 |
| Cyclosporiasis* | 50 | Rabies, human | - |
| Diphtheria | 1 | Rocky Mountain spotted fever (RMSF) | 534 |
| Encephalitis: California* | 60 | Streptococcal disease, invasive Group A | 2,014 |
| eastern equine* | 6 | Streptococcal toxic-shock syndrome* | 36 |
| St. Louis* | 5 | Syphilis, congenital [¶] | 271 |
| western equine* | 1 | Tetanus | 31 |
| Ehrlichiosis | 149 | Toxic-shock syndrome | 113 |
| human granulocytic (HGE)* | 40 | Trichinosis | 9 |
| human monocytic (HME)* | 93 | Typhoid fever | 294 |
| Hansen Disease* | 20 | Yellow fever | 1 |
| Hantavirus pulmonary syndrome* [†] | 117 | | |
| Hemolytic uremic syndrome, post-diarrheal* | | | |

-:no reported cases

*Not notifiable in all states.

[†] Updated weekly from reports to the Division of Viral and Rickettsial Diseases, National Center for Infectious Diseases (NCID).

⁵ Updated monthly from reports to the Division of HIV/AIDS Prevention—Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention (NCHSTP), last update November 28, 1999.

[¶] Updated from reports to the Division of STD Prevention, NCHSTP.

TABLE II. Provisional cases of selected notifiable diseases, United States, weeks ending December 11, 1999, and December 12, 1998 (49th Week)

| Reporting Area | AIDS | | Chlamydia | | Cryptosporidiosis | | <i>Escherichia coli</i> O157:H7* | | | |
|----------------|------------|-----------|-----------|-----------|-------------------|-----------|----------------------------------|-----------|-----------|-----------|
| | Cum. 1999† | Cum. 1998 | Cum. 1999 | Cum. 1998 | Cum. 1999 | Cum. 1998 | NETSS | | PHLIS | |
| | | | | | | | Cum. 1999 | Cum. 1998 | Cum. 1999 | Cum. 1998 |
| UNITED STATES | 40,933 | 42,308 | 558,680 | 558,222 | 2,265 | 3,566 | 3,319 | 2,832 | 2,225 | 2,118 |
| NEW ENGLAND | 2,090 | 1,664 | 19,769 | 18,990 | 158 | 147 | 393 | 324 | 341 | 274 |
| Maine | 75 | 28 | 904 | 963 | 30 | 31 | 39 | 36 | - | - |
| N.H. | 45 | 34 | 903 | 914 | 19 | 16 | 34 | 46 | 33 | 45 |
| Vt. | 16 | 18 | 438 | 389 | 36 | 26 | 32 | 21 | 20 | 18 |
| Mass. | 1,338 | 844 | 8,616 | 7,862 | 52 | 67 | 170 | 144 | 183 | 154 |
| R.I. | 96 | 119 | 2,159 | 2,171 | 6 | 7 | 27 | 13 | 26 | 1 |
| Conn. | 520 | 621 | 6,749 | 6,691 | 15 | U | 91 | 64 | 79 | 56 |
| MID. ATLANTIC | 10,473 | 11,353 | 55,879 | 58,213 | 411 | 559 | 308 | 294 | 92 | 86 |
| Upstate N.Y. | 1,196 | 1,322 | N | N | 169 | 328 | 246 | 213 | - | - |
| N.Y. City | 5,571 | 6,520 | 21,963 | 24,764 | 116 | 206 | 11 | 14 | 17 | 13 |
| N.J. | 1,932 | 2,007 | 10,095 | 11,130 | 36 | 25 | 51 | 67 | 46 | 52 |
| Pa. | 1,774 | 1,504 | 23,821 | 22,319 | 90 | N | N | N | 29 | 21 |
| E.N. CENTRAL | 2,801 | 3,061 | 81,247 | 94,937 | 564 | 720 | 687 | 448 | 484 | 367 |
| Ohio | 448 | 645 | 26,294 | 25,697 | 66 | 71 | 246 | 123 | 199 | 76 |
| Ind. | 320 | 473 | 10,586 | 10,458 | 38 | 59 | 107 | 101 | 64 | 54 |
| Ill. | 1,345 | 1,188 | 24,169 | 25,116 | 67 | 84 | 221 | 110 | 81 | 80 |
| Mich. | 555 | 577 | 20,198 | 20,469 | 48 | 38 | 113 | 114 | 76 | 69 |
| Wis. | 133 | 178 | U | 13,197 | 345 | 468 | N | N | 64 | 88 |
| W.N. CENTRAL | 940 | 827 | 33,074 | 33,165 | 202 | 334 | 586 | 470 | 406 | 398 |
| Minn. | 178 | 163 | 6,441 | 6,660 | 78 | 142 | 229 | 195 | 178 | 209 |
| Iowa | 77 | 62 | 4,649 | 4,245 | 55 | 65 | 115 | 91 | 73 | 59 |
| Mo. | 449 | 400 | 12,427 | 11,885 | 29 | 26 | 60 | 51 | 64 | 63 |
| N. Dak. | 6 | 5 | 707 | 977 | 18 | 30 | 17 | 12 | 14 | 15 |
| S. Dak. | 15 | 15 | 1,496 | 1,477 | 7 | 25 | 47 | 35 | 62 | 38 |
| Nebr. | 65 | 66 | 3,128 | 2,657 | 14 | 35 | 97 | 50 | - | - |
| Kans. | 150 | 116 | 4,226 | 5,264 | 1 | 11 | 21 | 36 | 15 | 14 |
| S. ATLANTIC | 11,305 | 11,023 | 119,300 | 108,094 | 373 | 341 | 341 | 245 | 163 | 168 |
| Del. | 159 | 152 | 2,604 | 2,461 | - | 3 | 6 | - | 3 | 2 |
| Md. | 1,344 | 1,482 | 10,616 | 6,888 | 17 | 19 | 42 | 42 | 4 | 14 |
| D.C. | 637 | 808 | N | N | 8 | 25 | 1 | 1 | U | U |
| Va. | 782 | 908 | 13,268 | 12,983 | 27 | 20 | 73 | N | 59 | 52 |
| W. Va. | 64 | 77 | 1,240 | 2,293 | 3 | 2 | 14 | 13 | 11 | 10 |
| N.C. | 739 | 753 | 20,705 | 20,644 | 33 | N | 74 | 56 | 52 | 47 |
| S.C. | 919 | 720 | 11,346 | 16,770 | - | - | 20 | 15 | 14 | 12 |
| Ga. | 1,581 | 1,173 | 30,893 | 22,576 | 132 | 127 | 36 | 76 | - | - |
| Fla. | 5,080 | 4,950 | 28,628 | 23,479 | 153 | 145 | 75 | 42 | 20 | 31 |
| E.S. CENTRAL | 1,796 | 1,681 | 42,694 | 38,802 | 35 | 25 | 132 | 118 | 58 | 64 |
| Ky. | 255 | 262 | 7,014 | 6,083 | 7 | 10 | 46 | 35 | - | - |
| Tenn. | 706 | 621 | 13,081 | 13,021 | 11 | 9 | 54 | 53 | 38 | 40 |
| Ala. | 449 | 455 | 12,004 | 9,704 | 12 | N | 26 | 24 | 16 | 20 |
| Miss. | 386 | 343 | 10,595 | 9,994 | 5 | 6 | 6 | 6 | 4 | 4 |
| W.S. CENTRAL | 4,177 | 5,129 | 79,259 | 84,486 | 84 | 909 | 128 | 102 | 124 | 106 |
| Ark. | 188 | 189 | 5,585 | 3,871 | 2 | 6 | 15 | 11 | 8 | 10 |
| La. | 813 | 874 | 11,220 | 14,301 | 22 | 16 | 9 | 5 | 14 | 7 |
| Okla. | 123 | 274 | 7,763 | 8,878 | 12 | N | 31 | 24 | 27 | 9 |
| Tex. | 3,053 | 3,792 | 54,691 | 57,436 | 48 | 887 | 73 | 62 | 75 | 80 |
| MOUNTAIN | 1,608 | 1,478 | 29,725 | 31,557 | 98 | 122 | 320 | 360 | 224 | 246 |
| Mont. | 13 | 28 | 1,496 | 1,205 | 13 | 10 | 25 | 16 | - | 5 |
| Idaho | 22 | 28 | 1,631 | 1,917 | 8 | 17 | 65 | 41 | 43 | 25 |
| Wyo. | 11 | 3 | 741 | 665 | 1 | 2 | 15 | 53 | 14 | 55 |
| Colo. | 290 | 286 | 5,417 | 7,963 | 14 | 19 | 107 | 89 | 88 | 69 |
| N. Mex. | 82 | 203 | 3,870 | 3,699 | 42 | 47 | 13 | 19 | 6 | 20 |
| Ariz. | 819 | 588 | 11,767 | 10,890 | 12 | 18 | 37 | 43 | 23 | 26 |
| Utah | 142 | 128 | 2,021 | 2,053 | N | N | 38 | 75 | 48 | 22 |
| Nev. | 229 | 214 | 2,782 | 3,165 | 8 | 9 | 20 | 24 | 2 | 24 |
| PACIFIC | 5,743 | 6,092 | 97,733 | 89,978 | 340 | 409 | 424 | 471 | 333 | 409 |
| Wash. | 337 | 386 | 11,370 | 10,356 | N | N | 167 | 109 | 159 | 130 |
| Oreg. | 208 | 166 | 5,698 | 5,376 | 93 | 67 | 74 | 107 | 68 | 100 |
| Calif. | 5,089 | 5,364 | 76,276 | 69,991 | 247 | 338 | 171 | 248 | 94 | 163 |
| Alaska | 15 | 17 | 1,770 | 1,791 | - | 1 | 1 | 7 | 1 | - |
| Hawaii | 94 | 159 | 2,619 | 2,464 | - | 3 | 11 | - | 11 | 16 |
| Guam | 10 | 1 | 299 | 404 | - | - | N | N | U | U |
| P.R. | 1,180 | 1,601 | U | U | - | N | 9 | 5 | U | U |
| V.I. | 35 | 31 | U | U | U | U | U | U | U | U |
| Amer. Samoa | - | - | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | - | U | U | U | U | U | U | U | U |

N: Not notifiable U: Unavailable -: no reported cases C.N.M.I.: Commonwealth of Northern Mariana Islands

*Individual cases may be reported through both the National Electronic Telecommunications System for Surveillance (NETSS) and the Public Health Laboratory Information System (PHLIS).

†Updated monthly from reports to the Division of HIV/AIDS Prevention—Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention, last update November 28, 1999.

TABLE II. (Cont'd.) Provisional cases of selected notifiable diseases, United States, weeks ending December 11, 1999, and December 12, 1998 (49th Week)

| Reporting Area | Gonorrhea | | Hepatitis C/NA,NB | | Legionellosis | | Lyme Disease | |
|----------------|-----------|-----------|-------------------|-----------|---------------|-----------|--------------|-----------|
| | Cum. 1999 | Cum. 1998 | Cum. 1999 | Cum. 1998 | Cum. 1999 | Cum. 1998 | Cum. 1999 | Cum. 1998 |
| UNITED STATES | 306,829 | 333,630 | 2,992 | 3,151 | 896 | 1,221 | 12,207 | 15,094 |
| NEW ENGLAND | 6,177 | 5,724 | 14 | 58 | 78 | 84 | 3,389 | 4,533 |
| Maine | 71 | 63 | 2 | - | 3 | 1 | 41 | 78 |
| N.H. | 105 | 88 | - | - | 8 | 7 | 23 | 43 |
| Vt. | 44 | 35 | 7 | 6 | 14 | 7 | 23 | 11 |
| Mass. | 2,383 | 2,147 | 2 | 49 | 28 | 33 | 945 | 690 |
| R.I. | 543 | 394 | 3 | 3 | 11 | 21 | 464 | 650 |
| Conn. | 3,031 | 2,997 | - | - | 14 | 15 | 1,893 | 3,061 |
| MID. ATLANTIC | 35,884 | 36,402 | 92 | 205 | 182 | 309 | 6,920 | 8,402 |
| Upstate N.Y. | 6,395 | 6,923 | 57 | 102 | 56 | 107 | 3,760 | 3,910 |
| N.Y. City | 11,762 | 11,305 | - | - | 9 | 35 | 39 | 230 |
| N.J. | 5,962 | 7,491 | - | U | 18 | 18 | 922 | 1,802 |
| Pa. | 11,765 | 10,683 | 35 | 103 | 99 | 149 | 2,199 | 2,460 |
| E.N. CENTRAL | 53,864 | 65,262 | 1,423 | 648 | 243 | 398 | 176 | 754 |
| Ohio | 15,957 | 16,870 | 4 | 8 | 79 | 125 | 73 | 46 |
| Ind. | 5,791 | 6,132 | 1 | 5 | 43 | 75 | 21 | 37 |
| Ill. | 17,967 | 20,791 | 41 | 40 | 23 | 52 | 12 | 14 |
| Mich. | 14,149 | 15,424 | 786 | 455 | 60 | 80 | 1 | 12 |
| Wis. | U | 6,045 | 591 | 140 | 38 | 66 | 69 | 645 |
| W.N. CENTRAL | 14,198 | 16,724 | 299 | 43 | 51 | 63 | 288 | 226 |
| Minn. | 2,484 | 2,578 | 10 | 11 | 13 | 7 | 220 | 173 |
| Iowa | 1,155 | 1,415 | - | 8 | 15 | 10 | 19 | 26 |
| Mo. | 7,179 | 8,847 | 277 | 15 | 14 | 16 | 26 | 12 |
| N. Dak. | 71 | 77 | 1 | - | 2 | - | 1 | - |
| S. Dak. | 186 | 209 | - | - | 3 | 3 | - | - |
| Nebr. | 1,297 | 1,120 | 5 | 5 | 4 | 19 | 10 | 4 |
| Kans. | 1,826 | 2,478 | 6 | 4 | - | 8 | 12 | 11 |
| S. ATLANTIC | 89,820 | 89,821 | 193 | 115 | 146 | 140 | 1,123 | 867 |
| Del. | 1,582 | 1,454 | 1 | - | 14 | 13 | 64 | 66 |
| Md. | 9,012 | 9,135 | 41 | 21 | 32 | 35 | 785 | 608 |
| D.C. | 3,316 | 4,009 | 1 | - | 4 | 8 | 6 | 4 |
| Va. | 9,015 | 9,106 | 11 | 12 | 38 | 20 | 118 | 68 |
| W. Va. | 387 | 824 | 17 | 7 | N | N | 17 | 13 |
| N.C. | 18,440 | 17,841 | 34 | 25 | 15 | 14 | 73 | 57 |
| S.C. | 6,744 | 10,728 | 22 | 11 | 11 | 11 | 7 | 7 |
| Ga. | 20,955 | 18,686 | 1 | 9 | 3 | 8 | - | 5 |
| Fla. | 20,369 | 18,038 | 65 | 30 | 29 | 31 | 53 | 39 |
| E.S. CENTRAL | 34,186 | 37,438 | 243 | 267 | 45 | 64 | 92 | 111 |
| Ky. | 3,192 | 3,577 | 21 | 20 | 20 | 26 | 10 | 26 |
| Tenn. | 10,498 | 11,366 | 95 | 160 | 21 | 23 | 50 | 44 |
| Ala. | 10,812 | 12,322 | 1 | 4 | 4 | 8 | 19 | 24 |
| Miss. | 9,684 | 10,173 | 126 | 83 | - | 7 | 13 | 17 |
| W.S. CENTRAL | 43,893 | 52,174 | 314 | 543 | 23 | 31 | 43 | 31 |
| Ark. | 2,984 | 3,800 | 18 | 22 | - | 1 | 4 | 7 |
| La. | 8,880 | 12,326 | 102 | 112 | 2 | 4 | - | 7 |
| Okla. | 3,792 | 4,960 | 15 | 16 | 3 | 12 | 4 | 2 |
| Tex. | 28,237 | 31,088 | 179 | 393 | 18 | 14 | 35 | 15 |
| MOUNTAIN | 8,881 | 8,665 | 146 | 362 | 47 | 71 | 18 | 18 |
| Mont. | 54 | 44 | 5 | 7 | - | 2 | - | - |
| Idaho | 80 | 168 | 7 | 86 | 3 | 2 | 5 | 6 |
| Wyo. | 34 | 33 | 45 | 90 | - | 1 | 3 | 1 |
| Colo. | 2,316 | 1,956 | 22 | 31 | 12 | 18 | - | - |
| N. Mex. | 802 | 894 | 8 | 96 | 1 | 2 | 1 | 4 |
| Ariz. | 4,185 | 3,982 | 45 | 11 | 7 | 17 | 2 | 1 |
| Utah | 216 | 217 | 6 | 21 | 18 | 21 | 5 | - |
| Nev. | 1,194 | 1,371 | 8 | 20 | 6 | 8 | 2 | 6 |
| PACIFIC | 19,926 | 21,420 | 268 | 910 | 81 | 61 | 158 | 152 |
| Wash. | 2,013 | 1,850 | 20 | 22 | 17 | 12 | 10 | 7 |
| Oreg. | 827 | 803 | 22 | 19 | N | N | 14 | 21 |
| Calif. | 16,436 | 17,987 | 226 | 815 | 63 | 47 | 134 | 123 |
| Alaska | 275 | 300 | - | - | 1 | 1 | - | 1 |
| Hawaii | 375 | 480 | - | 54 | - | 1 | N | N |
| Guam | 38 | 67 | 1 | 1 | - | 2 | - | 1 |
| P.R. | 328 | 363 | - | - | - | - | N | N |
| V.I. | U | U | U | U | U | U | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U |
| C.N.M.I. | U | U | U | U | U | U | U | U |

N: Not notifiable

U: Unavailable

-: no reported cases

TABLE II. (Cont'd.) Provisional cases of selected notifiable diseases, United States, weeks ending December 11, 1999, and December 12, 1998 (49th Week)

| Reporting Area | Malaria | | Rabies, Animal | | Salmonellosis* | | | |
|----------------|-----------|-----------|----------------|-----------|----------------|-----------|-----------|-----------|
| | Cum. 1999 | Cum. 1998 | Cum. 1999 | Cum. 1998 | NETSS | | PHLIS | |
| | | | | | Cum. 1999 | Cum. 1998 | Cum. 1999 | Cum. 1998 |
| UNITED STATES | 1,271 | 1,408 | 5,668 | 6,928 | 36,293 | 40,176 | 29,030 | 32,380 |
| NEW ENGLAND | 63 | 69 | 862 | 1,397 | 2,086 | 2,419 | 2,025 | 2,205 |
| Maine | 3 | 5 | 171 | 232 | 128 | 162 | 99 | 64 |
| N.H. | 2 | 5 | 50 | 77 | 136 | 178 | 140 | 215 |
| Vt. | 4 | 1 | 88 | 65 | 91 | 138 | 85 | 109 |
| Mass. | 24 | 26 | 216 | 489 | 1,113 | 1,267 | 1,118 | 1,296 |
| R.I. | 5 | 14 | 93 | 97 | 121 | 142 | 147 | 34 |
| Conn. | 25 | 18 | 244 | 437 | 497 | 532 | 436 | 487 |
| MID. ATLANTIC | 320 | 406 | 1,092 | 1,536 | 4,610 | 6,313 | 4,082 | 5,566 |
| Upstate N.Y. | 67 | 87 | 776 | 1,055 | 1,305 | 1,536 | 1,268 | 1,309 |
| N.Y. City | 167 | 230 | U | U | 1,298 | 1,825 | 1,173 | 1,407 |
| N.J. | 48 | 56 | 166 | 213 | 989 | 1,402 | 685 | 1,334 |
| Pa. | 38 | 33 | 150 | 268 | 1,018 | 1,550 | 956 | 1,516 |
| E.N. CENTRAL | 140 | 141 | 146 | 123 | 5,153 | 6,024 | 3,273 | 4,656 |
| Ohio | 18 | 15 | 36 | 57 | 1,257 | 1,445 | 1,011 | 1,103 |
| Ind. | 19 | 10 | 13 | 12 | 512 | 645 | 406 | 509 |
| Ill. | 54 | 57 | 10 | N | 1,495 | 1,853 | 399 | 1,512 |
| Mich. | 39 | 47 | 87 | 35 | 920 | 1,115 | 906 | 1,041 |
| Wis. | 10 | 12 | - | 19 | 969 | 966 | 551 | 491 |
| W.N. CENTRAL | 72 | 91 | 664 | 686 | 2,120 | 2,191 | 2,183 | 2,251 |
| Minn. | 41 | 56 | 107 | 114 | 619 | 550 | 657 | 636 |
| Iowa | 13 | 7 | 153 | 147 | 264 | 352 | 197 | 285 |
| Mo. | 14 | 14 | 14 | 41 | 689 | 592 | 876 | 820 |
| N. Dak. | - | 2 | 137 | 138 | 51 | 59 | 49 | 67 |
| S. Dak. | - | 1 | 163 | 151 | 93 | 120 | 115 | 127 |
| Nebr. | - | 1 | 3 | 7 | 185 | 174 | 78 | 46 |
| Kans. | 4 | 10 | 87 | 88 | 219 | 344 | 211 | 270 |
| S. ATLANTIC | 341 | 302 | 2,031 | 2,248 | 8,560 | 8,249 | 6,002 | 5,934 |
| Del. | 1 | 3 | 43 | 49 | 138 | 74 | 153 | 116 |
| Md. | 93 | 86 | 381 | 424 | 841 | 877 | 952 | 866 |
| D.C. | 18 | 19 | - | - | 69 | 83 | U | U |
| Va. | 70 | 56 | 554 | 534 | 1,206 | 1,057 | 943 | 835 |
| W. Va. | 3 | 2 | 106 | 76 | 163 | 147 | 148 | 158 |
| N.C. | 31 | 29 | 404 | 538 | 1,269 | 1,243 | 1,243 | 1,383 |
| S.C. | 17 | 6 | 133 | 143 | 675 | 605 | 479 | 527 |
| Ga. | 28 | 36 | 231 | 290 | 1,474 | 1,631 | 1,644 | 1,494 |
| Fla. | 80 | 65 | 179 | 194 | 2,725 | 2,532 | 440 | 555 |
| E.S. CENTRAL | 24 | 32 | 252 | 264 | 1,995 | 2,245 | 1,062 | 1,528 |
| Ky. | 7 | 7 | 35 | 31 | 393 | 347 | - | 124 |
| Tenn. | 8 | 16 | 93 | 135 | 513 | 574 | 509 | 686 |
| Ala. | 7 | 6 | 123 | 96 | 575 | 668 | 476 | 561 |
| Miss. | 2 | 3 | 1 | 2 | 514 | 656 | 77 | 157 |
| W.S. CENTRAL | 16 | 54 | 94 | 28 | 3,598 | 4,699 | 3,546 | 3,102 |
| Ark. | 3 | 1 | 14 | 28 | 626 | 589 | 120 | 367 |
| La. | 10 | 14 | - | - | 334 | 744 | 568 | 787 |
| Okla. | 2 | 3 | 80 | N | 406 | 468 | 320 | 225 |
| Tex. | 1 | 36 | - | - | 2,232 | 2,898 | 2,538 | 1,723 |
| MOUNTAIN | 43 | 61 | 197 | 246 | 2,918 | 2,435 | 2,411 | 1,938 |
| Mont. | 4 | 1 | 59 | 53 | 81 | 76 | 1 | 43 |
| Idaho | 3 | 8 | 5 | N | 125 | 118 | 98 | 94 |
| Wyo. | 1 | - | 44 | 64 | 67 | 63 | 49 | 57 |
| Colo. | 17 | 18 | 1 | 42 | 679 | 518 | 689 | 488 |
| N. Mex. | 2 | 12 | 9 | 6 | 362 | 288 | 245 | 255 |
| Ariz. | 8 | 9 | 66 | 48 | 913 | 798 | 762 | 663 |
| Utah | 4 | 1 | 8 | 27 | 506 | 341 | 514 | 122 |
| Nev. | 4 | 12 | 5 | 6 | 185 | 233 | 53 | 216 |
| PACIFIC | 252 | 252 | 330 | 400 | 5,253 | 5,601 | 4,446 | 5,200 |
| Wash. | 27 | 20 | - | - | 634 | 493 | 795 | 666 |
| Oreg. | 21 | 15 | 2 | 7 | 409 | 314 | 480 | 322 |
| Calif. | 192 | 207 | 321 | 370 | 3,833 | 4,457 | 2,875 | 3,881 |
| Alaska | 1 | 3 | 7 | 23 | 53 | 56 | 30 | 36 |
| Hawaii | 11 | 7 | - | - | 324 | 281 | 266 | 295 |
| Guam | - | 2 | - | - | 24 | 42 | U | U |
| P.R. | - | - | 66 | 49 | 433 | 769 | U | U |
| V.I. | U | U | U | U | U | U | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U |
| C.N.M.I. | U | U | U | U | U | U | U | U |

N: Not notifiable U: Unavailable -: no reported cases

*Individual cases may be reported through both the National Electronic Telecommunications System for Surveillance (NETSS) and the Public Health Laboratory Information System (PHLIS).

TABLE II. (Cont'd.) Provisional cases of selected notifiable diseases, United States, weeks ending December 11, 1999, and December 12, 1998 (49th Week)

| Reporting Area | Shigellosis* | | | | Syphilis (Primary & Secondary) | | Tuberculosis | |
|----------------|--------------|--------------|--------------|--------------|-----------------------------------|--------------|---------------|---------------|
| | NETSS | | PHLIS | | Cum. 1999 | Cum. 1998 | Cum. 1999† | Cum. 1998† |
| | Cum. 1999 | Cum. 1998 | Cum. 1999 | Cum. 1998 | | | | |
| UNITED STATES | 15,097 | 21,128 | 7,476 | 11,831 | 6,146 | 6,697 | 13,220 | 16,101 |
| NEW ENGLAND | 821 | 401 | 786 | 355 | 57 | 76 | 406 | 414 |
| Maine | 5 | 14 | - | - | - | 1 | 18 | 11 |
| N.H. | 17 | 16 | 17 | 20 | 1 | 2 | 10 | - |
| Vt. | 6 | 7 | 4 | 4 | 3 | 4 | 2 | 5 |
| Mass. | 703 | 258 | 687 | 253 | 35 | 43 | 232 | 239 |
| R.I. | 23 | 36 | 18 | 13 | 2 | 1 | 39 | 52 |
| Conn. | 67 | 70 | 60 | 65 | 16 | 25 | 105 | 107 |
| MID. ATLANTIC | 897 | 2,296 | 454 | 1,657 | 186 | 310 | 2,380 | 2,882 |
| Upstate N.Y. | 266 | 613 | 67 | 220 | 23 | 36 | 304 | 360 |
| N.Y. City | 281 | 689 | 82 | 575 | 79 | 79 | 1,264 | 1,363 |
| N.J. | 194 | 650 | 155 | 608 | 51 | 101 | 479 | 583 |
| Pa. | 156 | 344 | 150 | 254 | 33 | 94 | 333 | 576 |
| E.N. CENTRAL | 2,843 | 2,842 | 1,274 | 1,517 | 1,328 | 978 | 1,186 | 1,588 |
| Ohio | 411 | 495 | 136 | 141 | 87 | 128 | 228 | 221 |
| Ind. | 324 | 171 | 101 | 43 | 646 | 201 | 93 | 152 |
| Ill. | 1,048 | 1,518 | 592 | 1,261 | 365 | 396 | 508 | 766 |
| Mich. | 474 | 262 | 368 | 4 | 230 | 194 | 272 | 344 |
| Wis. | 586 | 396 | 77 | 68 | U | 59 | 85 | 105 |
| W.N. CENTRAL | 1,069 | 1,035 | 721 | 600 | 108 | 131 | 447 | 467 |
| Minn. | 238 | 298 | 229 | 325 | 9 | 9 | 187 | 146 |
| Iowa | 66 | 66 | 48 | 45 | 9 | 2 | 50 | 51 |
| Mo. | 638 | 190 | 352 | 129 | 72 | 99 | 152 | 163 |
| N. Dak. | 3 | 10 | 2 | 3 | - | - | 6 | 10 |
| S. Dak. | 18 | 32 | 10 | 23 | - | 1 | 17 | 17 |
| Nebr. | 69 | 367 | 35 | 19 | 8 | 7 | 16 | 28 |
| Kans. | 37 | 72 | 45 | 56 | 10 | 13 | 19 | 52 |
| S. ATLANTIC | 2,385 | 4,132 | 485 | 1,233 | 1,925 | 2,439 | 2,784 | 3,032 |
| Del. | 13 | 44 | 9 | 37 | 8 | 21 | 12 | 34 |
| Md. | 157 | 197 | 58 | 66 | 310 | 643 | 248 | 279 |
| D.C. | 51 | 37 | U | U | 59 | 85 | 47 | 102 |
| Va. | 129 | 192 | 61 | 87 | 148 | 144 | 265 | 280 |
| W. Va. | 8 | 11 | 5 | 8 | 2 | 3 | 37 | 41 |
| N.C. | 200 | 339 | 86 | 179 | 421 | 691 | 394 | 448 |
| S.C. | 123 | 178 | 62 | 94 | 245 | 309 | 218 | 270 |
| Ga. | 227 | 1,051 | 85 | 240 | 396 | 276 | 556 | 514 |
| Fla. | 1,477 | 2,083 | 119 | 522 | 336 | 267 | 1,007 | 1,064 |
| E.S. CENTRAL | 1,064 | 1,445 | 483 | 1,123 | 1,084 | 1,163 | 847 | 1,152 |
| Ky. | 229 | 145 | - | 45 | 99 | 103 | 166 | 157 |
| Tenn. | 600 | 801 | 426 | 852 | 602 | 545 | 334 | 436 |
| Ala. | 111 | 445 | 47 | 219 | 202 | 270 | 291 | 355 |
| Miss. | 124 | 54 | 10 | 7 | 181 | 245 | 56 | 204 |
| W.S. CENTRAL | 2,438 | 4,434 | 2,337 | 1,392 | 898 | 1,022 | 1,462 | 2,328 |
| Ark. | 74 | 201 | 23 | 61 | 79 | 107 | 161 | 143 |
| La. | 118 | 332 | 128 | 281 | 208 | 409 | U | 278 |
| Okla. | 456 | 617 | 153 | 191 | 175 | 92 | 122 | 155 |
| Tex. | 1,790 | 3,284 | 2,033 | 859 | 436 | 414 | 1,179 | 1,752 |
| MOUNTAIN | 1,127 | 1,246 | 722 | 728 | 223 | 229 | 427 | 534 |
| Mont. | 9 | 8 | - | 3 | 1 | - | 13 | 19 |
| Idaho | 28 | 19 | 12 | 14 | 1 | 2 | 15 | 11 |
| Wyo. | 3 | 3 | 1 | 1 | - | 1 | 3 | 4 |
| Colo. | 193 | 222 | 155 | 159 | 2 | 10 | U | 67 |
| N. Mex. | 139 | 289 | 89 | 173 | 11 | 22 | 59 | 65 |
| Ariz. | 599 | 594 | 395 | 324 | 200 | 175 | 215 | 205 |
| Utah | 66 | 46 | 64 | 34 | 2 | 4 | 40 | 48 |
| Nev. | 90 | 65 | 6 | 20 | 6 | 15 | 82 | 115 |
| PACIFIC | 2,453 | 3,297 | 214 | 3,226 | 337 | 349 | 3,281 | 3,704 |
| Wash. | 117 | 219 | 99 | 188 | 64 | 27 | 168 | 242 |
| Oreg. | 95 | 190 | 85 | 151 | 10 | 5 | 99 | 126 |
| Calif. | 2,205 | 2,830 | - | 2,830 | 259 | 313 | 2,793 | 3,119 |
| Alaska | 3 | 9 | 3 | 7 | 1 | 1 | 53 | 51 |
| Hawaii | 33 | 49 | 27 | 50 | 3 | 3 | 168 | 166 |
| Guam | 8 | 36 | U | U | 1 | 1 | 11 | 84 |
| P.R. | 106 | 62 | U | U | 151 | 167 | 41 | 140 |
| V.I. | U | U | U | U | U | U | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U |
| C.N.M.I. | U | U | U | U | U | U | U | U |

N: Not notifiable U: Unavailable -: no reported cases

*Individual cases may be reported through both the National Electronic Telecommunications System for Surveillance (NETSS) and the Public Health Laboratory Information System (PHLIS).

†Cumulative reports of provisional tuberculosis cases for 1999 are unavailable ("U") for some areas using the Tuberculosis Information System (TIMS).

TABLE III. Provisional cases of selected notifiable diseases preventable by vaccination, United States, weeks ending December 11, 1999, and December 12, 1998 (49th Week)

| Reporting Area | <i>H. influenzae</i> , invasive | | Hepatitis (Viral), by type | | | | Measles (Rubeola) | | | | | |
|----------------|---------------------------------|-----------|----------------------------|-----------|-----------|-----------|-------------------|-----------|-----------|-----------|-----------|-----------|
| | Cum. 1999† | Cum. 1998 | A | | B | | Indigenous | | Imported* | | Total | |
| | | | Cum. 1999 | Cum. 1998 | Cum. 1999 | Cum. 1998 | 1999 | Cum. 1999 | 1999 | Cum. 1999 | Cum. 1999 | Cum. 1998 |
| UNITED STATES | 1,088 | 1,003 | 15,794 | 21,068 | 6,003 | 9,059 | - | 60 | 1 | 25 | 85 | 90 |
| NEW ENGLAND | 94 | 68 | 283 | 281 | 134 | 212 | - | 6 | - | 5 | 11 | 3 |
| Maine | 8 | 3 | 14 | 20 | 1 | 5 | - | - | - | - | - | - |
| N.H. | 21 | 10 | 18 | 15 | 16 | 19 | - | - | - | 1 | 1 | - |
| Vt. | 5 | 9 | 19 | 17 | 3 | 10 | - | - | - | - | - | 1 |
| Mass. | 36 | 39 | 108 | 119 | 41 | 77 | - | 5 | - | 3 | 8 | 2 |
| R.I. | 6 | 6 | 21 | 17 | 34 | 68 | U | - | U | - | - | - |
| Conn. | 18 | 1 | 103 | 93 | 39 | 33 | - | 1 | - | 1 | 2 | - |
| MID. ATLANTIC | 169 | 166 | 913 | 1,638 | 555 | 1,167 | - | - | - | 2 | 2 | 14 |
| Upstate N.Y. | 76 | 62 | 256 | 346 | 172 | 231 | - | - | - | 2 | 2 | 2 |
| N.Y. City | 41 | 43 | 300 | 579 | 186 | 409 | - | - | - | - | - | - |
| N.J. | 49 | 51 | 112 | 331 | 41 | 194 | U | - | U | - | - | 8 |
| Pa. | 3 | 10 | 245 | 382 | 156 | 333 | - | - | - | - | - | 4 |
| E.N. CENTRAL | 159 | 171 | 2,628 | 3,441 | 625 | 1,362 | - | 1 | - | 2 | 3 | 16 |
| Ohio | 56 | 46 | 628 | 312 | 88 | 74 | - | - | - | - | - | 1 |
| Ind. | 23 | 43 | 107 | 156 | 43 | 107 | - | 1 | - | 1 | 2 | 3 |
| Ill. | 66 | 62 | 646 | 761 | 1 | 225 | - | - | - | - | - | 1 |
| Mich. | 13 | 13 | 1,180 | 2,029 | 469 | 463 | - | - | - | 1 | 1 | 10 |
| Wis. | 1 | 7 | 67 | 183 | 24 | 493 | - | - | - | - | - | 1 |
| W.N. CENTRAL | 88 | 87 | 874 | 1,273 | 344 | 391 | - | 1 | - | - | 1 | - |
| Minn. | 47 | 66 | 95 | 124 | 54 | 49 | - | 1 | - | - | 1 | - |
| Iowa | 10 | 3 | 143 | 394 | 39 | 53 | - | - | - | - | - | - |
| Mo. | 22 | 10 | 534 | 590 | 207 | 235 | - | - | - | - | - | - |
| N. Dak. | 1 | - | 3 | 3 | 2 | 4 | - | - | - | - | - | - |
| S. Dak. | 1 | 1 | 9 | 32 | 1 | 2 | - | - | - | - | - | - |
| Nebr. | 3 | 1 | 50 | 26 | 14 | 21 | - | - | - | - | - | - |
| Kans. | 4 | 6 | 40 | 104 | 27 | 27 | U | - | U | - | - | - |
| S. ATLANTIC | 252 | 176 | 1,950 | 1,925 | 1,168 | 991 | - | 14 | - | 6 | 20 | 8 |
| Del. | - | 1 | 2 | 6 | 1 | 4 | - | - | - | - | - | 1 |
| Md. | 66 | 52 | 339 | 394 | 165 | 132 | - | - | - | - | - | 1 |
| D.C. | 5 | - | 58 | 64 | 24 | 18 | U | - | U | - | - | - |
| Va. | 20 | 18 | 171 | 199 | 96 | 99 | - | 14 | - | 4 | 18 | 2 |
| W. Va. | 7 | 6 | 39 | 7 | 23 | 10 | - | - | - | - | - | - |
| N.C. | 35 | 24 | 156 | 123 | 212 | 227 | - | - | - | - | - | - |
| S.C. | 6 | 3 | 47 | 38 | 65 | 46 | - | - | - | - | - | - |
| Ga. | 67 | 44 | 446 | 638 | 159 | 138 | - | - | - | - | - | 2 |
| Fla. | 46 | 28 | 692 | 456 | 423 | 317 | - | - | - | 2 | 2 | 2 |
| E.S. CENTRAL | 62 | 61 | 390 | 382 | 414 | 479 | - | 2 | - | - | 2 | 2 |
| Ky. | 7 | 7 | 62 | 30 | 42 | 47 | - | 2 | - | - | 2 | - |
| Tenn. | 35 | 36 | 174 | 211 | 211 | 266 | - | - | - | - | - | 1 |
| Ala. | 17 | 15 | 50 | 73 | 78 | 72 | - | - | - | - | - | 1 |
| Miss. | 3 | 3 | 104 | 68 | 83 | 94 | - | - | - | - | - | - |
| W.S. CENTRAL | 46 | 53 | 3,612 | 3,857 | 803 | 1,993 | - | 10 | - | 4 | 14 | - |
| Ark. | 2 | - | 68 | 79 | 69 | 104 | - | 5 | - | - | 5 | - |
| La. | 7 | 21 | 73 | 114 | 77 | 163 | U | - | U | - | - | - |
| Okla. | 33 | 29 | 435 | 591 | 129 | 108 | - | - | - | - | - | - |
| Tex. | 4 | 3 | 3,036 | 3,073 | 528 | 1,618 | - | 5 | - | 4 | 9 | - |
| MOUNTAIN | 105 | 110 | 1,231 | 2,994 | 543 | 783 | - | 4 | - | - | 4 | 5 |
| Mont. | 3 | - | 17 | 93 | 17 | 5 | - | - | - | - | - | - |
| Idaho | 1 | 2 | 43 | 231 | 29 | 46 | - | - | - | - | - | - |
| Wyo. | 1 | 1 | 7 | 37 | 13 | 9 | - | - | - | - | - | - |
| Colo. | 11 | 21 | 206 | 324 | 91 | 102 | - | - | - | - | - | - |
| N. Mex. | 18 | 7 | 50 | 147 | 169 | 306 | - | - | - | - | - | - |
| Ariz. | 56 | 55 | 715 | 1,760 | 139 | 170 | - | 1 | - | - | 1 | 5 |
| Utah | 11 | 5 | 62 | 186 | 37 | 65 | - | 2 | - | - | 2 | - |
| Nev. | 4 | 19 | 131 | 216 | 48 | 80 | U | 1 | U | - | 1 | - |
| PACIFIC | 113 | 111 | 3,913 | 5,277 | 1,417 | 1,681 | - | 22 | 1 | 6 | 28 | 42 |
| Wash. | 7 | 9 | 372 | 927 | 73 | 108 | - | - | - | - | - | 1 |
| Oreg. | 40 | 40 | 238 | 422 | 100 | 193 | - | 9 | - | - | 9 | - |
| Calif. | 48 | 49 | 3,271 | 3,858 | 1,213 | 1,352 | - | 13 | - | 4 | 17 | 8 |
| Alaska | 9 | 4 | 12 | 17 | 17 | 13 | - | - | - | - | - | 33 |
| Hawaii | 9 | 9 | 20 | 53 | 14 | 15 | - | - | 1 | 2 | 2 | - |
| Guam | - | - | 2 | 1 | 2 | 2 | U | 1 | U | - | 1 | - |
| P.R. | 1 | 2 | 187 | 79 | 145 | 240 | - | - | - | - | - | - |
| V.I. | U | U | U | U | U | U | U | U | U | U | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | U | U | U | U | U | U | U | U | U | U | U | U |

N: Not notifiable U: Unavailable -: no reported cases

*For imported measles, cases include only those resulting from importation from other countries.

†Of 212 cases among children aged <5 years, serotype was reported for 107 and of those, 31 were type b.

TABLE III. (Cont'd.) Provisional cases of selected notifiable diseases preventable by vaccination, United States, weeks ending December 11, 1999, and December 12, 1998 (49th Week)

| Reporting Area | Meningococcal Disease | | Mumps | | | Pertussis | | | Rubella | | |
|----------------|-----------------------|-----------|-------|-----------|-----------|-----------|-----------|-----------|---------|-----------|-----------|
| | Cum. 1999 | Cum. 1998 | 1999 | Cum. 1999 | Cum. 1998 | 1999 | Cum. 1999 | Cum. 1998 | 1999 | Cum. 1999 | Cum. 1998 |
| UNITED STATES | 2,197 | 2,484 | 4 | 322 | 614 | 127 | 5,560 | 6,494 | 1 | 232 | 352 |
| NEW ENGLAND | 107 | 112 | - | 8 | 9 | 7 | 686 | 1,009 | - | 7 | 38 |
| Maine | 5 | 6 | - | - | - | - | - | 5 | - | - | - |
| N.H. | 13 | 12 | - | 1 | - | - | 78 | 121 | - | - | - |
| Vt. | 5 | 5 | - | 1 | - | 4 | 75 | 76 | - | - | - |
| Mass. | 61 | 56 | - | 4 | 6 | 3 | 469 | 748 | - | 7 | 8 |
| R.I. | 7 | 8 | U | 2 | 1 | U | 33 | 13 | U | - | 1 |
| Conn. | 16 | 25 | - | - | 2 | - | 31 | 46 | - | - | 29 |
| MID. ATLANTIC | 204 | 266 | 2 | 35 | 191 | 18 | 913 | 620 | - | 25 | 149 |
| Upstate N.Y. | 64 | 76 | 1 | 14 | 12 | 11 | 723 | 317 | - | 21 | 114 |
| N.Y. City | 50 | 32 | - | 3 | 155 | - | 10 | 46 | - | - | 19 |
| N.J. | 47 | 57 | U | - | 6 | U | 12 | 28 | U | 1 | 14 |
| Pa. | 43 | 101 | 1 | 18 | 18 | 7 | 168 | 229 | - | 3 | 2 |
| E.N. CENTRAL | 372 | 379 | - | 43 | 77 | 48 | 542 | 830 | - | 2 | - |
| Ohio | 126 | 133 | - | 18 | 28 | 44 | 268 | 279 | - | - | - |
| Ind. | 67 | 72 | - | 5 | 7 | 1 | 74 | 173 | - | 1 | - |
| Ill. | 96 | 99 | - | 11 | 10 | 1 | 82 | 127 | - | 1 | - |
| Mich. | 45 | 44 | - | 7 | 29 | 2 | 66 | 69 | - | - | - |
| Wis. | 38 | 31 | - | 2 | 3 | - | 52 | 182 | - | - | - |
| W.N. CENTRAL | 231 | 216 | - | 13 | 32 | 17 | 421 | 574 | - | 124 | 40 |
| Minn. | 50 | 32 | - | 1 | 13 | 17 | 226 | 337 | - | 5 | - |
| Iowa | 43 | 43 | - | 7 | 11 | - | 70 | 71 | - | 29 | - |
| Mo. | 93 | 76 | - | 1 | 3 | - | 61 | 35 | - | 3 | 2 |
| N. Dak. | 4 | 5 | - | 1 | 2 | - | 18 | 4 | - | - | - |
| S. Dak. | 11 | 8 | - | - | - | - | 7 | 8 | - | - | - |
| Nebr. | 12 | 17 | - | - | - | - | 4 | 17 | - | 87 | - |
| Kans. | 18 | 35 | U | 3 | 3 | U | 35 | 102 | U | - | 38 |
| S. ATLANTIC | 403 | 427 | 1 | 50 | 47 | 7 | 414 | 322 | 1 | 37 | 19 |
| Del. | 8 | 2 | - | - | - | - | 5 | 5 | - | - | - |
| Md. | 54 | 34 | - | 7 | - | 1 | 108 | 63 | - | 1 | 1 |
| D.C. | 2 | 3 | U | 2 | - | U | 1 | 1 | U | - | - |
| Va. | 53 | 45 | - | 10 | 8 | - | 51 | 41 | - | - | 1 |
| W. Va. | 8 | 17 | - | - | - | - | 3 | 4 | - | - | - |
| N.C. | 46 | 57 | - | 8 | 11 | 3 | 93 | 98 | - | 35 | 13 |
| S.C. | 43 | 55 | 1 | 5 | 7 | - | 18 | 27 | - | - | - |
| Ga. | 59 | 97 | - | 4 | 1 | - | 40 | 27 | - | - | - |
| Fla. | 130 | 117 | - | 14 | 20 | 3 | 95 | 56 | 1 | 1 | 4 |
| E.S. CENTRAL | 144 | 195 | - | 13 | 18 | - | 89 | 148 | - | 1 | 2 |
| Ky. | 31 | 37 | - | - | 1 | - | 25 | 79 | - | - | - |
| Tenn. | 59 | 68 | - | - | 2 | - | 40 | 37 | - | - | 2 |
| Ala. | 32 | 53 | - | 10 | 8 | - | 21 | 26 | - | 1 | - |
| Miss. | 22 | 37 | - | 3 | 7 | - | 3 | 6 | - | - | - |
| W.S. CENTRAL | 174 | 290 | - | 33 | 59 | 1 | 158 | 359 | - | 15 | 88 |
| Ark. | 35 | 30 | - | - | 13 | 1 | 19 | 82 | - | 6 | - |
| La. | 34 | 55 | U | 3 | 7 | U | 3 | 9 | U | - | - |
| Okla. | 31 | 40 | - | 1 | - | - | 12 | 32 | - | - | - |
| Tex. | 74 | 165 | - | 29 | 39 | - | 124 | 236 | - | 9 | 88 |
| MOUNTAIN | 137 | 141 | - | 28 | 39 | 21 | 737 | 1,169 | - | 16 | 5 |
| Mont. | 4 | 4 | - | - | - | - | 2 | 13 | - | - | - |
| Idaho | 13 | 13 | - | 3 | 7 | - | 139 | 232 | - | - | - |
| Wyo. | 5 | 8 | - | - | 1 | - | 2 | 8 | - | - | - |
| Colo. | 35 | 28 | - | 5 | 6 | 8 | 207 | 324 | - | 1 | - |
| N. Mex. | 14 | 26 | N | N | N | 9 | 200 | 98 | - | - | 1 |
| Ariz. | 42 | 39 | - | 8 | 6 | 4 | 117 | 191 | - | 13 | 1 |
| Utah | 16 | 13 | - | 7 | 5 | - | 59 | 262 | - | 1 | 2 |
| Nev. | 8 | 10 | U | 5 | 14 | U | 11 | 41 | U | 1 | 1 |
| PACIFIC | 425 | 458 | 1 | 99 | 142 | 8 | 1,600 | 1,463 | - | 5 | 11 |
| Wash. | 63 | 64 | - | 2 | 11 | 6 | 609 | 329 | - | - | 6 |
| Oreg. | 77 | 85 | N | N | N | - | 58 | 89 | - | - | - |
| Calif. | 271 | 301 | 1 | 82 | 104 | 2 | 894 | 1,005 | - | 5 | 3 |
| Alaska | 6 | 3 | - | 3 | 3 | - | 5 | 15 | - | - | - |
| Hawaii | 8 | 5 | - | 12 | 24 | - | 34 | 25 | - | - | 2 |
| Guam | 2 | 2 | U | 1 | 5 | U | 1 | 1 | U | - | - |
| P.R. | 7 | 11 | - | - | 7 | 1 | 20 | 9 | - | - | 14 |
| V.I. | U | U | U | U | U | U | U | U | U | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | U | U | U | U | U | U | U | U | U | U | U |

N: Not notifiable

U: Unavailable

-: no reported cases

TABLE IV. Deaths in 122 U.S. cities,* week ending
December 11, 1999 (49th Week)

| Reporting Area | All Causes, By Age (Years) | | | | | | P&J [†] Total | Reporting Area | All Causes, By Age (Years) | | | | | | P&J [†] Total |
|------------------------------|----------------------------|-------|-------|-------|------|----|---------------------------|-----------------------|----------------------------|-------|-------|-------|------|-----|---------------------------|
| | All Ages | ≥65 | 45-64 | 25-44 | 1-24 | <1 | | | All Ages | ≥65 | 45-64 | 25-44 | 1-24 | <1 | |
| NEW ENGLAND | 562 | 421 | 86 | 38 | 11 | 6 | 46 | S. ATLANTIC | 1,014 | 676 | 186 | 90 | 35 | 27 | 81 |
| Boston, Mass. | 153 | 105 | 24 | 15 | 6 | 3 | 15 | Atlanta, Ga. | U | U | U | U | U | U | U |
| Bridgeport, Conn. | 43 | 38 | 4 | 1 | - | - | 3 | Baltimore, Md. | 118 | 72 | 30 | 10 | 5 | 1 | 11 |
| Cambridge, Mass. | 22 | 17 | 2 | 2 | - | 1 | 2 | Charlotte, N.C. | 93 | 66 | 16 | 7 | 3 | 1 | 8 |
| Fall River, Mass. | 35 | 30 | 5 | - | - | - | 1 | Jacksonville, Fla. | 141 | 99 | 24 | 13 | 5 | - | 15 |
| Hartford, Conn. | 46 | 33 | 10 | 2 | - | 1 | 1 | Miami, Fla. | 98 | 68 | 18 | 8 | 3 | 1 | 9 |
| Lowell, Mass. | 28 | 20 | 7 | 1 | - | - | 3 | Norfolk, Va. | 55 | 32 | 9 | 7 | 2 | 5 | 1 |
| Lynn, Mass. | 7 | 5 | 1 | 1 | - | - | - | Richmond, Va. | 56 | 33 | 11 | 6 | 4 | 2 | 5 |
| New Bedford, Mass. | 25 | 22 | 1 | 1 | 1 | - | 1 | Savannah, Ga. | 56 | 39 | 12 | 3 | 1 | 1 | 6 |
| New Haven, Conn. | 43 | 30 | 7 | 4 | 2 | - | 6 | St. Petersburg, Fla. | 59 | 47 | 5 | 5 | - | 2 | 5 |
| Providence, R.I. | 49 | 40 | 5 | 3 | 1 | - | 2 | Tampa, Fla. | 225 | 166 | 37 | 10 | 8 | 4 | 19 |
| Somerville, Mass. | 3 | 1 | 2 | - | - | - | - | Washington, D.C. | 88 | 41 | 24 | 9 | 4 | 10 | 2 |
| Springfield, Mass. | 39 | 28 | 8 | 2 | 1 | - | 4 | Wilmington, Del. | 25 | 13 | - | 12 | - | - | - |
| Waterbury, Conn. | 9 | 6 | - | 2 | - | 1 | 2 | E.S. CENTRAL | 928 | 631 | 180 | 64 | 25 | 27 | 76 |
| Worcester, Mass. | 60 | 46 | 10 | 4 | - | - | 6 | Birmingham, Ala. | 177 | 115 | 37 | 11 | 4 | 9 | 21 |
| MID. ATLANTIC | 2,429 | 1,697 | 496 | 162 | 35 | 38 | 110 | Chattanooga, Tenn. | 84 | 61 | 19 | 3 | 1 | - | 4 |
| Albany, N.Y. | 47 | 30 | 11 | 3 | 1 | 2 | 3 | Knoxville, Tenn. | 78 | 59 | 13 | 4 | - | 2 | 6 |
| Allentown, Pa. | U | U | U | U | U | U | U | Lexington, Ky. | 85 | 49 | 24 | 5 | 5 | 2 | 6 |
| Buffalo, N.Y. | 102 | 81 | 11 | 5 | 3 | 1 | 9 | Memphis, Tenn. | 215 | 136 | 43 | 23 | 6 | 7 | 20 |
| Camden, N.J. | 42 | 27 | 6 | 5 | 2 | 2 | 2 | Mobile, Ala. | 79 | 62 | 8 | 5 | 2 | 2 | 1 |
| Elizabeth, N.J. | 10 | 9 | 1 | - | - | - | 2 | Montgomery, Ala. | 64 | 46 | 12 | 5 | - | 1 | 6 |
| Erie, Pa. | 38 | 30 | 5 | 3 | - | - | 2 | Nashville, Tenn. | 146 | 103 | 24 | 8 | 7 | 4 | 12 |
| Jersey City, N.J. | 50 | 26 | 14 | 7 | - | 3 | - | W.S. CENTRAL | 1,066 | 719 | 215 | 88 | 22 | 22 | 60 |
| New York City, N.Y. | 1,236 | 862 | 260 | 84 | 10 | 20 | 22 | Austin, Tex. | 79 | 56 | 12 | 8 | 1 | 2 | 4 |
| Newark, N.J. | 66 | 26 | 20 | 15 | 3 | 2 | 3 | Baton Rouge, La. | 30 | 21 | 6 | 1 | 1 | 1 | 1 |
| Paterson, N.J. | 32 | 18 | 10 | 3 | - | 1 | 1 | Corpus Christi, Tex. | 50 | 38 | 9 | 3 | - | - | 5 |
| Philadelphia, Pa. | 419 | 296 | 88 | 22 | 9 | 4 | 29 | Dallas, Tex. | 200 | 131 | 42 | 22 | 3 | 2 | 5 |
| Pittsburgh, Pa. [§] | 59 | 36 | 14 | 5 | 4 | - | 2 | El Paso, Tex. | 95 | 65 | 17 | 6 | 3 | 4 | 3 |
| Reading, Pa. | 28 | 21 | 7 | - | - | - | 1 | Ft. Worth, Tex. | 111 | 77 | 25 | 8 | 1 | - | 11 |
| Rochester, N.Y. | 121 | 95 | 19 | 4 | 1 | 2 | 16 | Houston, Tex. | U | U | U | U | U | U | U |
| Schenectady, N.Y. | 26 | 20 | 5 | 1 | - | - | 2 | Little Rock, Ark. | 65 | 47 | 13 | 2 | - | 3 | 4 |
| Scranton, Pa. | 43 | 37 | 3 | 2 | 1 | - | 2 | New Orleans, La. | 108 | 55 | 29 | 14 | 6 | 4 | 4 |
| Syracuse, N.Y. | 54 | 39 | 11 | 3 | 1 | - | 7 | San Antonio, Tex. | 170 | 116 | 32 | 17 | 2 | 3 | 11 |
| Trenton, N.J. | 30 | 20 | 9 | - | - | 1 | 5 | Shreveport, La. | 33 | 23 | 6 | 2 | 2 | - | 1 |
| Utica, N.Y. | 26 | 24 | 2 | - | - | - | 2 | Tulsa, Okla. | 125 | 90 | 24 | 5 | 3 | 3 | 11 |
| Yonkers, N.Y. | U | U | U | U | U | U | U | MOUNTAIN | 1,082 | 742 | 210 | 80 | 31 | 19 | 91 |
| E.N. CENTRAL | 2,072 | 1,388 | 421 | 148 | 60 | 55 | 162 | Albuquerque, N.M. | 120 | 86 | 19 | 9 | 5 | 1 | 12 |
| Akron, Ohio | 62 | 50 | 8 | 2 | 1 | 1 | 6 | Boise, Idaho | 40 | 28 | 7 | 3 | 1 | 1 | 3 |
| Canton, Ohio | 34 | 28 | 5 | 1 | - | - | 4 | Colo. Springs, Colo. | 59 | 44 | 8 | 3 | 1 | 3 | 4 |
| Chicago, Ill. | 413 | 249 | 96 | 33 | 19 | 16 | 38 | Denver, Colo. | 106 | 63 | 21 | 10 | 5 | 7 | 15 |
| Cincinnati, Ohio | 58 | 40 | 11 | 5 | 1 | 1 | 5 | Las Vegas, Nev. | 260 | 183 | 55 | 18 | 4 | - | 14 |
| Cleveland, Ohio | 131 | 75 | 30 | 14 | 6 | 6 | 8 | Ogden, Utah | 18 | 15 | 2 | 1 | - | - | - |
| Columbus, Ohio | 188 | 133 | 35 | 15 | 3 | 2 | 18 | Phoenix, Ariz. | 187 | 114 | 45 | 14 | 9 | 5 | 11 |
| Dayton, Ohio | 153 | 107 | 31 | 10 | 3 | 2 | 14 | Pueblo, Colo. | 35 | 27 | 4 | 3 | 1 | - | 3 |
| Detroit, Mich. | 213 | 121 | 49 | 29 | 8 | 6 | 20 | Salt Lake City, Utah | 121 | 83 | 24 | 9 | 3 | 2 | 18 |
| Evansville, Ind. | 42 | 31 | 9 | 2 | - | - | 1 | Tucson, Ariz. | 136 | 99 | 25 | 10 | 2 | - | 11 |
| Fort Wayne, Ind. | 57 | 39 | 11 | 1 | 3 | 3 | - | PACIFIC | 1,598 | 1,131 | 300 | 104 | 39 | 23 | 131 |
| Gary, Ind. | 10 | 6 | 4 | - | - | - | - | Berkeley, Calif. | 20 | 14 | 4 | 1 | - | 1 | 2 |
| Grand Rapids, Mich. | 57 | 38 | 12 | 2 | 2 | 3 | 6 | Fresno, Calif. | 141 | 96 | 29 | 8 | 5 | 3 | 18 |
| Indianapolis, Ind. | 185 | 127 | 41 | 9 | 4 | 4 | 8 | Glendale, Calif. | 18 | 12 | 4 | - | 1 | 1 | 1 |
| Lansing, Mich. | 47 | 34 | 8 | 5 | - | - | 5 | Honolulu, Hawaii | 81 | 61 | 14 | 4 | 1 | 1 | 5 |
| Milwaukee, Wis. | 128 | 99 | 20 | 5 | 2 | 2 | 9 | Long Beach, Calif. | 77 | 64 | 7 | 5 | 1 | - | 15 |
| Peoria, Ill. | 40 | 28 | 6 | 3 | 2 | 1 | 3 | Los Angeles, Calif. | 337 | 223 | 65 | 29 | 13 | 7 | 13 |
| Rockford, Ill. | 49 | 32 | 12 | 3 | 2 | - | 2 | Pasadena, Calif. | 24 | 18 | 6 | - | - | - | 3 |
| South Bend, Ind. | 63 | 48 | 12 | 1 | - | 2 | 5 | Portland, Oreg. | 194 | 129 | 51 | 8 | 4 | 2 | 15 |
| Toledo, Ohio | 84 | 62 | 12 | 7 | - | 3 | 6 | Sacramento, Calif. | U | U | U | U | U | U | U |
| Youngstown, Ohio | 58 | 41 | 9 | 1 | 4 | 3 | 4 | San Diego, Calif. | 183 | 131 | 29 | 10 | 6 | 7 | 14 |
| W.N. CENTRAL | 935 | 671 | 178 | 45 | 18 | 23 | 70 | San Francisco, Calif. | U | U | U | U | U | U | U |
| Des Moines, Iowa | 100 | 76 | 16 | 6 | 1 | 1 | 7 | San Jose, Calif. | 162 | 112 | 33 | 13 | 4 | - | 13 |
| Duluth, Minn. | 55 | 42 | 8 | 3 | 1 | 1 | 1 | Santa Cruz, Calif. | 33 | 30 | 1 | 2 | - | - | 5 |
| Kansas City, Kans. | 25 | 17 | 6 | 1 | - | 1 | 3 | Seattle, Wash. | 150 | 101 | 30 | 14 | 4 | 1 | 14 |
| Kansas City, Mo. | 111 | 77 | 24 | 6 | 2 | 2 | 5 | Spokane, Wash. | 61 | 47 | 7 | 7 | - | - | 9 |
| Lincoln, Nebr. | 42 | 34 | 6 | 2 | - | - | 3 | Tacoma, Wash. | 117 | 93 | 20 | 3 | - | - | 4 |
| Minneapolis, Minn. | 222 | 172 | 38 | 3 | 6 | 3 | 28 | TOTAL | 11,686 [¶] | 8,076 | 2,272 | 819 | 276 | 240 | 827 |
| Omaha, Nebr. | 90 | 65 | 14 | 7 | 1 | 3 | 5 | | | | | | | | |
| St. Louis, Mo. | 141 | 72 | 43 | 12 | 5 | 9 | - | | | | | | | | |
| St. Paul, Minn. | 68 | 54 | 12 | 1 | 1 | - | 13 | | | | | | | | |
| Wichita, Kans. | 81 | 62 | 11 | 4 | 1 | 3 | 5 | | | | | | | | |

U: Unavailable - : no reported cases

*Mortality data in this table are voluntarily reported from 122 cities in the United States, most of which have populations of 100,000 or more. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included.

[†]Pneumonia and influenza.

[§]Because of changes in reporting methods in this Pennsylvania city, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks.

[¶]Total includes unknown ages.

Additional information and applications are available from Emory University, International Health Dept. (PIA), 1518 Clifton Rd., N.E., Room 746, Atlanta, GA 30322; telephone (404) 727-3485; fax (404) 727-4590; or email pvaleri@sph.emory.edu.

Erratum: Vol. 48, No. RR-14

In the *MMWR Recommendations and Reports*, "Neuraminidase Inhibitors for Treatment of Influenza A and B Infections," the fifth sentence in the Summary on page 1 and the first sentence in the Conclusion on page 6 should read: "Amantadine was approved for prophylaxis of influenza A(H2N2) infection in the United States in 1966 and was approved for prophylaxis and treatment of influenza A infection in 1976; rimantadine was approved for treatment and prophylaxis of influenza A infection in 1993."

**Contributors to the Production of the *MMWR* (Weekly)
Weekly Notifiable Disease Morbidity Data and 122 Cities Mortality Data**

Samuel L. Groseclose, D.V.M., M.P.H.

State Support Team

Robert Fagan
Jose Aponte
Paul Gangarosa, M.P.H.
Gerald Jones
David Nitschke
Carol A. Worsham

CDC Operations Team

Carol M. Knowles
Deborah A. Adams
Willie J. Anderson
Fredrick Browder
Patsy A. Hall
Kathryn Snavelly

The *Morbidity and Mortality Weekly Report (MMWR) Series* is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/> or from CDC's file transfer protocol server at <ftp.cdc.gov>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (888) 232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Director, Centers for Disease Control
and Prevention
Jeffrey P. Koplan, M.D., M.P.H.

Acting Deputy Director for Science
and Public Health, Centers for
Disease Control and Prevention
Lynne S. Wilcox, M.D., M.P.H.

Acting Director,
Epidemiology Program Office
Barbara R. Holloway, M.P.H.

Editor, *MMWR* Series
John W. Ward, M.D.

Managing Editor,
MMWR (weekly)
Karen L. Foster, M.A.

Writers-Editors,
MMWR (weekly)
Jill Crane
David C. Johnson
Teresa F. Rutledge
Caran R. Wilbanks
Desktop Publishing
Morie M. Higgins