

MORBIDITY AND MORTALITY

WEEKLY REPORT

# April 20, 2001 / Vol. 50 / No. 15

- 85 50 Years of the Epidemic Intelligence Service
- 285 Mortality During a Famine Ethiopia, 2000
   289 Fatal and Severe Hepatitis Associated With Bifampin and Pyrazinamida
  - Rifampin and Pyrazinamide for the Treatment of Latent Tuberculosis Infection—New York and Georgia, 2000
- 291 Cluster of Tuberculosis Cases Among Exotic Dancers and Their Close Contacts — Kansas, 1994–2000
- 293 Outbreaks of *Escherichia coli* O157:H7 Infections Among Children Associated With Farm Visits — Pennsylvania and Washington

# 50 Years of the Epidemic Intelligence Service

This issue of *MMWR* commemorates the 50th anniversary of the Epidemic Intelligence Service (EIS). In 1951, EIS was established by CDC following the start of the Korean War as an early-warning system against biologic warfare and manmade epidemics. EIS officers selected for 2-year field assignments were primarily medical doctors and other health professionals, such as sanitarians, dentists, and veterinarians, who focused on infectious disease outbreaks. EIS has expanded to include a range of public health professionals, such as postdoctoral scientists in statistics, epidemiology, microbiology, anthropology, sociology, and behavioral sciences. The scope of work also has expanded to include chronic disease, environmental health, unintentional injury, violence prevention, and workplace health and safety. Since 1951, approximately 2500 EIS officers have responded to requests for epidemiologic assistance within the United States and throughout the world. Each year, EIS officers are involved in several hundred investigations of disease and injury problems, enabling CDC and its public health partners to make recommendations to improve the public's health and safety. Additional information about EIS and its 50th anniversary is available at http://www.cdc.gov/eis.

# Mortality During a Famine — Gode District, Ethiopia, July 2000

Recurrent famine has been a major cause of mortality in the Horn of Africa (1,2). In Ethiopia, three consecutive years of drought led to widespread loss of livestock, population displacement, and malnutrition, placing an estimated 10 million persons at risk for starvation in 2000 (3). A large proportion of the population of the Gode district in Somali region was displaced in a search for food and food aid (CDC, unpublished data, 2000). From April through July 2000, nongovernmental organizations (NGOs) opened feeding centers in the Gode district. Because no vital statistics or public health surveillance system existed in the district, and no representative mortality or morbidity data were available, during July 2000, CDC, in collaboration with Save the Children U.S., the Office of Foreign Disaster Assistance of the U.S. Agency for International Development, and the United Nations Children's Fund (UNICEF), conducted a mortality survey. This report summarizes the results of this survey, which found persistently high levels of mortality, with measles representing an important cause of mortality in children aged

# U.S. DEPARTMENT OF HEALTH & HUMAN SERVICES

## Mortality During a Famine — Continued

<5 years and 5–14 years. Mass measles vaccination with vitamin A distribution is an important intervention during the acute phase of famines in sub-Saharan Africa.

During a two-stage cluster survey in Gode district, the collaborating agencies collected retrospective mortality data from December 9, 1999, through July 31, 2000. A sample size of 3832 persons was required to achieve a 95% confidence interval (CI) with 2% precision around an estimated cumulative incidence of mortality of 10%. The design effect is the factor by which the sample size calculated for a simple random sample needs to be multiplied to account for the dependence of a given variable within a cluster. Although a design effect of two generally is assumed for nutrition surveys, deaths were expected to be more clustered than malnutrition, and a design effect of four was used in this survey. After adjusting for more recent estimates by NGOs involved in food distribution and by the Ethiopian army conducting comprehensive headcounts, the 1994 census (4) was used as the basis for the sampling frame. In the first stage of the survey, 30 clusters were assigned proportionally to village population size. In the second stage, households were selected using Expanded Program on Immunization methods (5). A household was defined as a group of persons who normally lived together and shared meals. Age at death and month of death were identified. Cause of death was assigned using standard case definitions for easily recognized causes of death. Analysis was performed using Epilnfo version 6.04b (6).

A total of 595 households comprising 4032 persons was surveyed. In stable, developing countries, the crude mortality rate (CMR) is generally  $\leq$ 0.5 deaths per 10,000 persons per day and the mortality rate for children aged <5 years (<5MR) is  $\leq$ 1 per 10,000 persons per day (7). During the study period, the CMR was 3.2 (95% Cl=2.4–3.8), three times the cut-off level of one per 10,000 per day used to define an emergency (7). The CMR peaked in January 2000 at 6.3 but during July was still 2.0. During the study period, the <5MR was 6.8 (95% Cl=5.3–8.0). The <5MR was highest in December 1999 at 12.5 but during July 2000, was 5.5, above the emergency threshold for <5MR of 2–4 (7). Of the 293 deaths that occurred during the study period, 158 (54%) were in children aged <5 years, and 73 (25%) were in children aged 5–14 years. Measles and malnutrition (without an accompanying major communicable disease) each contributed to approximately one fourth of the 159 deaths among children aged <5 years; diarrhea was reported as the cause of death for approximately one third of deaths in this age group (Figure 1). Measles also contributed to 12 (17%) of 72 deaths among children aged 5–14 years.

As a result of these findings, the following emergency measures were recommended: 1) accelerating plans for a mass measles vaccination campaign and vitamin A distribution targeting children aged 9 months–5 years; 2) extending coverage of the campaign to include children aged 6 months–14 years; 3) implementing water and sanitation programs to prevent diarrheal disease; 4) continuing treatment for severely malnourished children in therapeutic feeding centers and moderately malnourished children in supplementary feeding programs; and 5) ongoing monitoring of malnutrition (Figure 2) and mortality using cross-sectional surveys in the absence of a regular mortality surveillance system.

Reported by: A Teklehaimanot, I Jabr, A Paganini, United Nations Children's Fund. Disaster Preparedness and Prevention Commission, Government of Ethiopia; F Assefa, T Degefie, M Shibeshi, K Zeynu, A Makki, P Brandrup, M Clark, Save the Children US, Gode, Ethiopia. P Morris, R Machmer, K Lattu, Office of Foreign Disaster Assistance, US Agency for International Development. International Emergency and Refugee Health Br, Div of Emergency Environmental Health Svcs, National Center for Environmental Health; and an EIS Officer, CDC. Mortality During a Famine — Continued





**Editorial Note:** The age distribution for mortality during the famine in Ethiopia is similar to other famine- and emergency-affected populations. Children, particularly those aged <5 years, usually account for most deaths in such situations ( $\mathcal{B}$ ). Malnutrition, diarrheal diseases, acute respiratory infection, malaria, and measles account for 60%–95% of reported deaths in famines and complex emergencies (7). For children aged <5 years, measles is a leading cause of mortality during these emergencies. Most famines occur in areas of rural sub-Saharan Africa, where measles vaccination coverage is rarely adequate to prevent measles outbreaks during periods of mass displacement and malnutrition. Mass measles vaccination campaigns targeting children aged 6 months-5 years are likely to be cost-effective in such situations ( $\mathcal{B}$ ) and may prevent many more deaths than more high-profile interventions (e.g., feeding centers). The large proportion of measles-related deaths among children aged 5–14 years identified in this survey highlights the importance of extending coverage to children aged >5 years when measles-related mortality is high in this age group ( $\mathcal{B}$ ).

The findings in this report are subject to at least three limitations. First, data are subject to recall bias; as a result, the study period was limited to 8 months, and the beginning of the study period was defined by a religious date known to the entire population. Second, only households present on the day of the survey were sampled, possibly resulting in an underestimation of mortality because households in which all members had died during the famine could not have been selected. Finally, because no surveillance system and no birth and death registration existed in the district, comparing verbal reports of mothers with case definitions was used to determine causes of death. Inadequate sensitivity and specificity of case definitions could have resulted in some misclassification of causes of death.

## Mortality During a Famine — Continued

FIGURE 2. Ethiopian child being weighed with a Salter scale, 2000



Guidelines for humanitarian interventions prioritize interventions to be implemented: rapid assessment, measles vaccination with vitamin A distribution, water and sanitation programs, and food aid (10). In refugee camps, mass measles vaccination campaigns accompanied by vitamin A distribution and water and sanitation programs have become standard practice. This report underscores the importance of these programs in the acute phase of famines in sub-Saharan Africa. Such programs are more difficult to implement in widely dispersed famine-affected populations than in refugee or internally displaced camps, particularly in remote areas, such as the Somali region of Ethiopia, that have no cold chain and poor health infrastructure. Even though food aid and feeding centers also are a priority during famine, attracting a large concentration of susceptible persons to feeding centers may increase transmission of infectious diseases such as measles and diarrhea. Public health programs targeting major causes of mortality should be integrated with feeding programs during famine from the outset of the humanitarian response.

# References

- 1. Lindtjorn B. Famine in southern Ethiopia, 1985–86: population structure, nutritional state and incidence of death. BMJ 1990;301:1123–7.
- 2. Murray M, Murray A, Murray N, Murray M. Somali food shelters in the Ogaden famine and their impact on health. Lancet 1976;332:1283–5.
- 3. United Nations Children's Fund. Situation report. Addis Ababa, Ethiopia: United Nations Children's Fund, November 2000:1–10.
- Government of Ethiopia. The 1994 population and housing census of Ethiopia, results for Somali region. Addis Ababa, Ethiopia: Office of Population and Housing Census Commission, Central Statistical Authority, 1999:1–265.
- 5. Henderson R, Sundaresan T. Cluster sampling to assess immunization coverage: a review of experience with a simplified sampling method. Bull World Health Organ 1982;60:253–60.
- Dean AG, Dean JA, Burton AH, Dicker RC. EpiInfo version 6: a word processing, database and statistics program for epidemiology on microcomputers. Stone Mountain, Georgia: USD Incorporated, 1990.
- 7. CDC. Famine-affected, refugee, and displaced populations: recommendations for public health issues. MMWR 1992;41(no. RR-13).
- 8. Toole MJ, Waldman RJ. An analysis of mortality trends among refugee populations in Somalia, Sudan, and Thailand. Bull World Health Organ 1988;66:237–47.
- 9. Toole MJ, Steketee RW, Waldman RJ, Nieburg P. Measles prevention and control in emergency settings. Bull World Health Organ 1989;67:381–8.
- 10. SPHERE Project. Humanitarian charter and minimum standards in disaster response. Geneva, Switzerland: Steering Committee for Humanitarian Response, 1998.

# Fatal and Severe Hepatitis Associated With Rifampin and Pyrazinamide for the Treatment of Latent Tuberculosis Infection — New York and Georgia, 2000

One of the recommended treatments for latent tuberculosis infection (LTBI) is a 9-month regimen of isoniazid (INH); a 2-month regimen of rifampin (RIF) and pyrazinamide (PZA) is an alternative in some instances. In September 2000, a man in New York died of hepatitis after 5 weeks of RIF-PZA, and in December, a woman in Georgia was admitted to a hospital because of hepatitis after 7 weeks of this regimen. This report summarizes the findings of the investigations of these incidents, which underscore the need for clinical monitoring for adverse effects in all patients receiving treatment for LTBI.

# Case 1

A 53-year-old incarcerated man received 600 mg (6.7 mg/Kg) RIF and 1750 mg (19 mg/Kg) PZA daily after screening revealed a tuberculin skin test (TST) with 20 mm induration and no radiologic or clinical findings of active tuberculosis (TB). His risk factors for TB included previous work as a medical orderly, homelessness, and multiple incarcerations. He had a history of hypertensive heart disease and alcoholism without evidence of chronic liver disease. He was not known to inject drugs.

RIF-PZA was standard treatment for LTBI at the jail. Baseline and 1-month serum aminotransferase and bilirubin levels were measured routinely. The patient's baseline aminotransferase levels were slightly higher than the upper-normal limits. He was instructed to stop taking RIF-PZA if he developed symptoms suggestive of hepatitis. He also received 325 mg enteric-coated aspirin daily, 90 mg extended-release nifedipine, and 50 mg hydrochlorothiazide. Nurses supervised the administration of all medication to assure compliance.

Blood specimens tested on day 33 of treatment revealed alanine aminotransferase (ALT) 1734 U/L (normal range: 0–41 U/L), aspartate aminotransferase (AST) 1449 U/L (normal range: 0–38 U/L), and total bilirubin 4.2 mg/dL (normal range: 0–1.0 mg/dL). Blood cell counts showed leukocytosis. On day 35, RIF-PZA was discontinued when the test results were received. On the same day, a correctional officer urged the patient to visit the infirmary because of poor appetite and lassitude that had developed over several days; he declined. Five days after the cessation of RIF-PZA, the patient was evaluated in the infirmary for jaundice and altered mental status and was admitted to a hospital. Serum total bilirubin peaked at 17.8 mg/dL and blood ammonia at 378  $\mu$ mol/L (normal range: 17–47  $\mu$ mol/L). He died 3 days after admission.

On postmortem histology, the liver had bridging necrosis, lymphocytic infiltration, focal cholestasis, increased fibrosis, and micronodular cirrhosis. Results were negative for serum anti-A IgM, antibody to hepatitis B core antigen (anti-HBc), antibody to hepatitis B surface antigen (anti-HBs), and antibody to hepatitis C virus (anti-HCV). Antinuclear antibody (ANA) was undetectable. Hepatitis B and C were undetectable by polymerase chain reaction assays. The reported cause of death was liver necrosis and failure as a result of hepatitis following LTBI treatment.

# Case 2

A 59-year-old woman received 600 mg (7.2 mg/Kg) RIF and 2000 mg (24 mg/Kg) PZA daily after testing revealed a TST with 27 mm induration and no findings for active TB. She chose this regimen because of suspected exposure to drug-resistant TB and concern about liver injury from INH. In addition to RIF-PZA, she received beclomethasone

# Treatment of Latent Tuberculosis Infection - Continued

dipropionate nasal spray, budesonide inhalation powder, and albuterol inhalation aerosol for nasal allergies and asthma. She had no history of liver disease, rarely drank alcohol, and did not inject drugs. She was vaccinated against hepatitis A but not B. She had a history of anaphylactic reactions to penicillin and an estrogen sulfates blend. Baseline ALT and AST, bilirubin levels, and blood cell counts were normal. She was instructed to contact her health-care provider about adverse effects during treatment. On day 2 of treatment, she reported queasiness. On day 17, her blood tests were repeated: serum aminotransferase and bilirubin levels were normal, and her eosinophil count, which had been 157 cells/ $\mu$ L, was 510 cells/ $\mu$ L (normal range: 50–550 cells/ $\mu$ L).

She subsequently experienced malaise, anorexia, and feverishness, and she occasionally took one bismuth subsalicylate chewable tablet. On the 49th and last day of treatment, she returned to her health-care provider and was admitted to a hospital because of jaundice and altered mental status. AST was 986 U/L (normal range: 7–40 U/L), ALT 1735 U/L (normal range: 17–63 U/L), and total bilirubin 11.4 mg/dL (normal range: 0.1–1.1 mg/dL). The bilirubin peaked at 27.5 mg/dL after 14 days. Peak eosinophil count was 2580 cells/ $\mu$ L. No ova or protozoa were detected by stool examinations. Serum ANA was 1:640 (speckled pattern). Antibody (not IgM) to hepatitis A virus was detected. Test results were negative for hepatitis B surface antigen (HBsAg), anti-HBs, and anti-HCV. After receiving 40 mg prednisone daily, the symptoms and laboratory abnormalities slowly abated, and she was released after 25 days in the hospital.

Reported by: M DeMartino, MD, Nassau County Office of the Medical Examiner; J Maniscalco, A Greenberg, MD, Nassau County Dept of Health, Mineola; J Grabau, PhD, M Oxtoby, MD, E Foster, MS, Bur of Tuberculosis Control, P Smith, MD, State Epidemiologist, New York State Dept of Health. P Kozarsky, MD, C Pox, MD, Emory Univ School of Medicine, Atlanta, Georgia. National Institute of Diabetes and Diseases of the Digestive System and Kidneys, National Institutes of Health, Bethesda, Maryland. Occupational Health Clinic, Office of Health and Safety, Office of the Director, Hepatitis Br, Div of Viral and Rickettsial Diseases, National Center for Infectious Diseases; Div of Tuberculosis Elimination, National Center for HIV, STD, and TB Prevention, CDC.

**Editorial Note**: Case 1 is the first report to CDC of fatal hepatitis associated with the RIF-PZA regimen for LTBI, although sporadic cases of liver injury have been attributed to PZA used in treatment regimens for TB disease (1). Both cases illustrate that the usually well-tolerated regimens for LTBI occasionally can result in severe adverse effects and that clinical monitoring is crucial during treatment. In these cases, biochemical monitoring did not help to avoid severe liver injury and does not substitute for clinical monitoring (2). Idiosyncratic liver injury can be caused by hypersensitivity, as suspected for case 2, or by toxic drug metabolites. Other cases have implicated various medicines and alcohol as potential co-factors for INH liver injury (3,4). A similar association has not been assessed for RIF and PZA because of small case numbers.

Patients with LTBI and risk factors for active TB should be offered treatment (1,5). Health-care providers should instruct and frequently remind patients about the initial symptoms of hepatitis (e.g., fatigue, nausea, abdominal pain, and anorexia) and the importance of stopping medication if symptoms develop (2). In this report, both patients continued taking their medicines while symptoms were developing, a phenomenon also reported for INH-associated hepatitis (4).

CDC's Division of Tuberculosis Elimination is interested in receiving reports of severe hepatitis in patients being treated for LTBI. To report possible cases, telephone (404) 639-8125.

Treatment of Latent Tuberculosis Infection — Continued

#### References

- 1. Fox W, Mitchison DA. Short-course chemotherapy for tuberculosis. American Review of Respiratory Disease 1975;111:325–53.
- American Thoracic Society. Targeted tuberculin testing and treatment of latent tuberculosis infection. Am J Respir Crit Care Med 2000;161:S221–S247.
- 3. Millard PS, Wilcosky TC, Reade-Christopher SJ, Weber DJ. Isoniazid-related fatal hepatitis. West J Med 1996;164:486–91.
- 4. CDC. Severe isoniazid-associated hepatitis—New York, 1991–1993. MMWR 1993;42:545–7.
- CDC. Tuberculosis elimination revisited: obstacles, opportunities, and a renewed commitment—Advisory Council for the Elimination of Tuberculosis (ACET). MMWR 1999;48 (no. RR-9).

# Cluster of Tuberculosis Cases Among Exotic Dancers and Their Close Contacts — Kansas, 1994–2000

During January 2001, the Wichita-Sedgwick County Department of Community Health (WSCDCH), the Kansas Department of Health and Environment (KDHE), and CDC investigated a cluster of tuberculosis (TB) cases that occurred from 1994 to 2000 among women with a history of working as dancers in adult entertainment clubs (i.e., exotic dancers) and persons who were close contacts of exotic dancers. This report describes the results of the investigation and illustrates the need for early identification of TB clusters through ongoing surveillance and resources for health departments to respond rapidly to TB outbreaks.

As of April 2001, the TB control staff of WSCDCH and KDHE had identified 18 TB cases in this cluster that had been diagnosed from 1994 to 2000 (Figure 1). Of these, 14 (78%) were culture confirmed; all *Mycobacterium tuberculosis* isolates were susceptible to first-line anti-TB drugs. Eight patients were women (seven exotic dancers), seven were men, and three were children. Of the 15 adult patients, 14 were aged <45 years at the time of diagnosis. All dancers had cavitary pulmonary disease, an indication of increased infectiousness. All adult patients were voluntarily tested for human immunodeficiency virus infection and one was seropositive. Twelve (80%) of the 15 adult patients reported using cocaine, crack cocaine, or amphetamines, and 10 (67%) had been incarcerated at some time during 1994–2000. All 18 patients were started on directly observed therapy (DOT), and 17 completed treatment.

Evidence linking these cases included common occupation or known exposure to exotic dancers. Of the 11 nondancer patients, six were exposed to dancers outside of the clubs exclusively. Although dancer patients identified six clubs in which they worked during their potential infectious periods, no single club could be confirmed as the site of transmission to all other dancers. Shared drug-related activities may have linked the adult patients; however, no specific location of drug use was identified (1). Of the nine *M. tuberculosis* isolates tested, all had matching *IS6110* fingerprints, including isolates from six dancers (2).

Contact investigations of the nine infectious TB patients identified 344 contacts. Of 302 contacts with a tuberculin skin test (TST) placed and read, 76 (25%) were TST positive. Among 243 contacts eligible for 10-to-12 week postexposure TST, 32 (13%) had follow-up TST placed and read. Of these, 14 (44%) had TST conversion indicating recent *M. tuberculosis* infection. Among 72 contacts eligible for latent TB infection (LTBI) therapy, 54 (75%) initiated therapy. Of the 54 contacts who should have completed therapy by January 2001, six (11%) had documented completion.

*Tuberculosis Cases Among Exotic Dancers — Continued* 





\* n=18.

Reported by: C Magruder, MD, R Woodruff, G Minns, MD, V Barnett, P Baker, E Brady, T Julian, MPH, Wichita-Sedgwick County Dept of Community Health; G Pezzino, MD, M Reece, A Alejos, Kansas Dept of Health and Environment. MD Cave, PhD, National Tuberculosis Genotype Surveillance Network, Little Rock, Arkansas. R Rothenberg, MD, Emory Univ School of Medicine, Atlanta, Georgia. Div of Applied Public Health Training; Statistics and Epidemiology Br, Div of Prevention Research and Analytic Methods, Epidemiology Program Office; Field Services Br and Surveillance and Epidemiology Br, Div of Tuberculosis Elimination, National Center for HIV, STD, and TB prevention; and EIS officers, CDC.

**Editorial Note:** The findings in this report indicate the need for local health departments to have sufficient resources for ongoing surveillance for TB and capacity to rapidly respond during a time of increased demand. The cluster in Kansas occurred over a 7-year period and encompassed 18 patients.

The WSCDCH TB control staff consists of a full-time TB control nurse, a part-time physician consultant, and a full-time assistant. The nurse is primarily responsible for TB case management including DOT. In addition, in collaboration with the WSCDCH Health Surveillance Unit, the nurse is responsible for contact investigations and screening high-risk persons for TB with TST. Health departments in low incidence states such as Kansas (2.9\* per 100,000 population during 2000) may have limited resources to respond to outbreaks while maintaining the essential components of TB control, thus hampering efforts to eliminate TB (*3*).

<sup>\*</sup>Provisional 2000 data.

#### Tuberculosis Cases Among Exotic Dancers — Continued

Outbreaks of TB among persons who use illegal drugs and/or have been incarcerated can be difficult to investigate. Illegal drug users often belong to complex social networks, and members of these networks may be reluctant or unable to provide the names of their contacts to public health officials (4). Special techniques for exploring chains of transmission among members of complex social networks have been developed (5,6).

In this cluster investigation, follow-up rates of 10-to-12 week postexposure TST and completion rates of LTBI therapy were low. New approaches beyond traditional methods of TB contact investigations are necessary to follow-up contacts discovered through social network analysis. These approaches must assure that all contacts are assessed for LTBI and that those with LTBI complete therapy. This may require DOT for LTBI in an outbreak to prevent further *M. tuberculosis* transmission. The findings in this report underscore that all states, including those with very low TB incidence, should maintain TB control capacity and have outbreak response plans that include methods to augment this capacity during unexpected increases in *M. tuberculosis* transmission (7).

#### References

- 1. CDC. Crack cocaine use among persons with tuberculosis—Contra Costa County, California, 1987–1990. MMWR 1991;40:485–9.
- Van Emden J, Cave M, Crawford J, et al. Strain identification of *Mycobacterium tuberculo*sis by DNA fingerprinting: recommendations for a standardized methodology. J Clin Microbiol 1993;31:406–9.
- CDC. Essential components of a tuberculosis prevention and control program (ACET) MMWR 1995;44(no. RR-11).
- CDC. HIV-related tuberculosis in a transgender network—Baltimore, Maryland, and New York City area, 1998–2000. MMWR 2000;49:317–20.
- Rothenberg R, Narramore J. The relevance of social network concepts to sexually transmitted diseases control. Sex Transm Dis 1996;23:24–9.
- 6. Klovdahl A, Graviss E, Yaganehdoost A. Networks and tuberculosis: an undetected community outbreak involving public places. Soc Sci Med 2001;52:681–94.
- 7. Institute of Medicine. Ending neglect: the elimination of tuberculosis in the United States. Washington, DC: National Academy Press, 2000.

# Outbreaks of *Escherichia coli* O157:H7 Infections Among Children Associated With Farm Visits — Pennsylvania and Washington, 2000

During the spring and fall of 2000, outbreaks of *Escherichia coli* O157:H7 infections among school children in Pennsylvania and Washington resulted in 56 illnesses and 19 hospitalizations. Illness was associated with school and family visits to farms where children came into direct contact with farm animals. This report summarizes the findings of investigations of these outbreaks (Figure 1) and includes strategies to reduce the transmission of enteric pathogens from farm animals to children.

# Pennsylvania

During September–November 2000, the Montgomery County Health Department (MCHD) identified 51 persons who had diarrhea within 10 days of visiting a dairy farm (farm A) in Montgomery County. Fifteen (29%) persons had either *E. coli* O157 isolated from stool specimens or hemolytic-uremic syndrome (HUS); patients ranged in age from 1–52 years (median: 4 years), 26 (51%) were male, and dates of illness onset ranged from September 4 to November 8. Symptoms reported by the 51 patients included

Escherichia coli 0157:H7 Infections - Continued

# FIGURE 1. CDC investigator examines a calf at farm A — Pennsylvania, 2000



bloody diarrhea (37%), fever (45%), and vomiting (45%); 16 (31%) patients were hospitalized and eight (16%) developed HUS. *E. coli* O157 isolates were indistinguishable by pulsed-field gel electrophoresis (PFGE) and produced both Shiga toxins 1 and 2.

To identify risk factors, CDC, the Pennsylvania Department of Health, and MCHD conducted a case-control study among farm visitors during November 12–19. A confirmed case was defined as diarrhea in a person within 10 days of visiting farm A on or after September 1, with either *E. coli* O157 isolated from stool or HUS. A probable case was defined as diarrhea in a person within 10 days of visiting farm A on or after September 1. Controls also had visited farm A after September 1 but did not develop diarrhea within 10 days of the visit. Two controls per case were sought by sequential digit dialing and frequency matched by age group (i.e., <1 year, 1–4 years, 5–8 years, 9–12 years, 13–20 years, and ≥21 years). Fifty-one case-patients, or a parent or guardian for young children, and 92 controls were interviewed in the case-control study.

Case-patients were more likely than controls to have had contact with cattle (summary odds ratio [OR]=10.9; 95% confidence interval [CI]=1.7–70.7), an important farm animal reservoir for *E. coli* O157. Activities that promoted hand-mouth contact, such as nailbiting (summary OR=2.5; 95% CI=1.1–5.7) and purchasing food from an outdoor concession (summary OR=2.5; 95% CI=1.1–5.7), were more common among patients. Handwashing before eating was protective (summary OR=0.2; 95% CI=0.1–0.7). All 216 cattle on farm A were sampled by rectal swab, and 28 (13%) yielded *E. coli* O157 with a PFGE pattern indistinguishable from that isolated from patients. The same strain also was isolated from a railing surface. *E. coli* O157 was not isolated from 43 of the other animal species on the farm.

Among the 75,600 persons who visited farm A during the outbreak, most were preschool-aged or school-aged, groups at risk for serious *E. coli* O157 infection (1). No separate area was designated for interaction between visitors and farm animals.

### Escherichia coli 0157:H7 Infections — Continued

Visitors could touch cattle, calves, sheep, goats, llamas, chickens, and a pig and could eat and drink while interacting with animals. Handwashing facilities lacked soap and disposable towels, were out of children's reach, were few in number, and were unsupervised.

A total of 19,698 telephone calls were made to identify controls; 3497 household members were available. Household members were asked whether they had visited farm A since September 1 and whether they developed diarrhea within 10 days of the visit; 134 visited the farm during the outbreak, and 22 (16.4%) reported onset of diarrhea within 10 days of the visit. The expected rate of diarrhea from any cause in the general population during a 10-day period is approximately 7% (FoodNet Population Survey, unpublished data, 1998–1999). Because approximately 75,600 persons visited the farm during the outbreak, an estimated 7000 (9.4%) may have developed diarrhea associated with their visit. No further illness was reported after public access to animals was discontinued at farm A.

# Washington

During May–June 2000, five persons with culture-confirmed *E. coli* O157 infection were reported to the Snohomish Health District (SHD). Isolates from these persons were indistinguishable by PFGE. Dates of illness onset were May 21–31, and patients ranged in age from 2 to 14 years (median: 7 years); three were male. All five patients reported abdominal cramping and diarrhea, and four reported bloody diarrhea. Three patients, aged 2–6 years, were hospitalized, and one developed HUS. Four patients attending three elementary schools had visited a dairy farm (farm B) on May 18 or 24. The fifth patient had not visited farm B but had developed diarrhea after a sibling became ill following a farm B visit. Approximately 300 persons visited farm B during the outbreak, primarily preschool- and kindergarten-aged children accompanied by adults.

On May 31 and June 1, an investigation of farm B by SHD and the Washington Department of Health revealed that children were allowed to handle young poultry, rabbits, and goats. Goats, chickens, and a calf were kept in pens and could be touched through a fence. Children brought their own lunches and ate approximately 50 feet from the penned animals. Five animal stool samples collected from the farm were tested for *E. coli* O157; all were negative.

Farm B recommended that visitors bring antibacterial wipes to wash their hands; the farm also provided a communal rinse basin. No signs were posted instructing visitors to wash their hands after touching the animals. No further illness was reported after prevention measures were instituted, including distribution of instructional material and installation of handwashing stations with soap and running water.

Reported by: R Gage, MSPH, A Crielly, MS, M Baysinger, E Chernak, MD, G Herbert, A Johnson-Entsuah, MPH, Montgomery County Health Dept, Norristown; G Fraser, C Rinehardt, M Solomon, G Withers, MS, R Berman, MS, Bur of Laboratories, Lionville; M Moll, MD, J Rankin, DVM, Pennsylvania Dept of Health. J Carroll, M Ettinger, MS, S Henderson, M Mismas, D Patel, T Reed, E Smith, J Wozniak, MS, D Toney, PhD, J Pearson, DrPH, Virginia Div of Consolidated Laboratory Svcs, Richmond. J Hofmann, MD, Snohomish Health District, Everett; J Grendon, DVM, J Kobayashi, MD, Washington Dept of Health. Animal and Plant Health Inspection Svc, US Dept of Agriculture. Foodborne and Diarrheal Diseases Br, Div of Bacterial and Mycotic Diseases, National Center for Infectious Diseases; and an EIS Officer, CDC.

**Editorial Note**: The outbreaks described in this report were the first reported in the United States to be associated with direct transmission of *E. coli* O157 from farm animals to humans. An estimated 73,500 cases of illness, 2000 hospitalizations, and 60 deaths occur in the United States each year as the result of *E. coli* O157 infection (*2*); many *E. coli* O157 illnesses are associated with ingesting contaminated food or drink. However,

# Escherichia coli 0157:H7 Infections — Continued

during 1996 and 1997, visiting a farm with cows was identified as an important risk factor for *E. coli* O157 infection; 8% of persons aged  $\geq$ 6 years with *E. coli* O157 infection reported visiting a farm with cows during the preceding 7 days compared with 1% of controls (*3*).

Two random-digit–dial telephone surveys of 9000 persons were conducted during 1996–1997 and 1998–1999; 2% reported having visited a petting zoo during the preceding 5–7 days (4,5). In 1999 in Ontario, Canada, an *E. coli* O157 outbreak among visitors to a petting zoo resulted in 159 illnesses (6). In the United Kingdom, farm visit-related outbreaks of *E. coli* O157 infections have been reported among children (7). Such outbreak have led to the development of guidelines to prevent *E. coli*-related illnesses in these countries (6,8).

Of the 44 state and territorial public health departments responding to a national CDC survey in June 2000, none had laws to control exposure of humans to enteric pathogens at venues where the public has access to farm animals, and no federal laws exist that address this public health issue. Following these U.S. farm-associated outbreaks, CDC, in collaboration with the Zoonoses Working Group, National Association of State Public Health Veterinarians, U.S. Department of Agriculture, Animal and Plant Health Inspection Services, and other groups, drafted measures to reduce the risk for farm animal-human transmission of enteric infections (see box).

Before July 1, 2001, comments about prevention measures can be mailed to Strategies, Foodborne and Diarrheal Diseases Branch, Division of Bacterial and Mycotic Diseases, CDC, 1600 Clifton Road, MS A-38, Atlanta, Georgia 30333, or e-mailed to zcn0@cdc.gov.

# References

- 1. Boyce TG, Swerdlow DL, Griffin PM. *Escherichia coli* O157:H7 and the hemolytic-uremic syndrome. N Engl J Med 1995;333:364–8.
- Mead PS, Slutsker L, Dietz V, et al. Food-related illness and death in the United States. Emerg Infect Dis 1999;5:607–25.
- Kassenborg H, Hedberg C, Evans M. Case-control study of sporadic *Escherichia coli* O157:H7 infections in 5 FoodNet sites (California, Connecticut, Georgia, Minnesota, and Oregon). Presented at the 1st International Conference on Emerging Infectious Diseases, Atlanta, Georgia, 1998.
- CDC. Foodborne diseases active surveillance network (FoodNet): population survey atlas of exposures: 1998–1999. Atlanta, Georgia: US Department of Health and Human Services, CDC, 1999.
- CDC. Foodborne diseases active surveillance network (FoodNet): population survey atlas of exposures: 1996–1997. Atlanta, Georgia: US Department of Health and Human Services, CDC, 1997.
- Warshawsky B, Henry B, Gutmanis I, et al. An *Escherichia coli* O157:H7 outbreak associated with an animal exhibit: Middlesex-London Health Unit Investigation and Recommendations, 1999. Available at http://www.healthunit.com/reportsresearch.htm. Accessed April 2001.
- Milne LM, Plom A, Strudley I, et al. *Escherichia coli* O157 incident associated with a farm open to members of the public. Communicable Disease & Public Health 1999;2:22–6.
- Health and Safety Executive. Avoiding ill health at open farms: advice to farmers. Sudbury, England: HSE Books, 2000; revised ed.,vol. 23. Available at http://www.hsebooks.co.uk/ index2.html. Accessed April 2001.

Escherichia coli 0157:H7 Infections - Continued

# Reducing the Risk for Transmission of Enteric Pathogens at Petting Zoos, Open Farms, Animal Exhibits, and Other Venues Where the Public Has Contact With Farm Animals

- Information should be provided. Persons providing public access to farm animals should inform visitors about the risk for transmission of enteric pathogens from farm animals to humans, and strategies for prevention of such transmission. This should include public information and training of facility staff. Visitors should be made aware that certain farm animals pose greater risk for transmitting enteric infections to humans than others. Such animals include calves and other young ruminant animals, young poultry, and ill animals. When possible, information should be provided before the visit.
- Venues should be designed to minimize risk. Farm animal contact is not appropriate at food service establishments and infant care settings, and special care should be taken with school-aged children. At venues where farm animal contact is desired, layout should provide a separate area where humans and animals interact and an area where animals are not allowed. Food and beverages should be prepared, served, and consumed only in animalfree areas. Animal petting should occur only in the interaction area to facilitate close supervision and coaching of visitors. Clear separation methods such as double barriers should be present to prevent contact with animals and their environment other than in the interaction area.
- Handwashing facilities should be adequate. Handwashing stations should be available to both the animal-free area and the interaction area. Running water, soap, and disposable towels should be available so that visitors can wash their hands immediately after contact with the animals. Handwashing facilities should be accessible, sufficient for the maximum anticipated attendance, and configured for use by children and adults. Children aged <5 years should wash their hands with adult supervision. Staff training and posted signs should emphasize the need to wash hands after touching animals or their environment, before eating, and on leaving the interaction area. Communal basins do not constitute adequate handwashing facilities. Where running water is not available, hand sanitizers may be better than using nothing. However, CDC makes no recommendations about the use of hand sanitizers because of a lack of independently verified studies of efficacy in this setting.</li>
- Hand-mouth activities (e.g., eating and drinking, smoking, and carrying toys and pacifiers) should not be permitted in interaction areas.
- Persons at high risk for serious infections should observe heightened precaution. Farm animals should be handled by everyone as if the animals are colonized with human enteric pathogens. However, children aged <5 years, the elderly, pregnant women, and immunocompromised persons (e.g., those with HIV/AIDS) are at higher risk for serious infections. Such persons should weigh the risks for contact with farm animals. If allowed to have contact, children aged <5 years should be supervised closely by adults, with precautions strictly enforced.</li>
- Raw milk should not be served.

298



# FIGURE I. Selected notifiable disease reports, United States, comparison of provisional 4-week totals ending April 14, 2001, with historical data

- \* No rubella cases were reported for the current 4-week period yielding a ratio for week 15 of zero (0).
- <sup>†</sup> Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

		Cum. 2001		Cum. 2001
Anthrax Brucellosis*		- 16	Poliomyelitis, paralytic Psittacosis*	- 3
Cholera		-	Q fever*	4
Cyclosporiasis	*	31	Rabies, human	-
Diphtheria		-	Rocky Mountain spotted fever (RMSF)	29
Ehrlichiosis:	human granulocytic (HGE)*	11	Rubella, congenital syndrome	-
	human monocytic (HME)*	3	Streptococcal disease, invasive, group A	1,076
Encephalitis:	California serogroup viral*	-	Streptococcal toxic-shock syndrome*	17
	eastern equine*	-	Syphilis, congenital <sup>§</sup>	17
	St. Louis*	-	Tetanus	3
	western equine*	-	Toxic-shock syndrome	42
Hansen diseas	se (leprosy)*	16	Trichinosis	5
Hantavirus pu	Imonary syndrome*	3	Tularemia*	8
Hemolytic ure	mic syndrome, postdiarrheal*	15	Typhoid fever	46
HIV infection,	pediatric*†	37	Yellow fever	-
Plague		-		

# TABLE I. Summary of provisional cases of selected notifiable diseases, United States, cumulative, week ending April 14, 2001 (15th Week)

-: No reported cases. \*Not notifiable in all states.

<sup>1</sup>Updated monthly from reports to the Division of HIV/AIDS Prevention — Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention (NCHSTP). Last update February 27, 2001. <sup>§</sup>Updated from reports to the Division of STD Prevention, NCHSTP.

							Escherichia coli 0157:H7*			
	AI	DS Ourse	Chlan	nydia†	Cryptosp	oridiosis	NET		PH	LIS
<b>Reporting Area</b>	2001 <sup>§</sup>	2000	2001	2000	2001	2000	2001	2000	2001	2000
UNITED STATES	5,820	9,320	171,244	195,384	368	401	274	409	179	329
NEW ENGLAND Maine N.H. Vt. Mass. R.I. Conn.	200 3 12 9 118 24 34	653 11 9 - 439 20 174	5,749 316 327 169 2,532 800 1,605	6,723 376 319 161 2,846 686 2,335	13 - 5 4 2 2	26 3 1 8 7 2 5	31 4 5 1 16 5	39 3 4 1 18 - 13	26 3 - 14 2 4	38 3 4 2 14 - 15
MID. ATLANTIC Upstate N.Y. N.Y. City N.J. Pa.	1,180 29 740 241 170	2,343 102 1,428 481 332	14,795 N 7,432 1,287 6,076	18,692 N 7,947 3,857 6,888	44 21 21 1 1	84 21 58 1 4	25 20 5 N	52 47 4 1 N	15 10 1 4	51 38 1 6 6
E.N. CENTRAL Ohio Ind. III. Mich. Wis.	463 77 45 226 97 18	850 112 75 535 99 29	22,284 498 4,212 6,355 8,475 2,744	33,152 8,875 3,906 9,429 6,215 4,727	110 31 14 30 35	86 14 4 13 11 44	57 19 9 13 7	79 15 9 27 12 16	26 10 2 7 - 7	26 8 10 - 4 4
W.N. CENTRAL Minn. Iowa Mo. N. Dak. S. Dak. S. Dak. Nebr. Kans.	110 29 15 38 1 - 9 18	164 36 13 72 - 2 9 32	9,102 1,683 999 3,195 240 539 778 1,668	10,906 2,360 1,160 3,722 274 527 1,061 1,802	15 - 7 4 - 1 3 -	24 5 6 1 3 2 3	27 8 3 11 - 1 - 4	61 10 12 25 2 1 7 4	21 11 2 5 - 1 2	65 29 7 15 4 2 5 3
S. ATLANTIC Del. Md. D.C. Va. W. Va. N.C. S.C. Ga. Fla.	1,673 37 131 166 137 12 101 171 187 731	2,492 44 267 186 158 13 101 174 293 1,256	37,275 875 3,786 966 5,147 647 5,908 3,605 7,485 8,856	36,515 860 3,484 871 4,336 618 5,788 4,250 6,909 9,399	82 1 19 3 6 - 11 - 25 17	55 1 5 - 2 - 6 - 32 9	34 - 7 1 16 1 2 6	33 - - 6 2 8 2 3 7	14 - - 5 - 5 - 2 2	26 - U 7 1 2 1 7 7
E.S. CENTRAL Ky. Tenn. Ala. Miss.	360 51 132 95 82	343 56 133 100 54	13,708 2,352 4,238 3,818 3,300	14,847 2,295 4,221 4,916 3,415	11 1 2 4 4	13 - 2 7 4	11 1 6 4	22 8 7 1 6	8 2 5 - 1	19 7 10 2
W.S. CENTRAL Ark. La. Okla. Tex.	629 45 188 36 360	757 30 124 31 572	28,649 2,383 4,938 2,884 18,444	29,006 1,526 5,321 2,474 19,685	7 2 3 2	18 1 2 1 14	18 - - 6 12	24 4 - 4 16	21 - 8 5 8	38 3 8 3 24
MOUNTAIN Mont. Idaho Wyo. Colo. N. Mex. Ariz. Utah Nev.	241 5 - 40 15 93 23 60	289 5 4 1 62 40 92 30 55	8,637 471 529 219 805 1,520 3,607 279 1,207	11,288 348 556 217 3,254 1,365 3,712 746 1,090	37 3 5 12 8 1 8 -	27 1 3 2 8 1 3 7 2	32 3 5 15 1 5 2 1	34 8 4 3 12 5 1 1	17 - - 9 - 4 3 1	18 - 2 6 - 6 1 2
PACIFIC Wash. Oreg. Calif. Alaska Hawaii	964 117 38 798 2 9	1,429 141 35 1,215 5 33	31,045 4,001 268 25,202 688 886	34,255 3,850 1,851 27,037 705 812	49 N 2 47 -	68 U 2 66 -	39 9 5 25 -	65 10 9 40 1 5	31 8 5 16 - 2	48 21 9 13 1 4
Guam P.R. V.I. Amer. Samoa C.N.M.I.	5 158 1 - -	13 184 11 -	1,451 U U U	- U U U U	- U U U	- U U U	N - U U U	N 1 U U U		

 TABLE II. Provisional cases of selected notifiable diseases, United States, weeks ending April 14, 2001, and April 15, 2000 (15th Week)

N: Not rotifiable. U: Unavailable. -: No reported cases. C.N.M.L: Commonwealth of Northern Mariana Islands. \* Individual cases can be reported through both the National Electronic Telecommunications System for Surveillance (NETSS) and the Public Health Laboratory Information System (PHLIS). \* Chlamydia refers to genital infections caused by *C. trachomatis.* Totals reported to the Division of STD Prevention, NCHSTP. \* Updated monthly from reports to the Division of HIV/AIDS Prevention — Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention. Last update February 27, 2001.

	Gono	rrhea	Hepatit Non-A, I	is C; Von-B	Legionellosis		Listeriosis	Lyme Disease	
Reporting Area	Cum. 2001	Cum. 2000	Cum. 2001	Cum. 2000	Cum. 2001	Cum. 2000	Cum. 2001	Cum. 2001	Cum. 2000
UNITED STATES	81,486	99,721	521	946	190	204	82	567	1,184
NEW ENGLAND Maine N.H. Vt. Mass. R.I. Conn.	1,572 37 35 26 762 201 511	1,842 22 26 15 728 169 882	5 - 3 2 -	6 - 3 3 -	8 - 2 3 2 - 1	16 2 - 9 - 3	10 - - 6 - 4	141 - 42 1 19 - 79	171 - 18 - 63 - 90
MID. ATLANTIC Upstate N.Y. N.Y. City N.J. Pa.	9,030 2,017 3,210 797 3,006	10,387 1,686 3,371 2,202 3,128	23 14 - 9	198 13 176 9	17 11 3 2 1	45 16 5 2 22	9 3 2 1 3	272 216 - 56	806 322 29 98 357
E.N. CENTRAL Ohio Ind. III. Mich. Wis.	11,797 341 1,764 3,922 4,886 884	19,974 4,927 1,721 6,668 4,606 2,052	60 4 - 3 53 -	76 - - 9 67 -	58 32 6 - 14 6	59 26 9 6 10 8	8 1 - 5 1	13 13 - - U	28 4 - 1 - 23
W.N. CENTRAL Minn. Iowa Mo. N. Dak. S. Dak. S. Dak. Nebr. Kans.	3,866 504 307 2,008 9 58 248 732	4,600 893 272 2,260 14 75 351 735	141 - 136 - 2 3	139 - 132 - 2 5	15 1 4 7 - 2 1	11 1 3 5 - 1 - 1	2 - - - - - 1	20 13 4 - 1 1	18 6 - 6 - 1 5
S. ATLANTIC Del. Md. D.C. Va. W. Va. N.C. S.C. Ga. Fla.	22,754 453 2,310 913 2,633 139 4,801 2,592 3,933 4,980	27,488 464 2,399 636 2,893 167 5,136 4,936 4,334 6,523	31 - - - 3 7 2 - 9	23 2 4 - 1 2 8 - 6	28 - 7 1 4 N 2 - 2 12	38 3 11 3 N 5 2 2 12	16 - 2 - 2 1 - - 4 7	99 - 83 6 1 2 - - 1	128 16 93 - 8 4 4 - - 3
E.S. CENTRAL Ky. Tenn. Ala. Miss.	8,922 932 2,796 3,153 2,041	10,398 945 3,202 3,654 2,597	65 3 18 1 43	140 15 26 4 95	17 6 7 2 2	6 4 1 1	7 1 3 3	2 2 - -	1 - 1 -
W.S. CENTRAL Ark. La. Okla. Tex.	13,925 1,511 3,353 1,366 7,695	14,965 767 3,790 1,093 9,315	142 3 56 2 81	281 3 171 107	3 - 2 1 -	5 - 2 1 2	2 1 - 1	- - - -	9 - 2 - 7
MOUNTAIN Mont. Idaho Wyo. Colo. N. Mex. Ariz. Utah Nev.	2,712 26 26 958 272 956 26 432	3,064 8 20 990 296 1,249 89 386	23 - 1 3 8 7 1 - 3	30 1 12 4 9 - 3	15 - 1 4 1 6 1 2	13 - 1 - 6 1 2 3 -	7 - - 1 2 1 1 2	1 - - - - 1	
PACIFIC Wash. Oreg. Calif. Alaska Hawaii	6,908 898 45 5,715 87 163	7,003 706 241 5,864 81 111	31 9 1 21 -	53 6 12 35 -	29 5 N 24 -	11 5 N 6 -	21 2 19 -	19 2 1 16 N	23 2 21 N
Guam P.R. V.I. Amer. Samoa C.N.M.I.	- 364 U U U	- 140 U U U	- - U U U	- 1 U U U	2 U U U	- - U U U	- - - -	N U U U	N U U U

# TABLE II. (Cont'd) Provisional cases of selected notifiable diseases, United States,<br/>weeks ending April 14, 2001, and April 15, 2000 (15th Week)

N: Not notifiable. U: Unavailable. -: No reported cases.

					Salmonellosis*						
	Ma	aria	Rabies	, Animal	NE	rss	PHLIS				
Reporting Area	Cum. 2001	Cum. 2000	Cum. 2001	Cum. 2000	Cum. 2001	Cum. 2000	Cum. 2001	Cum. 2000			
UNITED STATES	209	258	1,240	1,590	5,496	6,922	4,564	6,454			
NEW ENGLAND Maine N.H. Vt. Mass. R.I. Conn.	17 1 5 - 10	10 1 - 1 6 - 2	137 20 5 26 38 16 32	182 47 3 11 55 12 54	443 38 37 20 263 23 62	438 32 25 34 258 9 80	424 17 33 22 232 35 85	469 22 29 40 254 31 93			
MID. ATLANTIC Upstate N.Y. N.Y. City N.J. Pa.	35 9 17 6 3	49 15 24 5 5	196 161 1 33 1	254 188 3 39 24	496 190 195 69 42	1,029 221 308 281 219	661 122 266 143 130	1,184 307 312 222 343			
E.N. CENTRAL Ohio Ind. III. Mich. Wis.	26 5 8 - 13	35 3 19 9 2	7 - 1 - 6 -	14 2 - 6 6	857 332 65 219 160 81	1,060 234 106 374 164 182	703 274 65 179 119 66	580 203 123 1 181 72			
W.N. CENTRAL Minn. Iowa Mo. N. Dak. S. Dak. Nebr. Kans.	7 1 2 - 1 2	14 - 1 - 3 6	90 15 16 5 14 13 - 27	130 23 17 4 24 35 - 27	365 71 60 125 1 24 31 53	329 37 43 108 4 18 53 66	375 136 53 127 9 12 - 38	442 130 49 135 18 25 38 47			
S. ATLANTIC Del. Md. D.C. Va. W. Va. N.C. S.C. Ga. Fla.	60 1 25 4 12 - 1 2 3 12	59 24 16 7 1 1	555 10 88 - 104 40 154 27 68 64	550 10 120 - - - - - - - - - - - - - - - - - - -	1,398 24 157 18 182 10 258 149 215 385	1,171 17 184 135 31 200 100 191 313	929 23 159 U 161 18 160 174 188 46	984 26 181 U 138 25 147 84 298 85			
E.S. CENTRAL Ky. Tenn. Ala. Miss.	8 2 3 3	10 2 1 6 1	35 5 25 5	56 9 33 14	345 61 93 138 53	352 72 85 114 81	174 33 98 31 12	273 48 121 88 16			
W.S. CENTRAL Ark. La. Okla. Tex.	3 - 1 1 1	3 - 3 -	80 - 21 59	291 - 20 271	400 58 60 31 251	657 63 70 62 462	382 29 125 30 198	425 36 82 56 251			
MOUNTAIN Mont. Idaho Wyo. Colo. N. Mex. Ariz. Utah Nev.	18 2 1 9 1 1 2 2	15 1 - 8 - 2 2 2	46 7 - 10 - 1 28 - -	48 10 - 22 - 3 13 -	432 16 19 13 127 56 127 48 26	613 21 37 9 186 56 162 95 47	351 4 13 109 47 108 47 23	572 35 11 177 50 158 92 49			
PACIFIC Wash. Oreg. Calif. Alaska Hawaii	35 1 32 1	63 4 17 40 - 2	94 - 66 28 -	65 - 57 8 -	760 94 15 642 9	1,273 83 88 1,032 16 54	565 144 61 284 76	1,525 156 109 1,194 18 48			
Guam P.R. V.I. Amer. Samoa C.N.M.I.	- U U U	2 U U U	42 U U U	- 18 U U U	75 U U U	92 U U U	U U U U U				

# TABLE II. (Cont'd) Provisional cases of selected notifiable diseases, United States, weeks ending April 14, 2001, and April 15, 2000 (15th Week)

N: Not notifiable. U: Unavailable. -: No reported cases. \* Individual cases can be reported through both the National Electronic Telecommunications System for Surveillance (NETSS) and the Public Health Laboratory Information System (PHLIS).

	NET	Shige	losis*		Sy (Drimory)	philis Secondary)	Tuberculosis		
-	Cum.	Cum.	Cum.	Cum.	Cum.	Cum.	Cum.	Cum.	
Reporting Area	2001	2000	2001	2000	2001	2000	2001	2000	
	2,820	4,433	1,466	2,830	1,400	1,871	2,387	3,192	
Maine	41	90 2	50	/0	-	- 23	90 5	90	
N.H. Vt.	1	1 1	1	2	-	-	6 1	2	
Mass. B.L	29 2	66 7	31 5	50 8	7	19 1	53 9	58 7	
Conn.	7	13	11	16	3	3	22	27	
MID. ATLANTIC	282 121	655 203	223	484 134	84 4	91 4	512 67	536 51	
N.Y. City	93	351	124	222	61	41	255	309	
Pa.	40 28	39	49 44	66	10	29	66	36	
E.N. CENTRAL	452	737	235	267	207	398	267	335	
Ind.	74	45 96	14	39 20	24 47	134	47 22	27	
III. Mich.	124 92	279 233	84 57	2 197	36 92	134 88	137 39	190 30	
Wis.	25	84	7	9	8	20	22	23	
W.N. CENTRAL Minn.	333 105	263 43	277 148	224 70	15 7	30 3	110 54	132 51	
lowa Mo.	69 80	44 136	61 52	54 81	- 6	8 15	9 30	11 52	
N. Dak.	9 19	1	1	1	-	-	-	-	
Nebr.	23	22	-	11	-	2	13	3	
	29 459	16	14	/ 156	569	2	-	12	
Del.	409	3	2	3	2	2	525	547	
D.C.	36 16	- 30	U	U	72 12	98 19	49 13	- 64	
Va. W. Va.	34 4	16 2	19 6	25 2	48	39 1	47 8	60 10	
N.C.	102	33	51 17	16	143 79	159	77 19	89 18	
Ga.	58 177	60	25	60	68 145	104	121	142	
F.S. CENTRAL	248	198	4 71	149	145	281	169	227	
Ky.	88	39	25	22	12	27	15	24	
Ala.	67	9	17	7	27	40	78	77	
WISS.	400	51 683	0 252	3 226	30 203	37 266	33 175	42 519	
Ark.	156	60	65	20	15	24	38	39	
Okla.	6	8	2	8	23	57	28	23	
ιέχ. Μοι ινιταίνι	219	535 276	132	160	123 54	118	78	432	
Mont.	- 5	1	-	- 17	-	-	- 4	4	
Wyo.	-	1	-	1	-	-	-	-	
N. Mex.	41 38	47 27	27	22 17	4	6	26 5	15	
Ariz. Utah	79 10	107 16	46 11	44 21	37 6	40	23 5	42 8	
Nev.	14	53	8	38	3	2	15	37	
PACIFIC Wash.	418 51	1,029 192	100 62	1,088 228	97 19	128 16	457 54	675 57	
Oreg. Calif	4 361	85 734	27	49 800	- 75	3 109	393	22 546	
Alaska	2	6 12	- 11	3	- 3	-	10	20 30	
Guam	-	-	U	U	-	-	-	-	
P.R.	7	14	Ŭ	Ŭ	96	52	38	21	
Amer. Samoa	Ŭ	Ŭ	Ŭ	Ŭ	Ŭ	Ŭ	Ŭ	Ŭ	
0.11.101.1.	0	U	0	0	0	0	U	U	

TABLE II. (Cont'd) Provis	sional cases of sel	ected notifiable of	liseases, Un	ited States,
weeks endin	g April 14, 2001, a	nd April 15, 2000	(15th Week)	

N: Not notifiable. U: Unavailable. -: No reported cases. \*Individual cases can be reported through both the National Electronic Telecommunications System for Surveillance (NETSS) and the Public Health Laboratory Information System (PHLIS).

	H. influ	uenzae,	Hepatitis (Viral), By Type				Measles (Rubeola)					
	Inva	asive	Α	-	В	-	Indige	nous	Impo	rted*	Tota	
Reporting Area	Cum. 2001 <sup>†</sup>	Cum. 2000	Cum. 2001	Cum. 2000	Cum. 2001	Cum. 2000	2001	Cum. 2001	2001	Cum. 2001	Cum. 2001	Cum. 2000
UNITED STATES	389	413	2,367	3,650	1,609	1,689	2	15	-	15	30	19
NEW ENGLAND	14	33	98	94	16	29	-	3	-	1	4	-
Maine N.H.	1	1 6	1 5	5 8	2 6	1 6	-	-	-	-	-	-
Vt.	-	3	2	3	1	3	-	1	-	-	1	-
R.I.	-	-	30 5	39 5	6	6	-	2 -	-	-	-	-
Conn.	-	4	50	34	-	12	-	-	-	-	-	-
MID. ATLANTIC	47 17	64 26	207	246	212	289	1	2	-	4	6	8
N.Y. City	18	21	85	132	115	161	-	-	-	-	-	8
N.J. Pa.	11 1	13	46 15	- 44	44 15	14 85	-	1	-	-	1	-
E.N. CENTRAL	49	64	264	507	192	166	-	-		7	7	3
Ohio	26 12	20	80	111	35	32	-	-	-	2	2	2
III.	4	25	59	217	14	2	-	-	-	3	3	-
Mich. Wis.	3	3 11	103	153 13	138	120 1	-	-	-	-	-	1
W.N. CENTRAL	18	12	135	320	58	87	-	4	-	-	4	-
Minn.	8	7	8	36	5	6 12	-	1	-	-	1	-
Mo.	8	4	41	199	37	56	-	3	-	-	3	-
N. Dak. S. Dak.	-	1	- 1	-	- 1	-	-	-	-	-	-	-
Nebr.	1	-	18 55	12	5	9	-	-	-	-	-	-
	-	-	50	41 275	5 270	4 290	-	-	-	-	-	-
Del.	-	-	510	6		205	-	-	-	-	-	-
Md. D.C.	38	28	70 14	47	44 3	47	-	2	-	1	3	-
Va.	9	20	42	46	39	39	-	-	-	-	-	-
N.C.	20	8	34	65	80	81	-	-	-	-	-	-
S.C. Ga.	2 31	5 26	17 167	12 48	1 94	2 45	-	- 1	-	-	- 1	-
Fla.	41	13	164	118	103	69	-	-	-	-	-	-
E.S. CENTRAL	25	18	80	164	99 11	120	-	-	-	-	-	-
Tenn.	12	6	38	57	39	54	-	-	-	-	-	-
Ala. Miss.	11 1	3	30 4	23 68	28 21	9 38	-	-	-	-	-	-
W.S. CENTRAL	9	23	326	692	212	192	-	1	-	-	1	-
Ark.	-	7	17	53	26	25	-	-	-	-	-	-
Okla.	7	16	53	106	25	40 23	-	-	-	-	-	-
Tex.	-	-	236	505	147	96	-	1	-	-	1	-
MOUNTAIN Mont.	73	48	239 4	255 1	160 1	132 3	-	-	-	1	1	2
Idaho	1	2	26	11	4	4	-	-	-	1	1	-
Colo.	- 15	- 11	28		34	28	-	-	-	-	-	-
N. Mex.	10 38	11 19	7 118	30 121	43 59	43 40	-	-	-	-	-	-
Utah	2	3	22	17	6	3	-	-	-	-	-	-
Nev.	7	2	33	17	13	11	-	-	-	-	-	2
PACIFIC Wash.	9 1	48 2	508 22	997 62	290 27	385 17	1	2	-	1	3	6 3
Oreg.	2	16	10	74	5	33	1	1	ū	-	1	-
Alaska	1	1	10	4	4	2	-	-	-	-	-	-
Hawaii	-	13	-	7	-	5	-	-	-	-	-	-
Guam P.R.	-	- 2	- 28	- 106	- 15	- 74	U U	-	U U	-	-	-
V.I.	U	Ū	Ŭ	Ŭ	Ŭ	Ŭ	Ŭ	U	Ŭ	U	U	U
C.N.M.L	U	U	U	U	U	U	U	U	U	U	U	U

# TABLE III. Provisional cases of selected notifiable diseases preventable<br/>by vaccination, United States, weeks ending April 14, 2001,<br/>and April 15, 2000 (15th Week)

N: Not notifiable. U: Unavailable. - : No reported cases. \*For imported measles, cases include only those resulting from importation from other countries. † Of 74 cases among children aged <5 years, serotype was reported for 35, and of those, 7 were type b.

Reporting Area         Cum. 2001         Cum. 2000         Cum. 2001         Cum. 2001	Rubella			
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	Cum. 2000			
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	22			
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	5 - 1 - 3 -			
N.J.21172Pa.1212-842Pa.1212-842Pa.102140-4178162210-1Ohio3824-167113131Ind.171759Ill.18386-1512Wis.9161-1740Wis.9161-1740Nox. CENTRAL5649246-4039Iowa15123-47N.Dak.2121Nebr.2312Nebr.2212-25-1032Nebr.2212-25-1032Nebr.23111<	1 5 2			
E.N. CENTRAL102140-4178162210-1Ohio3824-167113131Ind.171759III.1838-3411218-1Mich.20456-1512Wis.9161-1740Mon.6315N.O.21241-237No.21241-237Nok.211Nok.211Nok.211Nok.21112Nok.211-11Nok.211-1Nok.212-25-1032Nok.2212-	- -			
n.1030112101Wis.9161-1740W.N. CENTRAL5649246-4039Iowa15123-47-No.21241-237-N. Dak.211Nebr.231-12-Nebr.231-106S. ATLANTIC1641171516160104-1Del1Md.2212-25-1032NC.392211N.C.3922229-1Ga.2222229Fla.453229Ga.2222Md. </td <td>- -</td>	- -			
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	-			
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	1 - -			
Nebr.231-12Kans.82241-106S. ATLANTIC1641171516160104-1Del1Md.2212-25-1032D.C11Va.1819123-810N.C.39222-2328S.C.147-15-814Ga.22221-710E.S. CENTRAL60542-2935Ky.1011623Ala.24151-48	-			
S. ALLANTIC       164       117       1       5       16       1       60       104       -       1         Del.       -       -       -       -       -       -       -       1       -       -       1         Md.       22       12       -       2       5       -       10       32       -       -         D.C.       -       -       -       -       -       1       1       -       -       -         Va.       18       19       1       2       3       -       8       10       -       -       -         N.C.       39       22       -       -       2       -       23       28       -       -         S.C.       14       7       -       1       5       -       8       14       -       -         Ga.       22       22       -       -       -       2       9       -       1         Fla.       45       32       -       -       1       -       7       10       -       -         Fla.       10       11       -       -	1			
D.C.       1 <th1< th=""> <th1< th=""></th1<></th1<>	3 - -			
Inc.Image: Constraint of the second systemImage: Constraint of the second systemImage: Constraint of the second systemImage: Constraint of the second systemS.C.147-15-29-1Ga.222229-1Fla.45321-710E.S. CENTRAL60542-2935Ky.1011623Tenn.2223163Ala.24151-48Miss.451-31	-			
E.S. CENTRAL       60       54       -       -       2       -       29       35       -       -         Ky.       10       11       -       -       -       6       23       -       -         Tenn.       22       23       -       -       -       16       3       -       -         Ala.       24       15       -       -       1       -       4       8       -       -         Miss.       4       5       -       -       1       -       3       1       -       -	2 - 1			
Miss. 4 5 1 - 3 1	1 1 - -			
W.S. CENTRAL 123 91 - 5 14 - 21 29 Ark 9 5 - 1 1 - 2 5 -	- 3			
La. 41 26 - 2 3 - 1 3 Okla. 14 16 1 Tex. 59 44 - 2 10 - 17 21	- - 3			
MOUNTAIN         47         49         1         5         7         7         625         243         -         -           Mont.         -         1         -         -         1         2         5         1         -         -           Idaho         3         6         -         -         2         157         35         -         -	- -			
Wyo.     -     -     -     1     -     -     -     -       Colo.     18     12     1     2     1     2     134     151     -       N. Mex.     8     7     -     2     1     -     40     37     -	-			
Ariz.     9     10     -     -     -     1 $2/9$ 11     -     -       Utah     5     5     -     -     2     -     9     5     -     -       Nev.     4     2     -     -     2     -     1     3     -     -	-			
PACIFIC         148         178         -         14         63         2         85         256         -         -           Wash.         30         16         -         -         2         2         29         60         -         -           Oreg.         3         22         N         N         N         -         -         25         -         -           Calif.         114         135         U         13         55         U         56         158         U         -           Alaska         1         1         -         1         1         -         -         4         -         -           Hawaii         -         4         -         -         5         -         9         -         -	4 3 - 1 -			
Guam       -       -       U       -       -       U       -       -       U       -         P.R.       1       4       U       -       -       U       -       U       -       U       -       U       -       U       -       U       -       U       -       U       -       U       -       U       -       U       -       U       -       U       -       U       -       U       -       U       U       -       U       U       -       U	- - U U U			

# TABLE III. (Cont'd) Provisional cases of selected notifiable diseases preventable by vaccination, United States, weeks ending April 14, 2001, and April 15, 2000 (15th Week)

N: Not notifiable.

-: No reported cases.

		All Cau	ses, By	Age (Ye	ears)		P&I⁺		All Causes, By Age (Years)						P&I⁺
Reporting Area	All Ages	≥65	45-64	25-44	1-24	<1	Total	Reporting Area	All Ages	≥65	45-64	25-44	1-24	<1	Total
NEW ENGLAND Boston, Mass. Bridgeport, Conn. Cambridge, Mass. Fall River, Mass. Hartford, Conn. Lowell, Mass. Lynn, Mass. New Bedford, Mas New Haven, Conn Providence, R.I. Somerville, Mass. Springfield. Mass	577 152 25 16 33 28 25 15 5 5 28 25 5 37	405 84 20 14 25 20 20 20 12 13 21 88 5 26	113 36 4 2 7 6 5 2 3 3 15 - 9	27 10 - 1 1 - 1 4 4 - 2	23 17 - - - 3 -	9 5 - 1 - 2 -	86 16 5 - 6 5 7 3 4 11 - 7	S. ATLANTIC Atlanta, Ga. Baltimore, Md. Charlotte, N.C. Jacksonville, Fla Miami, Fla. Norfolk, Va. Richmond, Va. Savannah, Ga. St. Petersburg, F Tampa, Fla. Washington, D.C Wilmington, D.C	1,150 125 220 . 148 65 48 54 54 54 54 54 . 00 179 C. 107 L. 28	740 72 141 55 87 42 34 31 35 48 126 41 28	249 37 53 11 42 16 8 14 5 3 29 31	93 11 20 8 10 4 2 5 3 16 14	29 2 3 1 6 1 3 3 - 5 3 2 -	36 3 1 2 1 1 5 17	88 2 31 6 11 6 5 8 4 3 8 4 3
Waterbury, Conn. Worcester, Mass. MID. ATLANTIC Albany, N.Y. Allentown, Pa. Buffalo, N.Y. Camden, N.J. Elizabeth, N.J. Erie, Pa.§	33 71 2,248 50 17 82 23 6 42	26 51 1,563 36 14 57 15 3 33 33	7 14 475 9 2 16 3 - 7	- 3 126 3 1 7 2 3 1 2	- 2 41 1 - 1 2 -	- 1 38 1 - 1 1 - 1	5 17 116 4 2 1 3	E.S. CENTRAL Birmingham, Ali Chattanooga, Te Knoxville, Tenn. Lexington, Ky. Memphis, Tenn. Mobile, Ala. Montgomery, A Nashville, Tenn.	869 a. 212 ann. 48 94 59 . 176 69 Ia. 75 136	576 138 39 62 42 112 55 53 75	174 34 7 21 6 42 10 17 37	75 26 7 7 12 4 4 14	19 4 - 2 - 5 - 8	22 7 1 2 4 5 - 1 2	89 32 3 4 7 19 5 7 12
New York City, N.J. Paterson, N.J. Philadelphia, Pa. Pittsburgh, Pa.§ Reading, Pa. Rochester, N.Y. Schenectady, N.Y. Scranton, Pa.§ Syracuse, N.Y. Trenton, N.J. Utica, N.Y. Yonkers, N.Y.	7. 1,140 53 25 407 33 26 130 18 29 81 28 12 23 U	804 255 17 254 103 16 22 63 7 18 U	247 18 4 104 6 7 20 2 4 10 5 U	57 32 31 3 - 3 - 3 1 4 - U	20 - 9 3 - 2 - 1 - 1 - U	8 6 1 9 - 1 2 - 6 1 - U	55 1 2 12 7 12 2 2 10 1 2 U	W.S. CENTRAL Austin, Tex. Baton Rouge, La Corpus Christi, 1 Dallas, Tex. El Paso, Tex. Ft. Worth, Tex. Houston, Tex. Little Rock, Ark. New Orleans, La San Antonio, Te Shreveport, La. Tulsa, Okla.	1,516 98 57 195 57 138 323 73 . 58 x. 270 50 122	1,001 72 38 68 111 38 90 199 42 31 189 34 89	328 16 8 50 13 34 76 20 16 53 9 25	117 8 2 17 4 9 32 6 7 17 5 4	47 1 11 2 3 12 3 4 9 - 1	23 1 - 1 6 - 2 4 2 - 2 2 3	112 7 1 20 7 6 28 6 - 13 6 14
E.N. CENTRAL Akron, Ohio Canton, Ohio Chicago, III. Cincinnati, Ohio Columbus, Ohio Dayton, Ohio Detroit, Mich. Evansville, Ind. Fort Wayne, Ind. Gary, Ind. Grand Rapids, Mid Indianapolis, Ind. Lansing, Mich. Milwaukee, Wis. Peoria III.	1,699 66 33 0 69 113 288 128 128 128 128 31 79 20 ch. 46 208 48 95 202	1,209 522 U 47 755 190 96 102 199 63 14 41 153 37 70 34	339 7 7 U 19 27 69 23 50 5 10 6 5 39 5 19 3	94 2 1 2 9 20 6 16 5 4 - 8 2 4 3	32 1 2 U - 2 3 - 8 - 2 - 4 4 1 2	25 1 U 1 6 3 6 2 - 4 - 4 - 1	112 5 3 0 6 4 17 7 12 5 8 - - 15 4 13 4	MOUNTAIN Albuquerque, N Boise, Idaho Colo. Springs, C Denver, Colo. Las Vegas, Nev. Ogden, Utah Phoenix, Ariz. Pueblo, Colo. Salt Lake City, U Tucson, Ariz. PACIFIC Berkeley, Calif. Fresno, Calif. Glendale, Calif. Honolulu, Hawa	1,047 .M. 110 6 olo. 72 113 207 22 165 32 tah 118 162 1,252 21 146 U 146 U	757 93 29 57 63 148 19 106 29 87 126 924 13 112 U 50	187 14 12 9 26 45 3 29 3 19 27 207 6 25 U 9	65 3 4 3 15 12 - 18 - 4 6 73 1 5 U 1 1	27 - - 3 7 2 - 8 - 5 2 26 - 4 U 3	11 - - - - - - - - - - - - - - - - - -	91 11 4 1 8 8 2 12 3 17 5 95 1 9 U 3 1
Rockford, III. South Bend, Ind. Toledo, Ohio Youngstown, Ohi W.N. CENTRAL Des Moines, Iowa Duluth, Minn. Kansas City, Kans. Kansas City, Mo. Lincoln, Nebr. Minneapolis, Min Omaha, Nebr. St. Louis, Mo. St. Paul, Minn. Wichita, Kans.	-26 27 94 0 65 770 64 22 22 22 111 48 115 75 105 81 88	548 21 68 53 538 45 19 14 70 35 131 54 53 68 49	14 4 18 9 145 11 2 6 21 30 12 26 10 17	2 1 8 1 51 4 1 9 3 8 5 17 2 2	- 1 - 2 18 1 - 1 6 - 2 1 7 -	1 - - 18 3 - 1 5 - 4 2 2 1 -	<sup>1</sup> 3141 7192203210345 210345	Long Beach, Cal Los Angeles, Cal Pasadena, Calif. Portland, Oreg. San Caramento, Cal San Diego, Calif San Francisco, C San Jose, Calif. Santa Cruz, Calif. Seattle, Wash. Spokane, Wash. Tacoma, Wash. TOTAL	rr. 60 lif. U 32 U if. 191 . 162 . 162 . 248 f. 48 f. 48 f. 48 121 54 106 11,128 <sup>1</sup>	39 U 25 112 U 186 35 85 44 84 84 7,713	14 U 3 U 29 31 U 35 5 9 12 2,217	5 U 2 U 14 10 U 16 6 4 1 8 721	U 1 0 4 6 U 5 2 1 - - 262	2 U 1 U 5 3 U 6 - 2 - 2 02	10 4 12 12 22 6 9 7 4 860

# TABLE IV. Deaths in 122 U.S. cities,\* week ending April 14, 2001 (15th Week)

U: Unavailable. -: No reported cases. \*Mortality data in this table are voluntarily reported from 122 cities in the United States, most of which have populations of ≥100,000. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included. \*Pneumonia and influenza.

<sup>®</sup>Because of changes in reporting methods in this Pennsylvania city, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks. <sup>®</sup>Total includes unknown ages.

# Contributors to the Production of the MMWR (Weekly)

## Weekly Notifiable Disease Morbidity Data and 122 Cities Mortality Data

Samuel L. Groseclose, D.V.M., M.P.H.

State Support Team Robert Fagan Jose Aponte Gerald Jones David Nitschke Scott Noldy Carol A. Worsham

#### CDC Operations Team Carol M. Knowles Deborah A. Adams Willie J. Anderson Patsy A. Hall Mechele Hester Felicia J. Perry Pearl Sharp

# Informatics

T. Demetri Vacalis, Ph.D.

Michele D. Renshaw

Erica R. Shaver

The Morbidity and Mortality Weekly Report (MMWR) Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to *listserv@listserv.cdc.gov*. The body content should read SUBscribe mmwr-toc. Electronic copy also is available from CDC's World-Wide Web server at http://www.cdc.gov/mmwr or from CDC's file transfer protocol server at http://ftp.cdc.gov/pub/Publications/mmwr. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (888) 232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Director, Centers for Disease Control and Prevention Jeffrey P. Koplan, M.D., M.P.H.	Director, Epidemiology Program Office Stephen B. Thacker, M.D., M.Sc.	Writers-Editors, <i>MMWR</i> (Weekly) Jill Crane David C. Johnson					
Deputy Director for Science and Public Health, Centers for Disease Control and Prevention David W. Fleming, M.D.	Editor, <i>MMWR</i> Series John W. Ward, M.D. Acting Managing Editor, <i>MMWR</i> (Weekly) Teresa F. Rutledge	Desktop Publishing Lynda G. Cupell Morie M. Higgins					
☆U.S. Government Printing Office: 2001-633-173/48224Region IV							