

MMWR™

MORBIDITY AND MORTALITY WEEKLY REPORT

- 617 West Nile Virus Activity — Eastern United States, 2001
- 620 Global Progress Toward Laboratory Containment of Wild Polioviruses, June 2001
- 623 Heat-Related Deaths — Los Angeles County, California, 1999–2000, and United States, 1979–1998

West Nile Virus Activity — Eastern United States, 2001

In 2000, ArboNET, an enhanced human and animal surveillance system designed to monitor the geographic spread of West Nile virus (WNV) in the United States and to identify areas at increased risk for human infections with WNV, detected WNV activity in the District of Columbia and 12 states (1). This system, first implemented in the District of Columbia and 20 states along the Atlantic and Gulf coasts, was later expanded throughout the continental United States. This report summarizes ArboNET data from January 1 through July 25, 2001, which documents epizootic WNV activity in the southeast and indicates the need for widespread implementation of WNV prevention activities.

The first human infection in 2001 was identified in a 73-year-old man from Madison County, Florida, with illness onset on approximately July 15. He remains hospitalized with encephalitis. Equine surveillance identified three horses with neurologic disease attributed to WNV infection in Jefferson County, Florida, with illness onsets beginning on June 24. Avian mortality surveillance identified 142 WNV-infected birds from the District of Columbia (one bird) and 34 counties in nine states (Connecticut [four], Florida [21], Georgia [two], Maryland [51], Massachusetts [six], New Jersey [37], New York [16], Rhode Island [three], and Virginia [one]) (Figure 1). Crows accounted for 126 (89%) of the reported birds. In New York City, one live hatch-year house sparrow had antibody to WNV. One sentinel chicken from Duval County, Florida, seroconverted to WNV in a serum specimen drawn on July 10.

WNV also was detected in 38 mosquito pools collected in 10 counties in four states, representing at least nine species, including a pool of six *Culex salinarius* collected in Baltimore, Maryland, on July 11, a mixed pool of *Cx. pipiens*/*Cx. restuans* collected in Queens, New York, on July 3, a pool of *Ochlerotatus canadensis* collected on July 5 and a pool of *Cx. pipiens* collected on July 16 in Fairfield County, Connecticut, and 34 pools collected in seven New Jersey counties as early as May 31. The New Jersey mosquito pools included *Cx. pipiens* (17 pools), *Cx. restuans* (nine), *Culiseta melanura* (three), unidentified *Aedes*/*Oc.* species (two), *Ae. vexans* (one), *Oc. canadensis* (one), and *Oc. triseriatus* (one).

Reported by: ArboNET surveillance group in local and state health depts. National Wildlife Health Center, US Geologic Survey, Madison, Wisconsin. National Veterinary Svcs Laboratories, Veterinary Svcs, Animal and Plant Health Inspection Svc, US Dept of Agriculture, Ames, Iowa. Walter Reed Army Institute of Research, District of Columbia. US Air Force, Frederick, Maryland. Arbovirus Diseases Br, Div of Vector-Borne Infectious Diseases, National Center for Infectious Diseases, CDC.

*West Nile Virus Activity — Continued***FIGURE 1. Location of human and animal infection with West Nile virus — Eastern United States, 2001**

Editorial Note: The findings in this report demonstrate multifocal epizootic WNV activity across the eastern United States, including new areas in the southeast. In 2000, avian mortality surveillance indicated northward spread of epizootic activity from the New York City metropolitan area in late spring and early summer, and southward spread as far as North Carolina in late summer and early fall (1), a finding consistent with viral spread by migrating birds (2). The detection of WNV in Florida and southern Georgia in 2001, extends substantially the known distribution of this virus. Although first detected in these areas in 2001, WNV may have been introduced into these states earlier, but epizootic viral activity remained below the detection threshold of surveillance.

In 2000, avian mortality surveillance identified 4305 WNV-infected birds, 77 of which were identified by August 1 (1,3). The finding of 142 WNV-positive birds as of July 25, 2001, is twice the 2000 surveillance figures, and the geographic distribution of these birds differs between the 2 years. All of the birds identified in the early summer of 2000 were from four states (Connecticut, Massachusetts, New Jersey, and New York), compared with 44% of those identified as of July 25, 2001.

West Nile Virus Activity — Continued

Illness onset on approximately July 15 in the patient from Florida was the earliest of any person reported since the 1999 recognition of WNV in the United States. The extensive epizootic WNV activity and continued geographic expansion of the virus highlight the need for widespread implementation and intensification of surveillance, prevention, and control measures to minimize the risk for human and equine disease. Prevention activities have included the development and maintenance of long-term sustained mosquito-control programs using integrated pest management strategies and public education programs, emphasizing residential mosquito larval control and personal prevention measures to reduce mosquito exposure (4).

WNV detection in *Cx. salinarius* and *Ae. vexans* is of particular concern because these species more readily feed on mammals (including humans) than do *Cx. pipiens*, *Cx. restuans*, or *Cs. melanura*, which have a strong feeding preference for birds (5). However, mosquito-control programs in urban areas should continue to emphasize reduction of *Cx. pipiens* populations in the north and *Cx. quinquefasciatus* populations in the south. Although the role of these species in the direct transmission of WNV to humans is unclear, their role in the amplification of this virus and the closely related St. Louis encephalitis (SLE) virus in urban ecosystems is well established (6,7). The occurrence of WNV in Florida raises the possibility of transmission of this virus by other mosquito species such as *Cx. nigripalpus*, a primary vector of SLE virus in that state (8), and the possibility of a longer transmission season than is typical in regions with a more temperate climate.

The U.S. Geological Survey, CDC, and other federal, state, and local government agencies have collaborated to establish World-Wide Web-based maps to track the spread of WNV. These maps are available at http://cindi.usgs.gov/hazard/event/west_nile/west_nile.html. Additional information also is available from sites maintained by local and state health agencies. A partial listing of these sites is available at http://www.cdc.gov/ncidod/dvbid/westnile/city_states.htm.

References

1. Marfin AA, Petersen LR, Eidson M, et al. Widespread West Nile virus activity, eastern United States, 2000. *Emerg Infect Dis* 2001;7:730–5.
2. Rappole JH, Derricksen SR, Hubálek Z. Migratory birds and spread of West Nile virus in the Western Hemisphere. *Emerg Infect Dis* 2000;6:319–28.
3. CDC. Update: West Nile virus activity—Northeastern United States, January–August 7, 2000. *MMWR* 2000;49:714–7.
4. CDC. Epidemic/epizootic West Nile virus in the United States: revised guidelines for surveillance, prevention, and control, 2001. Available at <http://www.cdc.gov/ncidod/dvbid/westnile/resources/wnv-guidelines-apr-2001.pdf>. Accessed July 25, 2001.
5. CDC. Guidelines for arbovirus surveillance in the United States. Fort Collins, Colorado: US Department of Health and Human Services, Public Health Service, CDC, 1993. Available at <http://www.cdc.gov/ncidod/dvbid/arbor/arboguid.htm>. Accessed July 25, 2001.
6. Nasci RN, White DJ, Stirling H, et al. West Nile virus isolates from mosquitoes in New York and New Jersey, 1999. *Emerg Infect Dis* 2001;7:626–30.
7. Savage HM, Ceianu C, Nicolescu G, et al. Entomologic and avian investigations of an epidemic of West Nile fever in Romania in 1996, with serologic and molecular characterization of a virus isolate from mosquitoes. *Am J Trop Med Hyg* 1999;61:600–11.
8. Day JF. Predicting St. Louis encephalitis virus epidemics: lessons from recent, and not so recent, outbreaks. *Annu Rev Entomol* 2001;46:111–38.

Global Progress Toward Laboratory Containment of Wild Polioviruses, June 2001

When the World Health Assembly resolved to eradicate poliomyelitis in 1988, the estimated number of polio cases was 350,000; in 2000, approximately 3000 cases were reported (1). Two World Health Organization (WHO) regions (the Americas and Western Pacific) have been certified as polio-free, and a third (European Region) has been free of indigenous wild poliovirus transmission for nearly 3 years (3 years are required for WHO certification). As interruption of wild poliovirus circulation approaches, public health agencies are increasing efforts to minimize the risk for reintroduction of wild polioviruses from laboratory sources. This report describes the global plan for containing laboratory wild polioviruses and summarizes the steps being taken toward implementation.

Once wild poliovirus transmission ceases and laboratories are the only source of wild poliovirus, an increase in precautions will be needed to minimize the risk for reintroducing wild polioviruses from stored sources and for ensuring the safe handling and disposal of these materials, which include wild poliovirus infectious stocks, specimens from polio patients, and products of research or potentially infectious materials (i.e., throat, fecal, or environmental [water and sewage] specimens collected for any purpose at a time and in a geographic location where polio was endemic). Virology laboratories are the most likely sources of infectious materials, but other biomedical laboratories such as bacteriology, parasitology, gastroenterology, nutrition, pathology, and environmental also may store infectious materials.

The *WHO Global Action Plan for Laboratory Containment of Wild Polioviruses* (2), developed in collaboration with scientists, ministries of health, and vaccine manufacturers, was endorsed by a 1999 World Health Assembly resolution. The Global Certification Commission stated that a precondition of certification is adequate containment of wild polioviruses (3), and the plan outlines three implementation phases: preglobal eradication, postglobal eradication, and post-OPV (oral poliovirus vaccine) immunization.

During the preglobal eradication phase, countries in which wild poliovirus circulation has been interrupted appoint a national task force or coordinator to develop and oversee a national plan. The first step in the plan is to alert biomedical laboratories to the impending eradication of polio, encourage them to dispose appropriately of unneeded wild poliovirus or potentially infectious materials, and establish a national inventory of laboratories that retain such materials. The inventory will provide a list of laboratories to be informed of eradication progress and containment developments and to be notified when eradication occurs and implementation of additional biosafety requirements takes effect.

Many countries/territories are surveying and identifying laboratories for their capacity to store infectious materials (Table 1). Laboratories that do not have the capacity to store infectious materials or routinely do not keep specimens for long periods confirm their inability to serve as a storage facility and are eliminated as a site for wild poliovirus materials. Laboratories identified as having the capacity to store infectious materials are followed up to determine the materials they hold in storage.

The postglobal eradication phase begins soon after detection of the last wild poliovirus in the world. At that time, laboratories storing and handling infectious or potentially infectious materials prepare for certification by implementing biosafety conditions appropriate for the levels of risk presented by the materials under study and laboratory procedures in use. A further increase in biosafety requirements is anticipated when a

TABLE 1. Number and percentage of countries/territories with national task forces, national plans, initiated plans, identified biomedical laboratories, biomedical laboratories that handle infectious or potentially infectious material, and have submitted inventories, by World Health Organization (WHO) region, June 2001

| WHO region | No. countries/territories with task force or coordinator | | Region total (%) | | No. countries/territories with plan | | Region total (%) | | No. countries/territories that have initiated plan* | | Region total (%) | | No. laboratories identified for survey | | No. laboratories with potentially infectious materials | | No. countries/territories submitting an inventory of laboratories with potentially infectious materials | | Region total (%) | |
|------------------|----------------------------------------------------------|--------------|------------------|-------------------------------------|-------------------------------------|---------------|-----------------------------------------------------|--------------|-----------------------------------------------------|----------------------------------------|--------------------------------------------------------|---------------------------------------------------------------------------------------------------------|----------------------------------------|--------------|--------------------------------------------------------|--|---------------------------------------------------------------------------------------------------------|--|------------------|--|
| | No. countries/territories with task force or coordinator | Region total | (%) | No. countries/territories with plan | Region total | (%) | No. countries/territories that have initiated plan* | Region total | (%) | No. laboratories identified for survey | No. laboratories with potentially infectious materials | No. countries/territories submitting an inventory of laboratories with potentially infectious materials | Region total | (%) | | | | | | |
| African | 0 | 48 | — | 0 | 48 | — | 0 | 48 | — | 0 | 0 | 0 | 48 | — | | | | | | |
| Americas | 2 | 47 | (4%) | 2 | 47 | (4%) | 2 | 47 | (4%) | 16,781 | 21 | 0 | 47 | — | | | | | | |
| Eastern | | | | | | | | | | | | | | | | | | | | |
| Mediterranean | 17 | 24 | (71%) | 17 | 24 | (71%) | 7 | 24 | (29%) | 1,499 | 10 | 2 | 24 | (8%) | | | | | | |
| European | 48 | 51 | (98%) | 45 | 51 | (76%) | 36 | 51 | (53%) | 36,089 | 254 | 0 | 51 | — | | | | | | |
| South-East Asian | 7 | 10 | (70%) | 7 | 10 | (70%) | 1 | 10 | (10%) | 63 | 0 | 0 | 10 | — | | | | | | |
| Western Pacific | 36 | 36 | (100%) | 36 | 36 | (100%) | 36 | 36 | (100%) | 11,620 | 98 | 9 | 36 | (25%) | | | | | | |
| Total | 110 | 216 | (51%) | 107 | 216 | (50%) | 82 | 216 | (38%) | 66,052 | 383 | 11 | 216 | (5%) | | | | | | |

* Have initiated a survey of laboratories to identify those storing wild polioviruses and infectious or potentially infectious materials.

Containment of Wild Polioviruses — Continued

global decision is made on OPV cessation. WHO is working with manufacturers of inactive polio vaccine (IPV) to develop a plan for containing poliovirus strains used in manufacturing IPV and to formulate containment guidelines designed to minimize risk during the production of IPV.

The risk for accidental reintroduction of wild poliovirus into a community from a laboratory is possible if four conditions exist: 1) the presence of wild poliovirus infectious materials in a laboratory; 2) an event (e.g., break in standard procedure) that exposes workers to infectious materials containing poliovirus; 3) susceptible workers who replicate and shed the virus in their stool; and 4) susceptible persons in the community who are directly or indirectly exposed to an infected worker. Implementation of the plan cannot ensure absolute containment; however, it will minimize the likelihood of a situation in which the first three conditions occur. The fourth condition is linked to posteradication immunization policy decisions.

Progress is being made in implementing the first phase of laboratory containment (Table 1); 110 (51%) of 216 countries/territories have appointed a national task force and have created a plan. Eleven countries have submitted completed national inventories, and approximately 400 laboratories with wild poliovirus materials have been identified. In the Americas, laboratory containment activities are under way. Canada is in the final stages of preparing its national inventory and the United States is in the initial stage of its laboratory survey. In the Western Pacific, all member states have begun implementation and nine of 36 have finished their national inventory. Laboratory containment activities have increased substantially in the European Region as it prepares for certification; 48 of 51 member states have appointed a task force and 36 of these have started contacting laboratories. Although polio is still endemic in the South-East Asian, Eastern Mediterranean, and African regions, many polio-free countries in these regions have begun preparations for laboratory containment.

Reported by: Vaccines and Biologicals Dept, World Health Organization, Geneva, Switzerland. Respiratory and Enteric Viruses Br, Div of Viral and Rickettsial Diseases, National Center for Infectious Diseases; Vaccine Preventable Disease Eradication Div, National Immunization Program, CDC.

Editorial Note: Appropriate laboratory containment of wild poliovirus is critical to polio eradication. Progress toward implementation of the global plan is encouraging; a systematic and well-documented approach has been established to identify laboratories with infectious wild poliovirus or potentially infectious materials, and cooperation from laboratories and governments has been good throughout the world.

Implementing laboratory containment procedures is a complex process. Industrialized countries with well-developed research programs and laboratory infrastructure will require considerable time and effort for implementing survey and inventory activities. Countries with less developed biomedical research programs and laboratory infrastructure generally do not have laboratories that store infectious materials. Such countries can more easily compile a list of laboratories and identify those with infectious wild poliovirus or potentially infectious materials. Technical expertise for assisting countries with their national plans and implementing activities is available from members of the Global Laboratory Network for Polio Eradication, which comprises 124 national (or subnational) laboratories, 16 regional reference laboratories, and seven specialized laboratories.

The link between certification and laboratory containment activities has evolved; laboratory containment procedures were not part of the certification process when the

Containment of Wild Polioviruses — Continued

Americas was certified free of polio in 1994. The Pan American Health Organization is working with member governments to meet the requirements outlined in the global plan. The most progress toward completion of the first phase of the plan has been reported from the Western Pacific Region where laboratory containment activities were an integral part of the certification process. The European Region is integrating containment into the regional certification process.

WHO member states will be responsible for laboratory containment within their respective countries. The containment process will be monitored by national authorities, national committees for polio eradication, and the Regional and Global Certification commissions. Before global certification can occur, as anticipated in 2005, all countries of the world must demonstrate that they have minimized the risk for reintroducing wild poliovirus from their laboratories to a polio-free world.

References

1. World Health Assembly. Polio eradication by the year 2000. Resolutions of the 41st World Health Assembly. Geneva, Switzerland: World Health Organization, 1988 (Resolution 41.28).
2. Department of Vaccines and Biologicals. World Health Organization global action plan for laboratory containment of wild polioviruses. Geneva, Switzerland: World Health Organization, 1999 (Reference WHO/V&B/99.32).
3. Department of Vaccines and Biologicals. Report of the third meeting of the Global Commission for the Certification of the Eradication of Polio, July 9, 1998. Geneva, Switzerland: World Health Organization, 1999 (Reference WHO/EPI/GEN/981.17).

Heat-Related Deaths — Los Angeles County, California, 1999–2000, and United States, 1979–1998

Heat-related deaths typically occur during summer months. Many of these deaths are preventable. This report describes four cases of heat-related deaths in Los Angeles County, California, during 1999–2000, compares age-, sex-, and race-specific rates in Los Angeles County and the United States during 1979–1998, and summarizes trends in the United States during 1979–1998. Relatives, neighbors, and caretakers of persons at risk for heat-related death should frequently evaluate heat-related hazards, recognize symptoms of heat-related morbidity, and take appropriate preventive action.

Case Reports

Case 1. In June 1999, a 4-month-old boy was found dead in his parents' car. The child had been left in the car with windows closed for 6 hours. Death was attributed to hyperthermia. The temperature inside the car was 118 F (47.8 C), and the outside temperature was 96.0 F (35.6 C).

Case 2. In July 1999, an 81-year-old woman with a medical history of dementia and heart disease was found dead on the roof of the residential-care center where she lived. She had last been seen alive 64 hours earlier and had been reported missing for 24 hours before she was found. The decedent wore a "wanderer" bracelet that sounded an alarm when she exited through the front door of the center. The roof door was not equipped with an alarm but was usually kept locked. Death was attributed to hyperthermia. The ambient temperature on the roof was 96.0 F (35.6 C) at the time the decedent was found.

Case 3. In July 2000, a 46-year-old man was found confused and rolling on the pavement of a parking lot near his residence. When an ambulance arrived, he was unconscious and had had seizures. The local ambient temperature was 109.0 F (42.8 C) at the

Heat-Related Deaths — Continued

time he was found. At the emergency department, his temperature was 107.0 F (41.7 C). He died 2 days later in a hospital. Laboratory tests showed a blood alcohol level of 93 mg/dL (the legal blood alcohol limit in California is 80 mg/dL) and a positive screen for cocaine. Death was attributed to hyperthermia.

Case 4. In August 2000, a 65-year-old woman was found unresponsive in the backyard of her residence. She was admitted to a hospital where she died 2 days later. Her body temperature on admission was 108.0 F (42.2 C). The decedent had a medical history of insulin-dependent diabetes, hypertension, and heart disease. The underlying cause of death was hyperthermia.

U.S. Trends and Summary of Rates in Los Angeles County

During 1979–1998 (the most recent years for which national data are available), 7421 deaths in the United States were heat-related* with a median of 274 deaths per year (range: 148–1700), and a median heat-related death rate of 0.1 per 100,000 population (range: 0.1–0.8). Heat-related death rates appear to be stable over time in all age groups with the highest mortality among persons aged ≥ 65 years (Figure 1).

During 1979–1998, the age-adjusted[†] heat-related death rate in Los Angeles County was 44% lower than that in the general U.S. population (0.90 per 100,000 population versus 0.16). Los Angeles residents aged ≥ 65 years were more likely than residents aged < 65 years to die from exposure to excessive heat, but the rate ratio was smaller than in the general U.S. population (2.4 versus 7.4). Men in Los Angeles were more likely than women to die from exposure to excessive heat, and the rate ratio of 2.0 was the same as in the general U.S. population. Although blacks in Los Angeles County were more likely than whites to die from exposure to excessive heat, the rate ratio was smaller than in the general U.S. population (1.4 versus 4.9). Persons of other (nonblack and nonwhite[§]) races in Los Angeles County were less likely than whites to die from exposure to excessive heat, but the rate ratio was smaller than in the general U.S. population (0.5 versus 0.8).

Reported by: L Sathyavagiswaran, MD, Dept of the Coroner, Los Angeles County; JE Fielding, MD, D Dassy, MD, Los Angeles County Dept of Health Svcs. Health Studies Br, Div of Environmental Hazards and Health Effects, National Center for Environmental Health; and EIS officers, CDC.


Editorial Note: These case reports illustrate some of the risks for hyperthermia. The primary risk factors include age (i.e., increasing age, except children aged < 5 years who are at higher risk than older children), behavior (e.g., low fluid intake, excessive exercise, prolonged stay in nonair-conditioned places, and alcohol and/or drug use) (1), chronic disease (e.g., cardiac or mental illness) (2,3), prescription drugs (e.g., psychotropic medication) (3), living conditions (e.g., low income, residence in urban areas, no access to air-conditioning, and social isolation) (1), and prolonged outdoor activities (e.g., agricultural work and recreational running).

*Underlying cause of death attributed to "excessive heat exposure," classified according to the *International Classification of Diseases, Ninth Revision (ICD-9)*, code E900.0, "due to weather conditions" (deaths); code E900.1, "of manmade origin" (deaths); or code E900.9, "of unspecified origin" (deaths). Data were obtained from the Compressed Mortality File of CDC's National Center for Health Statistics, which contains information from death certificates filed in the 50 states and the District of Columbia.

[†] Rates were age-adjusted to the 2000 U.S. standard population.

[§] Race in the Compressed Mortality File was categorized as white, black, and other.

Heat-Related Deaths — Continued

FIGURE 1. Rate* of heat-related deaths, by age group — United States, 1979–1998

* Per 100,000 population.

Heat-related illness can begin as sunburn and fatigue and progress to heat cramps, heat exhaustion, and heatstroke. The two most serious types of heat-related illness are heat exhaustion (heavy sweating, paleness, muscle cramps, tiredness or weakness, dizziness or headache, nausea or vomiting, and faintness) and heatstroke (oral temperature of ≥ 103.0 F [≥ 39.4 C]; rapid, strong pulse; red, hot, and dry or sweaty skin; throbbing headache or dizziness; nausea; confusion; and unconsciousness). Untreated heat exhaustion can progress to heatstroke (4), a medical emergency that can develop in ≤ 24 hours (5). Even when treated, the death rate for heatstroke may be as high as 33% (6). Permanent neurologic damage occurs in up to 17% of survivors (7), and its likelihood increases with longer duration of heatstroke (4).

Spending time in an air-conditioned area is the most important factor in preventing heat-related deaths (2). During the 1999 heat wave in Cincinnati, Ohio, three of 18 heat-related deaths occurred in assisted-care facilities for persons with mental illness that did not have air-conditioning (8). The use of fans does not appear to be protective. If exposure to heat cannot be avoided, prevention measures should include reducing, eliminating, or rescheduling strenuous activities; frequently drinking water or nonalcoholic fluids; frequently taking showers; wearing light-weight and light-colored clothing; and avoiding direct sunshine.


Heat-Related Deaths — Continued

Because heat-related morbidity and mortality could increase with more periods of extreme heat in future summers (9), many cities have developed heat emergency response plans. These response plans use information on risk factors and meteorologic information to implement prevention strategies that reduce morbidity and mortality from excessive heat (1). A heat response plan also should address rolling energy black-outs in areas that use air-conditioning to mitigate many of the factors that increase the risk for heat-related morbidity and mortality. To defray energy costs, support of low-income populations may be necessary to allow the use of air-conditioning during summer months.

References

1. McGeehin MA, Mirabelli M. The potential impacts of climate variability and change on temperature-related morbidity and mortality in the United States. *Environ Health Perspec* 2001;109:185–90.
2. Semenza JC, Rubin CH, Falter KH, et al. Heat-related deaths during the 1995 heat wave in Chicago. *N Engl J Med* 1996;335:84–90.
3. Batscha CL. Heat stroke: keeping your clients cool in the summer. *J Psychosoc Nurs Ment Health Serv* 1997;35:12–7.
4. Knochel JP. Environmental heat illness: an eclectic review. *Arch Intern Med* 1974;133:841–64.
5. Jones TS, Liang AP, Kilbourne EM, et al. Morbidity and mortality associated with the July 1980 heat wave in St. Louis and Kansas City, Mo. *JAMA* 1982;247:3327–31.
6. Vicario SJ, Okabajue R, Haltom T. Rapid cooling in classic heatstroke: effect on mortality rates. *Am J Emerg Med* 1986;4:394–8.
7. Hart GR, Anderson RJ, Crumpler CP, Shulkin A, Reed G, Knochel JP. Epidemic classical heat stroke: clinical characteristics and course of 28 patients. *Medicine* 1982;61:189–97.
8. Kaiser R, Rubin CH, Henderson A, et al. Heat-related death and mental illness during the 1999 Cincinnati heat wave. *Am J Forensic Med Pathol*(in press).
9. Gaffen DJ. Increased summertime heat stress in the US. *Nature* 1998;396:529–30.

FIGURE I. Selected notifiable disease reports, United States, comparison of provisional 4-week totals ending July 21, 2001, with historical data



* No measles cases were reported for the current 4-week period yielding a ratio for week 29 of zero (0).

† Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

TABLE I. Summary of provisional cases of selected notifiable diseases, United States, cumulative, week ending July 21, 2001 (29th Week)

| | Cum. 2001 | | Cum. 2001 |
|-------------------------------------------|-----------|------------------------------------------|-----------|
| Anthrax | - | Poliomyelitis, paralytic | - |
| Brucellosis* | 38 | Psittacosis* | 7 |
| Cholera | 4 | Q fever* | 11 |
| Cyclosporiasis* | 63 | Rabies, human | 1 |
| Diphtheria | 1 | Rocky Mountain spotted fever (RMSF) | 217 |
| Ehrlichiosis: human granulocytic (HGE)* | 68 | Rubella, congenital syndrome | - |
| human monocytic (HME)* | 28 | Streptococcal disease, invasive, group A | 2,182 |
| Encephalitis: California serogroup viral* | 5 | Streptococcal toxic-shock syndrome* | 33 |
| eastern equine* | 1 | Syphilis, congenital† | 84 |
| St. Louis* | - | Tetanus | 13 |
| western equine* | - | Toxic-shock syndrome | 66 |
| Hansen disease (leprosy)* | 39 | Trichinosis | 11 |
| Hantavirus pulmonary syndrome*† | 4 | Tularemia* | 47 |
| Hemolytic uremic syndrome, postdiarrheal* | 52 | Typhoid fever | 139 |
| HIV infection, pediatric*§ | 98 | Yellow fever | - |
| Plague | 2 | | |

-: No reported cases.

*Not notifiable in all states.

† Updated monthly from reports to the Division of HIV/AIDS Prevention — Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention (NCHSTP). Last update June 26, 2001.

§ Updated from reports to the Division of STD Prevention, NCHSTP.

TABLE II. Provisional cases of selected notifiable diseases, United States, weeks ending July 21, 2001, and July 22, 2000 (29th Week)

| Reporting Area | AIDS | | Chlamydia [†] | | Cryptosporidiosis | | <i>Escherichia coli</i> O157:H7* | | | |
|----------------|------------------------|-----------|------------------------|-----------|-------------------|-----------|----------------------------------|-----------|-----------|-----------|
| | Cum. 2001 [§] | Cum. 2000 | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2000 | NETSS | | PHLIS | |
| | | | | | | | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2000 |
| UNITED STATES | 19,145 | 21,713 | 362,229 | 379,078 | 926 | 902 | 958 | 1,686 | 765 | 1,564 |
| NEW ENGLAND | 746 | 1,279 | 12,290 | 12,643 | 38 | 56 | 111 | 172 | 76 | 183 |
| Maine | 20 | 20 | 662 | 770 | 4 | 9 | 14 | 9 | 14 | 14 |
| N.H. | 17 | 21 | 698 | 575 | 2 | 6 | 14 | 14 | 11 | 18 |
| Vt. | 10 | 17 | 330 | 298 | 13 | 14 | 4 | 20 | 2 | 22 |
| Mass. | 411 | 837 | 5,573 | 5,344 | 12 | 17 | 58 | 80 | 28 | 72 |
| R.I. | 53 | 48 | 1,552 | 1,408 | 3 | 2 | 6 | 8 | 4 | 10 |
| Conn. | 235 | 336 | 3,475 | 4,248 | 4 | 8 | 15 | 41 | 17 | 47 |
| MID. ATLANTIC | 3,974 | 5,227 | 41,389 | 36,436 | 108 | 164 | 82 | 188 | 59 | 136 |
| Upstate N.Y. | 322 | 539 | 7,355 | 640 | 49 | 40 | 61 | 115 | 33 | 38 |
| N.Y. City | 1,996 | 2,852 | 15,718 | 15,193 | 51 | 91 | 4 | 14 | 7 | 8 |
| N.J. | 960 | 1,024 | 5,521 | 7,138 | 4 | 7 | 17 | 59 | 19 | 55 |
| Pa. | 696 | 812 | 12,795 | 13,465 | 4 | 26 | N | N | - | 35 |
| E.N. CENTRAL | 1,408 | 2,068 | 52,783 | 64,574 | 296 | 212 | 220 | 349 | 161 | 279 |
| Ohio | 237 | 290 | 7,727 | 17,278 | 62 | 24 | 62 | 56 | 47 | 72 |
| Ind. | 165 | 188 | 8,037 | 7,118 | 32 | 12 | 37 | 43 | 21 | 43 |
| Ill. | 665 | 1,191 | 14,031 | 18,527 | 1 | 34 | 47 | 96 | 41 | 69 |
| Mich. | 261 | 296 | 16,524 | 12,748 | 72 | 35 | 26 | 51 | 27 | 42 |
| Wis. | 80 | 103 | 6,464 | 8,903 | 129 | 107 | 48 | 103 | 25 | 53 |
| W.N. CENTRAL | 454 | 487 | 18,357 | 21,202 | 91 | 76 | 123 | 220 | 140 | 267 |
| Minn. | 85 | 86 | 3,412 | 4,352 | 32 | 11 | 36 | 47 | 63 | 80 |
| Iowa | 47 | 52 | 1,858 | 2,761 | 28 | 25 | 28 | 44 | 24 | 57 |
| Mo. | 218 | 226 | 6,616 | 7,279 | 10 | 12 | 22 | 58 | 29 | 56 |
| N. Dak. | 1 | 2 | 520 | 493 | 3 | 5 | 1 | 7 | 10 | 14 |
| S. Dak. | 18 | 4 | 957 | 996 | 5 | 8 | 9 | 14 | 8 | 19 |
| Nebr. | 39 | 31 | 1,696 | 2,052 | 13 | 12 | 16 | 35 | - | 31 |
| Kans. | 46 | 86 | 3,298 | 3,269 | - | 3 | 11 | 15 | 6 | 10 |
| S. ATLANTIC | 6,167 | 5,749 | 66,953 | 70,449 | 161 | 134 | 95 | 119 | 48 | 142 |
| Del. | 116 | 94 | 1,606 | 1,570 | 1 | 4 | 1 | 1 | 3 | - |
| Md. | 751 | 690 | 6,670 | 7,468 | 27 | 7 | 7 | 13 | 1 | 1 |
| D.C. | 465 | 389 | 1,663 | 1,771 | 9 | 5 | - | - | U | U |
| Va. | 501 | 380 | 9,657 | 8,997 | 10 | 4 | 24 | 24 | 18 | 29 |
| W. Va. | 49 | 31 | 1,275 | 1,185 | 1 | 3 | 3 | 8 | 1 | 5 |
| N.C. | 402 | 312 | 8,914 | 12,103 | 18 | 15 | 26 | 22 | 13 | 37 |
| S.C. | 350 | 455 | 6,208 | 5,105 | - | - | 2 | 8 | 3 | 9 |
| Ga. | 757 | 605 | 12,438 | 14,459 | 56 | 63 | 14 | 15 | 2 | 26 |
| Fla. | 2,776 | 2,793 | 18,522 | 17,791 | 39 | 33 | 18 | 28 | 7 | 35 |
| E.S. CENTRAL | 977 | 1,051 | 26,994 | 27,468 | 24 | 27 | 44 | 61 | 41 | 55 |
| Ky. | 201 | 127 | 4,872 | 4,418 | 3 | 3 | 15 | 21 | 21 | 19 |
| Tenn. | 293 | 438 | 8,022 | 7,946 | 5 | 6 | 20 | 23 | 18 | 28 |
| Ala. | 224 | 255 | 7,578 | 8,228 | 9 | 10 | 8 | 5 | - | 4 |
| Miss. | 259 | 231 | 6,522 | 6,876 | 7 | 8 | 1 | 12 | 2 | 4 |
| W.S. CENTRAL | 2,058 | 2,333 | 56,698 | 57,283 | 18 | 47 | 35 | 150 | 52 | 183 |
| Ark. | 104 | 111 | 4,094 | 3,587 | 3 | 1 | 4 | 36 | - | 30 |
| La. | 472 | 318 | 9,388 | 10,525 | 7 | 10 | 2 | 10 | 23 | 27 |
| Okla. | 107 | 184 | 5,815 | 4,732 | 6 | 4 | 12 | 9 | 14 | 7 |
| Tex. | 1,375 | 1,720 | 37,401 | 38,439 | 2 | 32 | 17 | 95 | 15 | 119 |
| MOUNTAIN | 714 | 806 | 19,639 | 22,173 | 62 | 42 | 115 | 179 | 77 | 136 |
| Mont. | 12 | 9 | 1,015 | 825 | 5 | 8 | 6 | 20 | - | - |
| Idaho | 15 | 13 | 909 | 1,031 | 7 | 3 | 15 | 23 | - | 18 |
| Wyo. | 1 | 7 | 454 | 410 | 1 | 5 | 5 | 10 | 1 | 6 |
| Colo. | 140 | 200 | 2,908 | 6,662 | 19 | 12 | 50 | 67 | 44 | 50 |
| N. Mex. | 56 | 87 | 3,066 | 2,732 | 12 | 2 | 8 | 6 | 6 | 7 |
| Ariz. | 295 | 224 | 7,769 | 7,070 | 4 | 2 | 15 | 28 | 9 | 22 |
| Utah | 63 | 81 | 906 | 1,339 | 12 | 8 | 10 | 21 | 16 | 27 |
| Nev. | 132 | 185 | 2,612 | 2,104 | 2 | 2 | 6 | 4 | 1 | 6 |
| PACIFIC | 2,647 | 2,713 | 67,126 | 66,850 | 128 | 144 | 133 | 248 | 111 | 183 |
| Wash. | 290 | 275 | 7,339 | 7,054 | N | U | 32 | 89 | 31 | 104 |
| Oreg. | 112 | 88 | 2,159 | 3,868 | 14 | 9 | 22 | 41 | 17 | 46 |
| Calif. | 2,204 | 2,255 | 54,101 | 52,618 | 111 | 135 | 69 | 101 | 60 | 25 |
| Alaska | 13 | 12 | 1,492 | 1,354 | - | - | 2 | 9 | - | 1 |
| Hawaii | 28 | 83 | 2,035 | 1,956 | 3 | - | 8 | 8 | 3 | 7 |
| Guam | 9 | 13 | - | 263 | - | - | N | N | U | U |
| P.R. | 580 | 706 | 1,611 | U | - | - | - | 5 | U | U |
| V.I. | 2 | 24 | 53 | - | - | - | - | - | U | U |
| Amer. Samoa | - | - | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | - | 69 | U | - | U | - | U | U | U |

N: Not notifiable. U: Unavailable. -: No reported cases. C.N.M.I.: Commonwealth of Northern Mariana Islands.

* Individual cases can be reported through both the National Electronic Telecommunications System for Surveillance (NETSS) and the Public Health Laboratory Information System (PHLIS).

[†] Chlamydia refers to genital infections caused by *C. trachomatis*. Totals reported to the Division of STD Prevention, NCHSTP.

[§] Updated monthly from reports to the Division of HIV/AIDS Prevention — Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention. Last update June 26, 2001.

TABLE II. (Cont'd) Provisional cases of selected notifiable diseases, United States, weeks ending July 21, 2001, and July 22, 2000 (29th Week)

| Reporting Area | Gonorrhea | | Hepatitis C: Non-A, Non-B | | Legionellosis | | Listeriosis | Lyme Disease | |
|----------------|--------------|--------------|------------------------------|--------------|---------------|--------------|--------------|--------------|--------------|
| | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2001 | Cum. 2000 |
| UNITED STATES | 165,190 | 189,589 | 1,190 | 1,864 | 425 | 446 | 223 | 2,580 | 6,811 |
| NEW ENGLAND | 3,387 | 3,566 | 14 | 15 | 20 | 25 | 29 | 728 | 1,709 |
| Maine | 73 | 46 | - | 1 | 1 | 2 | - | - | - |
| N.H. | 87 | 59 | - | - | 5 | 2 | 1 | 69 | 36 |
| Vt. | 41 | 32 | 6 | 3 | 4 | 2 | 1 | 3 | 12 |
| Mass. | 1,707 | 1,429 | 8 | 8 | 5 | 11 | 15 | 150 | 706 |
| R.I. | 399 | 341 | - | 3 | 1 | 3 | 1 | 109 | 78 |
| Conn. | 1,080 | 1,659 | - | - | 4 | 5 | 11 | 397 | 877 |
| MID. ATLANTIC | 20,055 | 20,566 | 48 | 402 | 45 | 109 | 32 | 1,135 | 3,853 |
| Upstate N.Y. | 4,396 | 3,752 | 34 | 17 | 28 | 32 | 14 | 891 | 1,106 |
| N.Y. City | 6,603 | 6,377 | - | - | 6 | 16 | 6 | 1 | 144 |
| N.J. | 2,951 | 4,155 | - | 360 | 5 | 9 | 7 | 85 | 1,727 |
| Pa. | 6,105 | 6,282 | 14 | 25 | 6 | 52 | 5 | 158 | 876 |
| E.N. CENTRAL | 28,066 | 37,999 | 105 | 144 | 116 | 118 | 26 | 145 | 491 |
| Ohio | 4,537 | 10,145 | 7 | 4 | 61 | 41 | 7 | 46 | 26 |
| Ind. | 3,295 | 3,268 | 1 | - | 12 | 22 | 4 | 3 | 10 |
| Ill. | 8,608 | 11,349 | 10 | 15 | - | 15 | - | - | 27 |
| Mich. | 9,481 | 9,447 | 87 | 125 | 29 | 21 | 13 | - | 13 |
| Wis. | 2,145 | 3,790 | - | - | 14 | 19 | 2 | 96 | 415 |
| W.N. CENTRAL | 7,630 | 9,309 | 397 | 329 | 34 | 28 | 6 | 108 | 83 |
| Minn. | 1,091 | 1,739 | 3 | 5 | 7 | 1 | - | 69 | 42 |
| Iowa | 428 | 589 | - | 1 | 6 | 6 | - | 18 | 4 |
| Mo. | 3,962 | 4,585 | 387 | 315 | 12 | 14 | 3 | 15 | 22 |
| N. Dak. | 16 | 39 | - | - | 1 | - | - | - | - |
| S. Dak. | 144 | 154 | - | - | 3 | 1 | - | - | - |
| Nebr. | 556 | 774 | 3 | 3 | 4 | 2 | 1 | 2 | 2 |
| Kans. | 1,433 | 1,429 | 4 | 5 | 1 | 4 | 2 | 4 | 13 |
| S. ATLANTIC | 41,409 | 49,208 | 59 | 54 | 94 | 80 | 37 | 365 | 551 |
| Del. | 907 | 903 | - | 2 | 2 | 5 | - | 22 | 103 |
| Md. | 3,712 | 4,994 | 10 | 6 | 22 | 25 | 4 | 224 | 338 |
| D.C. | 1,468 | 1,275 | - | 2 | 7 | - | - | 7 | 2 |
| Va. | 5,253 | 5,541 | - | 3 | 11 | 13 | 6 | 72 | 63 |
| W. Va. | 341 | 365 | 6 | 12 | N | N | 4 | 8 | 17 |
| N.C. | 7,980 | 9,829 | 10 | 13 | 5 | 8 | 2 | 16 | 20 |
| S.C. | 4,499 | 4,876 | 4 | 1 | 4 | 2 | 3 | 2 | 2 |
| Ga. | 6,762 | 9,099 | - | 2 | 6 | 4 | 8 | - | - |
| Fla. | 10,487 | 12,326 | 29 | 13 | 37 | 23 | 10 | 14 | 6 |
| E. S. CENTRAL | 17,389 | 19,653 | 124 | 267 | 34 | 15 | 11 | 17 | 22 |
| Ky. | 1,878 | 1,879 | 4 | 18 | 8 | 7 | 4 | 7 | 5 |
| Tenn. | 5,347 | 6,215 | 41 | 59 | 16 | 5 | 3 | 6 | 13 |
| Ala. | 5,921 | 6,509 | 2 | 7 | 8 | 2 | 4 | 4 | 2 |
| Miss. | 4,243 | 5,050 | 77 | 183 | 2 | 1 | - | - | 2 |
| W.S. CENTRAL | 27,325 | 29,905 | 161 | 503 | 5 | 18 | 5 | 7 | 42 |
| Ark. | 2,521 | 1,959 | 3 | 4 | - | - | 1 | - | 3 |
| La. | 6,539 | 7,440 | 74 | 268 | 2 | 7 | - | 1 | 3 |
| Okla. | 2,609 | 2,038 | 3 | 4 | 3 | 1 | 1 | - | - |
| Tex. | 15,656 | 18,468 | 81 | 227 | - | 10 | 3 | 6 | 36 |
| MOUNTAIN | 5,529 | 5,754 | 201 | 39 | 34 | 22 | 23 | 8 | 4 |
| Mont. | 53 | 26 | 1 | 2 | - | 1 | - | - | - |
| Idaho | 39 | 50 | 1 | 3 | 2 | 4 | 1 | 3 | 1 |
| Wyo. | 32 | 30 | 159 | 2 | 3 | - | 1 | 3 | 2 |
| Colo. | 1,748 | 1,763 | 13 | 6 | 10 | 6 | 3 | 1 | - |
| N. Mex. | 487 | 581 | 10 | 11 | 2 | 1 | 6 | - | - |
| Ariz. | 2,152 | 2,375 | 9 | 11 | 11 | 5 | 6 | - | - |
| Utah | 79 | 140 | 2 | - | 4 | 5 | 1 | - | - |
| Nev. | 939 | 789 | 6 | 4 | 2 | - | 5 | 1 | 1 |
| PACIFIC | 14,400 | 13,629 | 81 | 111 | 43 | 31 | 54 | 67 | 56 |
| Wash. | 1,607 | 1,217 | 16 | 16 | 6 | 11 | 3 | 2 | 3 |
| Oreg. | 307 | 503 | 9 | 21 | N | N | 1 | 5 | 4 |
| Calif. | 11,942 | 11,469 | 56 | 72 | 33 | 20 | 49 | 58 | 48 |
| Alaska | 213 | 182 | - | - | - | - | - | 2 | 1 |
| Hawaii | 331 | 258 | - | 2 | 4 | - | 1 | N | N |
| Guam | - | 26 | - | 2 | - | - | - | - | - |
| P.R. | 438 | 293 | 1 | 1 | 2 | - | - | N | N |
| V.I. | 6 | - | - | - | - | - | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | - | U | U |
| C.N.M.I. | 7 | U | - | U | - | U | - | - | U |

N: Not notifiable.

U: Unavailable.

-: No reported cases.

TABLE II. (Cont'd) Provisional cases of selected notifiable diseases, United States, weeks ending July 21, 2001, and July 22, 2000 (29th Week)

| Reporting Area | Malaria | | Rabies, Animal | | Salmonellosis* | | | |
|----------------|--------------|--------------|----------------|--------------|----------------|--------------|--------------|--------------|
| | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2000 | NETSS | | PHLIS | |
| | | | | | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2000 |
| UNITED STATES | 529 | 694 | 3,194 | 3,688 | 15,621 | 18,150 | 12,839 | 16,188 |
| NEW ENGLAND | 34 | 34 | 328 | 403 | 1,209 | 1,128 | 1,032 | 1,179 |
| Maine | 3 | 4 | 40 | 80 | 111 | 80 | 93 | 61 |
| N.H. | 2 | 1 | 7 | 8 | 106 | 72 | 108 | 79 |
| Vt. | - | 2 | 37 | 37 | 39 | 65 | 38 | 61 |
| Mass. | 11 | 13 | 116 | 129 | 679 | 669 | 460 | 659 |
| R.I. | 3 | 5 | 29 | 18 | 66 | 45 | 85 | 84 |
| Conn. | 15 | 9 | 99 | 131 | 208 | 197 | 248 | 235 |
| MID. ATLANTIC | 97 | 156 | 509 | 655 | 1,643 | 2,575 | 1,974 | 2,657 |
| Upstate N.Y. | 28 | 31 | 381 | 400 | 578 | 582 | 479 | 675 |
| N.Y. City | 44 | 84 | 12 | 6 | 460 | 664 | 661 | 679 |
| N.J. | 19 | 21 | 95 | 88 | 419 | 635 | 413 | 501 |
| Pa. | 6 | 20 | 21 | 161 | 186 | 694 | 421 | 802 |
| E.N. CENTRAL | 56 | 83 | 44 | 54 | 2,270 | 2,524 | 1,835 | 1,600 |
| Ohio | 14 | 12 | 16 | 13 | 694 | 585 | 544 | 583 |
| Ind. | 12 | 4 | 1 | - | 249 | 285 | 241 | 315 |
| Ill. | 1 | 43 | 4 | 9 | 575 | 816 | 429 | 1 |
| Mich. | 19 | 17 | 17 | 23 | 396 | 476 | 397 | 510 |
| Wis. | 10 | 7 | 6 | 9 | 356 | 362 | 224 | 191 |
| W.N. CENTRAL | 23 | 36 | 188 | 336 | 945 | 1,185 | 1,072 | 1,346 |
| Minn. | 6 | 13 | 20 | 50 | 259 | 262 | 355 | 361 |
| Iowa | 3 | 1 | 43 | 47 | 158 | 174 | 163 | 183 |
| Mo. | 8 | 9 | 16 | 23 | 252 | 375 | 357 | 446 |
| N. Dak. | - | 2 | 24 | 83 | 14 | 27 | 38 | 49 |
| S. Dak. | - | - | 21 | 64 | 72 | 46 | 63 | 58 |
| Nebr. | 2 | 5 | 4 | - | 69 | 109 | - | 88 |
| Kans. | 4 | 6 | 60 | 69 | 121 | 192 | 96 | 161 |
| S. ATLANTIC | 156 | 152 | 1,181 | 1,283 | 3,885 | 3,272 | 2,340 | 2,816 |
| Del. | 1 | 3 | 18 | 20 | 44 | 58 | 43 | 68 |
| Md. | 66 | 50 | 153 | 245 | 409 | 398 | 389 | 382 |
| D.C. | 10 | 12 | - | - | 39 | 32 | U | U |
| Va. | 30 | 31 | 236 | 332 | 747 | 456 | 495 | 463 |
| W. Va. | 1 | 2 | 76 | 69 | 54 | 74 | 71 | 78 |
| N.C. | 7 | 11 | 324 | 313 | 549 | 432 | 459 | 488 |
| S.C. | 4 | 1 | 78 | 76 | 376 | 304 | 345 | 268 |
| Ga. | 8 | 4 | 174 | 157 | 564 | 522 | 351 | 822 |
| Fla. | 29 | 38 | 122 | 71 | 1,103 | 996 | 187 | 247 |
| E.S. CENTRAL | 15 | 22 | 110 | 105 | 935 | 952 | 714 | 800 |
| Ky. | 5 | 6 | 11 | 15 | 171 | 188 | 110 | 141 |
| Tenn. | 7 | 5 | 71 | 56 | 264 | 218 | 302 | 367 |
| Ala. | 3 | 10 | 28 | 34 | 296 | 257 | 235 | 246 |
| Miss. | - | 1 | - | - | 204 | 289 | 67 | 46 |
| W.S. CENTRAL | 6 | 42 | 504 | 529 | 1,226 | 2,266 | 1,079 | 1,398 |
| Ark. | 3 | 1 | 19 | - | 287 | 281 | 92 | 245 |
| La. | 1 | 7 | - | 1 | 249 | 395 | 344 | 309 |
| Okla. | 1 | 4 | 43 | 36 | 144 | 176 | 132 | 141 |
| Tex. | 1 | 30 | 442 | 492 | 546 | 1,414 | 511 | 703 |
| MOUNTAIN | 29 | 29 | 127 | 143 | 1,098 | 1,387 | 755 | 1,323 |
| Mont. | 2 | 1 | 20 | 39 | 40 | 61 | - | - |
| Idaho | 3 | 2 | 2 | 1 | 72 | 76 | 4 | 70 |
| Wyo. | - | - | 20 | 37 | 34 | 40 | 22 | 32 |
| Colo. | 15 | 14 | - | - | 300 | 414 | 276 | 393 |
| N. Mex. | 1 | - | 6 | 13 | 128 | 122 | 106 | 121 |
| Ariz. | 3 | 5 | 76 | 50 | 329 | 323 | 216 | 347 |
| Utah | 3 | 3 | 2 | 2 | 123 | 210 | 108 | 217 |
| Nev. | 2 | 4 | 1 | 1 | 72 | 141 | 23 | 143 |
| PACIFIC | 113 | 140 | 203 | 180 | 2,410 | 2,861 | 2,038 | 3,069 |
| Wash. | 4 | 13 | - | - | 231 | 236 | 358 | 349 |
| Oreg. | 5 | 23 | - | 4 | 120 | 174 | 167 | 223 |
| Calif. | 96 | 96 | 166 | 152 | 1,839 | 2,308 | 1,332 | 2,357 |
| Alaska | 1 | - | 37 | 24 | 24 | 31 | 2 | 23 |
| Hawaii | 7 | 8 | - | - | 196 | 112 | 179 | 117 |
| Guam | - | - | - | - | - | 17 | U | U |
| P.R. | 3 | 4 | 61 | 42 | 310 | 314 | U | U |
| V.I. | - | - | - | - | - | - | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | U | U | U | 7 | U | U | U |

N: Not notifiable. U: Unavailable. -: No reported cases.

* Individual cases can be reported through both the National Electronic Telecommunications System for Surveillance (NETSS) and the Public Health Laboratory Information System (PHLIS).

TABLE II. (Cont'd) Provisional cases of selected notifiable diseases, United States, weeks ending July 21, 2001, and July 22, 2000 (29th Week)

| Reporting Area | Shigellosis* | | | | Syphilis (Primary & Secondary) | | Tuberculosis | |
|----------------|--------------|--------------|--------------|--------------|-----------------------------------|--------------|--------------|--------------|
| | NETSS | | PHLIS | | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2000 |
| | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2000 | | | | |
| UNITED STATES | 7,538 | 11,190 | 3,766 | 6,191 | 2,986 | 3,370 | 6,346 | 7,543 |
| NEW ENGLAND | 120 | 205 | 106 | 193 | 28 | 48 | 227 | 215 |
| Maine | 6 | 6 | 1 | - | - | 1 | 7 | 8 |
| N.H. | 2 | 4 | 2 | 7 | 1 | 1 | 11 | 8 |
| Vt. | 3 | 2 | 2 | - | 2 | - | 2 | 3 |
| Mass. | 79 | 145 | 63 | 131 | 16 | 32 | 122 | 124 |
| R.I. | 8 | 12 | 14 | 19 | 3 | 3 | 21 | 23 |
| Conn. | 22 | 36 | 24 | 36 | 6 | 11 | 64 | 49 |
| MID. ATLANTIC | 606 | 1,565 | 498 | 977 | 265 | 162 | 1,211 | 1,220 |
| Upstate N.Y. | 329 | 444 | 64 | 161 | 19 | 6 | 167 | 147 |
| N.Y. City | 182 | 686 | 236 | 437 | 139 | 69 | 629 | 650 |
| N.J. | 40 | 293 | 133 | 247 | 54 | 36 | 276 | 295 |
| Pa. | 55 | 142 | 65 | 132 | 53 | 51 | 139 | 128 |
| E.N. CENTRAL | 1,457 | 2,366 | 654 | 700 | 506 | 711 | 668 | 720 |
| Ohio | 852 | 157 | 357 | 131 | 46 | 44 | 111 | 154 |
| Ind. | 128 | 884 | 23 | 107 | 102 | 225 | 49 | 74 |
| Ill. | 209 | 657 | 143 | 2 | 122 | 246 | 343 | 325 |
| Mich. | 155 | 469 | 116 | 424 | 219 | 163 | 130 | 117 |
| Wis. | 113 | 199 | 15 | 36 | 17 | 33 | 35 | 50 |
| W.N. CENTRAL | 813 | 1,117 | 685 | 946 | 35 | 43 | 231 | 269 |
| Minn. | 237 | 292 | 282 | 347 | 17 | 6 | 122 | 88 |
| Iowa | 254 | 269 | 215 | 207 | 1 | 10 | 18 | 23 |
| Mo. | 138 | 402 | 111 | 282 | 8 | 22 | 59 | 100 |
| N. Dak. | 13 | 4 | 7 | 4 | - | - | 3 | 2 |
| S. Dak. | 87 | 4 | 50 | 3 | - | - | 8 | 11 |
| Nebr. | 41 | 39 | - | 44 | 1 | 2 | 21 | 11 |
| Kans. | 43 | 107 | 20 | 59 | 8 | 3 | - | 34 |
| S. ATLANTIC | 1,163 | 1,369 | 344 | 515 | 1,092 | 1,108 | 1,298 | 1,522 |
| Del. | 5 | 9 | 4 | 10 | 7 | 5 | 9 | 7 |
| Md. | 62 | 79 | 33 | 43 | 126 | 163 | 111 | 144 |
| D.C. | 29 | 22 | U | U | 21 | 21 | 15 | 11 |
| Va. | 116 | 229 | 56 | 186 | 66 | 69 | 126 | 150 |
| W. Va. | 6 | 3 | 7 | 3 | - | 2 | 17 | 19 |
| N.C. | 211 | 68 | 101 | 47 | 250 | 315 | 192 | 216 |
| S.C. | 143 | 66 | 67 | 54 | 149 | 122 | 119 | 150 |
| Ga. | 124 | 126 | 57 | 108 | 169 | 206 | 235 | 318 |
| Fla. | 467 | 767 | 19 | 64 | 304 | 205 | 474 | 507 |
| E.S. CENTRAL | 760 | 522 | 335 | 315 | 346 | 496 | 414 | 507 |
| Ky. | 289 | 158 | 142 | 48 | 26 | 53 | 71 | 58 |
| Tenn. | 53 | 222 | 60 | 241 | 191 | 303 | 147 | 195 |
| Ala. | 149 | 30 | 113 | 23 | 70 | 67 | 144 | 171 |
| Miss. | 269 | 112 | 20 | 3 | 59 | 73 | 52 | 83 |
| W.S. CENTRAL | 1,011 | 1,806 | 683 | 528 | 380 | 454 | 668 | 1,123 |
| Ark. | 379 | 116 | 155 | 42 | 21 | 60 | 81 | 115 |
| La. | 108 | 168 | 106 | 97 | 74 | 112 | - | 71 |
| Okla. | 21 | 65 | 10 | 23 | 37 | 70 | 75 | 86 |
| Tex. | 503 | 1,457 | 412 | 366 | 248 | 212 | 512 | 851 |
| MOUNTAIN | 466 | 499 | 253 | 340 | 122 | 122 | 228 | 284 |
| Mont. | 1 | 5 | - | - | - | - | - | 6 |
| Idaho | 21 | 36 | - | 23 | - | 1 | 7 | 4 |
| Wyo. | 2 | 2 | - | 2 | - | 1 | 2 | 1 |
| Colo. | 93 | 88 | 80 | 46 | 23 | 5 | 66 | 41 |
| N. Mex. | 64 | 52 | 40 | 34 | 10 | 10 | 11 | 28 |
| Ariz. | 224 | 199 | 99 | 135 | 78 | 100 | 90 | 121 |
| Utah | 28 | 37 | 26 | 43 | 7 | 1 | 16 | 26 |
| Nev. | 33 | 80 | 8 | 57 | 4 | 4 | 36 | 57 |
| PACIFIC | 1,142 | 1,741 | 208 | 1,677 | 212 | 226 | 1,401 | 1,683 |
| Wash. | 97 | 324 | 119 | 291 | 32 | 36 | 124 | 141 |
| Oreg. | 40 | 104 | 61 | 65 | 4 | 8 | 52 | 50 |
| Calif. | 971 | 1,283 | - | 1,298 | 170 | 181 | 1,115 | 1,347 |
| Alaska | 4 | 6 | 1 | 3 | - | - | 26 | 66 |
| Hawaii | 30 | 24 | 27 | 20 | 6 | 1 | 84 | 79 |
| Guam | - | 26 | U | U | - | 2 | - | 32 |
| P.R. | 6 | 21 | U | U | 259 | 99 | 54 | 70 |
| V.I. | - | - | U | U | - | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | U | U |
| C.N.M.I. | 4 | U | U | U | - | U | 19 | U |

N: Not notifiable. U: Unavailable. -: No reported cases.

*Individual cases can be reported through both the National Electronic Telecommunications System for Surveillance (NETSS) and the Public Health Laboratory Information System (PHLIS).

TABLE III. Provisional cases of selected notifiable diseases preventable by vaccination, United States, weeks ending July 21, 2001, and July 22, 2000 (29th Week)

| Reporting Area | <i>H. influenzae</i> , Invasive | | Hepatitis (Viral), By Type | | | | Measles (Rubeola) | | | | | |
|----------------|---------------------------------|-----------|----------------------------|-----------|-----------|-----------|-------------------|-----------|-----------|-----------|-----------|-----------|
| | Cum. 2001 [†] | Cum. 2000 | A | | B | | Indigenous | | Imported* | | Total | |
| | | | Cum. 2001 | Cum. 2000 | Cum. 2001 | Cum. 2000 | 2001 | Cum. 2001 | 2001 | Cum. 2001 | Cum. 2001 | Cum. 2000 |
| UNITED STATES | 795 | 752 | 5,021 | 7,034 | 3,347 | 3,809 | - | 43 | - | 31 | 74 | 58 |
| NEW ENGLAND | 43 | 59 | 233 | 203 | 58 | 62 | - | 4 | - | 1 | 5 | 3 |
| Maine | 1 | 1 | 5 | 10 | 5 | 5 | - | - | - | - | - | - |
| N.H. | - | 9 | 10 | 17 | 11 | 11 | - | - | - | - | - | - |
| Vt. | 2 | 4 | 6 | 6 | 3 | 6 | - | 1 | - | - | 1 | 3 |
| Mass. | 32 | 29 | 74 | 83 | 3 | 6 | - | 2 | - | 1 | 3 | - |
| R.I. | 2 | 1 | 11 | 7 | 12 | 9 | - | - | - | - | - | - |
| Conn. | 6 | 15 | 127 | 80 | 24 | 25 | - | 1 | - | - | 1 | - |
| MID. ATLANTIC | 96 | 141 | 421 | 754 | 447 | 656 | - | 2 | - | 9 | 11 | 20 |
| Upstate N.Y. | 42 | 52 | 148 | 126 | 78 | 68 | - | 1 | - | 4 | 5 | 9 |
| N.Y. City | 25 | 39 | 171 | 271 | 261 | 314 | - | - | - | - | - | 10 |
| N.J. | 26 | 29 | 70 | 130 | 64 | 110 | - | - | - | 1 | 1 | - |
| Pa. | 3 | 21 | 32 | 227 | 44 | 164 | - | 1 | - | 4 | 5 | 1 |
| E.N. CENTRAL | 107 | 111 | 558 | 903 | 417 | 403 | - | - | - | 10 | 10 | 6 |
| Ohio | 48 | 36 | 136 | 154 | 66 | 69 | - | - | - | 3 | 3 | 2 |
| Ind. | 32 | 12 | 51 | 30 | 24 | 28 | - | - | - | 4 | 4 | - |
| Ill. | 10 | 41 | 157 | 389 | 62 | 61 | - | - | - | 3 | 3 | 3 |
| Mich. | 5 | 7 | 175 | 279 | 265 | 226 | - | - | - | - | - | 1 |
| Wis. | 12 | 15 | 39 | 51 | - | 19 | - | - | - | - | - | - |
| W.N. CENTRAL | 38 | 34 | 219 | 482 | 108 | 171 | - | 4 | - | - | 4 | 1 |
| Minn. | 21 | 16 | 16 | 129 | 13 | 20 | - | 2 | - | - | 2 | 1 |
| Iowa | - | - | 18 | 50 | 14 | 17 | - | - | - | - | - | - |
| Mo. | 11 | 11 | 59 | 213 | 53 | 90 | - | 2 | - | - | 2 | - |
| N. Dak. | 4 | 2 | 2 | 2 | - | 2 | - | - | - | - | - | - |
| S. Dak. | - | - | 1 | - | 1 | - | - | - | - | - | - | - |
| Nebr. | 1 | 3 | 27 | 21 | 14 | 26 | - | - | - | - | - | - |
| Kans. | 1 | 2 | 96 | 67 | 13 | 16 | - | - | - | - | - | - |
| S. ATLANTIC | 242 | 179 | 1,140 | 720 | 725 | 646 | - | 3 | - | 1 | 4 | 2 |
| Del. | - | - | - | 10 | - | 9 | - | - | - | - | - | - |
| Md. | 57 | 51 | 149 | 89 | 88 | 76 | - | 2 | - | 1 | 3 | - |
| D.C. | - | - | 27 | 15 | 11 | 19 | - | - | - | - | - | - |
| Va. | 18 | 29 | 68 | 85 | 85 | 83 | - | - | - | - | - | 2 |
| W. Va. | 8 | 4 | 7 | 45 | 16 | 6 | - | - | - | - | - | - |
| N.C. | 32 | 17 | 85 | 97 | 111 | 142 | - | - | - | - | - | - |
| S.C. | 5 | 7 | 38 | 30 | 16 | 5 | - | - | - | - | - | - |
| Ga. | 61 | 47 | 457 | 116 | 176 | 101 | - | 1 | - | - | 1 | - |
| Fla. | 61 | 24 | 309 | 233 | 222 | 205 | - | - | - | - | - | - |
| E.S. CENTRAL | 56 | 33 | 184 | 269 | 225 | 268 | - | 2 | - | - | 2 | - |
| Ky. | 2 | 12 | 40 | 32 | 17 | 53 | - | 2 | - | - | 2 | - |
| Tenn. | 28 | 14 | 78 | 95 | 118 | 124 | - | - | - | - | - | - |
| Ala. | 25 | 5 | 58 | 37 | 51 | 27 | - | - | - | - | - | - |
| Miss. | 1 | 2 | 8 | 105 | 39 | 64 | - | - | - | - | - | - |
| W.S. CENTRAL | 29 | 42 | 608 | 1,301 | 359 | 586 | - | 1 | - | - | 1 | - |
| Ark. | - | - | 43 | 97 | 56 | 64 | - | - | - | - | - | - |
| La. | 3 | 12 | 46 | 45 | 28 | 83 | - | - | - | - | - | - |
| Okla. | 26 | 28 | 86 | 158 | 63 | 78 | - | - | - | - | - | - |
| Tex. | - | 2 | 433 | 1,001 | 212 | 361 | - | 1 | - | - | 1 | - |
| MOUNTAIN | 110 | 76 | 476 | 485 | 335 | 280 | - | - | - | 1 | 1 | 12 |
| Mont. | - | - | 8 | 3 | 2 | 3 | - | - | - | - | - | - |
| Idaho | 1 | 3 | 48 | 18 | 8 | 4 | - | - | - | 1 | 1 | - |
| Wyo. | 13 | 1 | 21 | 4 | 29 | - | - | - | - | - | - | - |
| Colo. | 25 | 16 | 41 | 115 | 70 | 46 | - | - | - | - | - | 2 |
| N. Mex. | 14 | 16 | 18 | 44 | 81 | 89 | - | - | - | - | - | - |
| Ariz. | 42 | 31 | 250 | 233 | 105 | 99 | - | - | - | - | - | - |
| Utah | 6 | 6 | 48 | 33 | 15 | 14 | - | - | - | - | - | 3 |
| Nev. | 9 | 3 | 42 | 35 | 25 | 25 | - | - | - | - | - | 7 |
| PACIFIC | 74 | 77 | 1,182 | 1,917 | 673 | 737 | - | 27 | - | 9 | 36 | 14 |
| Wash. | 1 | 3 | 64 | 167 | 72 | 44 | - | 13 | - | 2 | 15 | 3 |
| Oreg. | 16 | 22 | 49 | 124 | 43 | 60 | - | 3 | - | - | 3 | - |
| Calif. | 32 | 29 | 1,056 | 1,604 | 541 | 619 | - | 8 | - | 4 | 12 | 8 |
| Alaska | 3 | 4 | 12 | 11 | 5 | 6 | - | - | - | - | - | 1 |
| Hawaii | 22 | 19 | 1 | 11 | 12 | 8 | - | 3 | - | 3 | 6 | 2 |
| Guam | - | 1 | - | 1 | - | 9 | U | - | U | - | - | - |
| P.R. | 1 | 3 | 56 | 173 | 99 | 156 | - | - | - | - | - | 2 |
| V.I. | - | - | - | - | - | - | U | - | U | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | U | U | - | U | 20 | U | - | - | - | - | - | U |

N: Not notifiable. U: Unavailable. -: No reported cases.

*For imported measles, cases include only those resulting from importation from other countries.

[†] Of 160 cases among children aged <5 years, serotype was reported for 72, and of those, 11 were type b.

TABLE III. (Cont'd) Provisional cases of selected notifiable diseases preventable by vaccination, United States, weeks ending July 21, 2001, and July 22, 2000 (29th Week)

| Reporting Area | Meningococcal Disease | | Mumps | | | Pertussis | | | Rubella | | |
|----------------|-----------------------|-----------|-------|-----------|-----------|-----------|-----------|-----------|---------|-----------|-----------|
| | Cum. 2001 | Cum. 2000 | 2001 | Cum. 2001 | Cum. 2000 | 2001 | Cum. 2001 | Cum. 2000 | 2001 | Cum. 2001 | Cum. 2000 |
| UNITED STATES | 1,393 | 1,405 | 1 | 109 | 206 | 31 | 2,440 | 3,168 | 1 | 16 | 95 |
| NEW ENGLAND | 79 | 85 | - | - | 3 | 1 | 256 | 860 | - | - | 11 |
| Maine | 1 | 6 | - | - | - | - | - | 14 | - | - | - |
| N.H. | 10 | 9 | - | - | - | - | 25 | 62 | - | - | 2 |
| Vt. | 4 | 2 | - | - | - | 1 | 25 | 161 | - | - | - |
| Mass. | 45 | 50 | - | - | 1 | - | 190 | 580 | - | - | 8 |
| R.I. | 2 | 6 | - | - | 1 | - | 2 | 11 | - | - | - |
| Conn. | 17 | 12 | - | - | 1 | - | 14 | 32 | - | - | 1 |
| MID. ATLANTIC | 116 | 155 | - | 10 | 13 | 1 | 201 | 259 | - | 4 | 8 |
| Upstate N.Y. | 43 | 41 | - | 1 | 5 | 1 | 108 | 138 | - | 1 | 1 |
| N.Y. City | 29 | 33 | - | 6 | 5 | - | 33 | 44 | - | 2 | 7 |
| N.J. | 34 | 27 | - | - | - | - | 8 | - | - | 1 | - |
| Pa. | 10 | 54 | - | 3 | 3 | - | 52 | 77 | - | - | - |
| E.N. CENTRAL | 169 | 244 | - | 12 | 17 | 4 | 276 | 363 | - | 3 | 1 |
| Ohio | 58 | 55 | - | 1 | 7 | - | 168 | 182 | - | - | - |
| Ind. | 27 | 31 | - | 1 | - | 3 | 27 | 36 | - | 1 | - |
| Ill. | 20 | 61 | - | 8 | 5 | 1 | 30 | 28 | - | 2 | 1 |
| Mich. | 33 | 71 | - | 2 | 4 | - | 27 | 42 | - | - | - |
| Wis. | 31 | 26 | - | - | 1 | - | 24 | 75 | - | - | - |
| W.N. CENTRAL | 101 | 91 | - | 5 | 12 | 2 | 121 | 165 | - | 2 | 1 |
| Minn. | 15 | 7 | - | 2 | - | - | 31 | 75 | - | - | - |
| Iowa | 21 | 21 | - | - | 5 | - | 16 | 26 | - | 1 | - |
| Mo. | 37 | 46 | - | - | 4 | 2 | 55 | 33 | - | - | - |
| N. Dak. | 5 | 2 | - | - | - | - | - | 1 | - | - | - |
| S. Dak. | 4 | 5 | - | - | - | - | 3 | 3 | - | - | - |
| Nebr. | 10 | 4 | - | 1 | 1 | - | 3 | 4 | - | - | 1 |
| Kans. | 9 | 6 | - | 2 | 2 | - | 13 | 23 | - | 1 | - |
| S. ATLANTIC | 272 | 201 | - | 18 | 29 | 3 | 122 | 239 | 1 | 4 | 50 |
| Del. | 2 | - | - | - | - | - | - | 6 | - | - | - |
| Md. | 32 | 19 | - | 4 | 6 | - | 17 | 65 | 1 | 1 | - |
| D.C. | - | - | - | - | - | - | 1 | 2 | - | - | - |
| Va. | 28 | 34 | - | 2 | 5 | 1 | 13 | 33 | - | - | - |
| W. Va. | 10 | 9 | - | - | - | - | 1 | 1 | - | - | - |
| N.C. | 55 | 30 | - | 1 | 4 | 1 | 41 | 51 | - | - | 42 |
| S.C. | 25 | 15 | - | 1 | 9 | 1 | 23 | 20 | - | 2 | 6 |
| Ga. | 37 | 36 | - | 7 | 2 | - | 7 | 20 | - | - | - |
| Fla. | 83 | 58 | - | 3 | 3 | - | 19 | 41 | - | 1 | 2 |
| E.S. CENTRAL | 96 | 99 | - | 3 | 4 | 2 | 56 | 66 | - | 1 | 4 |
| Ky. | 16 | 20 | - | 1 | - | - | 11 | 32 | - | - | 1 |
| Tenn. | 43 | 40 | - | - | 2 | 1 | 24 | 19 | - | 1 | - |
| Ala. | 29 | 29 | - | - | 2 | 1 | 18 | 12 | - | - | 3 |
| Miss. | 8 | 10 | - | 2 | - | - | 3 | 3 | - | - | - |
| W.S. CENTRAL | 167 | 150 | - | 8 | 22 | 3 | 200 | 145 | - | - | 6 |
| Ark. | 12 | 8 | - | 1 | 1 | - | 7 | 14 | - | - | 1 |
| La. | 54 | 35 | - | 2 | 4 | - | 2 | 9 | - | - | 1 |
| Okla. | 21 | 21 | - | - | - | - | 1 | 9 | - | - | - |
| Tex. | 80 | 86 | - | 5 | 17 | 3 | 190 | 113 | - | - | 4 |
| MOUNTAIN | 74 | 64 | - | 7 | 14 | 10 | 912 | 412 | - | 1 | 2 |
| Mont. | 3 | 4 | - | - | 1 | 3 | 13 | 12 | - | - | - |
| Idaho | 7 | 6 | - | - | - | - | 164 | 41 | - | - | - |
| Wyo. | 6 | - | - | 1 | 1 | - | 1 | 2 | - | - | - |
| Colo. | 25 | 20 | - | 1 | - | 2 | 162 | 229 | - | 1 | 1 |
| N. Mex. | 11 | 6 | - | 2 | 1 | 2 | 63 | 70 | - | - | - |
| Ariz. | 11 | 19 | - | 1 | 3 | - | 460 | 40 | - | - | 1 |
| Utah | 7 | 6 | - | 1 | 4 | 3 | 40 | 12 | - | - | - |
| Nev. | 4 | 3 | - | 1 | 4 | - | 9 | 6 | - | - | - |
| PACIFIC | 319 | 316 | 1 | 46 | 92 | 5 | 296 | 659 | - | 1 | 12 |
| Wash. | 46 | 33 | - | 1 | 3 | 3 | 82 | 202 | - | - | 7 |
| Oreg. | 24 | 38 | N | N | N | 2 | 29 | 63 | - | - | - |
| Calif. | 239 | 232 | 1 | 27 | 71 | - | 165 | 357 | - | - | 5 |
| Alaska | 2 | 5 | - | 1 | 7 | - | 2 | 12 | - | - | - |
| Hawaii | 8 | 8 | - | 17 | 11 | - | 18 | 25 | - | 1 | - |
| Guam | - | - | U | - | 11 | U | - | 3 | U | - | 1 |
| P.R. | 3 | 7 | - | - | - | - | 2 | 4 | - | - | - |
| V.I. | - | - | U | U | U | U | - | - | U | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | U | - | - | U | - | - | U | - | - | U |

N: Not notifiable.

U: Unavailable.

- : No reported cases.

TABLE IV. Deaths in 122 U.S. cities,* week ending
July 21, 2001 (29th Week)

| Reporting Area | All Causes, By Age (Years) | | | | | | P&I† Total | Reporting Area | All Causes, By Age (Years) | | | | | | P&I† Total |
|---------------------|----------------------------|-------|-------|-------|------|----|---------------|-----------------------|----------------------------|-------|-------|-------|------|-----|---------------|
| | All Ages | ≥65 | 45-64 | 25-44 | 1-24 | <1 | | | All Ages | ≥65 | 45-64 | 25-44 | 1-24 | <1 | |
| NEW ENGLAND | 541 | 375 | 96 | 45 | 15 | 10 | 58 | S. ATLANTIC | 1,329 | 840 | 310 | 105 | 46 | 26 | 96 |
| Boston, Mass. | 131 | 78 | 31 | 12 | 5 | 5 | 11 | Atlanta, Ga. | 141 | 89 | 32 | 10 | 7 | 3 | - |
| Bridgeport, Conn. | 34 | 23 | 9 | 2 | - | - | 3 | Baltimore, Md. | 197 | 104 | 60 | 21 | 6 | 6 | 22 |
| Cambridge, Mass. | 22 | 14 | 5 | 3 | - | - | 3 | Charlotte, N.C. | 103 | 62 | 25 | 8 | 3 | 5 | 7 |
| Fall River, Mass. | 33 | 25 | 5 | 3 | - | - | 2 | Jacksonville, Fla. | 143 | 97 | 34 | 7 | 4 | 1 | 18 |
| Hartford, Conn. | 74 | 50 | 14 | 6 | 3 | 1 | 9 | Miami, Fla. | 117 | 78 | 24 | 9 | 5 | 1 | 17 |
| Lowell, Mass. | 25 | 18 | 3 | 3 | 1 | - | 1 | Norfolk, Va. | 41 | 21 | 7 | 5 | 4 | 3 | 1 |
| Lynn, Mass. | 12 | 10 | 1 | 1 | - | - | 3 | Richmond, Va. | 64 | 41 | 12 | 5 | 3 | 2 | 4 |
| New Bedford, Mass. | 27 | 23 | 1 | 3 | - | - | 5 | Savannah, Ga. | 58 | 44 | 10 | 2 | 2 | - | 6 |
| New Haven, Conn. | 46 | 30 | 8 | 3 | 2 | 3 | 5 | St. Petersburg, Fla. | 54 | 41 | 8 | 2 | 2 | 1 | 5 |
| Providence, R.I. | U | U | U | U | U | U | U | Tampa, Fla. | 182 | 120 | 43 | 14 | 3 | 2 | 14 |
| Somerville, Mass. | 5 | 4 | 1 | - | - | - | 1 | Washington, D.C. | 201 | 128 | 42 | 22 | 7 | 2 | 2 |
| Springfield, Mass. | 47 | 36 | 6 | 3 | 2 | - | 6 | Wilmington, Del. | 28 | 15 | 13 | - | - | - | - |
| Waterbury, Conn. | 27 | 23 | 2 | 2 | - | - | 5 | E.S. CENTRAL | 840 | 580 | 160 | 62 | 27 | 10 | 60 |
| Worcester, Mass. | 58 | 41 | 10 | 4 | 2 | 1 | 4 | Birmingham, Ala. | 143 | 105 | 27 | 9 | 1 | - | 15 |
| MID. ATLANTIC | 2,096 | 1,483 | 413 | 125 | 40 | 35 | 107 | Chattanooga, Tenn. | 96 | 75 | 17 | 2 | 1 | 1 | 2 |
| Albany, N.Y. | 54 | 37 | 12 | 1 | 2 | 2 | 5 | Knoxville, Tenn. | 85 | 62 | 19 | 1 | 1 | 2 | 2 |
| Allentown, Pa. | 16 | 15 | 1 | - | - | - | 1 | Lexington, Ky. | 67 | 43 | 8 | 11 | 5 | - | 4 |
| Buffalo, N.Y. | 76 | 54 | 15 | 6 | 1 | 1 | 7 | Memphis, Tenn. | 161 | 105 | 27 | 13 | 13 | 3 | 16 |
| Camden, N.J. | 33 | 20 | 5 | 5 | 2 | 1 | 5 | Mobile, Ala. | 100 | 73 | 15 | 9 | 1 | 2 | 3 |
| Elizabeth, N.J. | 16 | 14 | 2 | - | - | - | - | Montgomery, Ala. | 43 | 31 | 8 | 3 | 1 | - | 7 |
| Erie, Pa.§ | 44 | 37 | 5 | 1 | - | 1 | 5 | Nashville, Tenn. | 145 | 86 | 39 | 14 | 4 | 2 | 11 |
| Jersey City, N.J. | 45 | 30 | 10 | 2 | 2 | 1 | - | W.S. CENTRAL | 1,363 | 841 | 275 | 126 | 86 | 35 | 86 |
| New York City, N.Y. | 1,116 | 766 | 239 | 72 | 20 | 19 | 50 | Austin, Tex. | 84 | 58 | 15 | 7 | 4 | - | 6 |
| Newark, N.J. | U | U | U | U | U | U | U | Baton Rouge, La. | 44 | 23 | 15 | 5 | 1 | - | 4 |
| Paterson, N.J. | 12 | 8 | 2 | - | 1 | 1 | - | Corpus Christi, Tex. | 41 | 25 | 10 | 3 | 1 | 2 | - |
| Philadelphia, Pa. | 304 | 202 | 68 | 24 | 6 | 4 | 8 | Dallas, Tex. | U | U | U | U | U | U | U |
| Pittsburgh, Pa.§ | 42 | 31 | 7 | 2 | 1 | 1 | 4 | El Paso, Tex. | 106 | 68 | 24 | 7 | 3 | 4 | 9 |
| Reading, Pa. | 29 | 24 | 3 | 1 | 1 | - | 1 | Ft. Worth, Tex. | 138 | 91 | 31 | 9 | 4 | 3 | 5 |
| Rochester, N.Y. | 113 | 90 | 15 | 4 | 3 | 1 | 8 | Houston, Tex. | 471 | 233 | 96 | 64 | 55 | 23 | 27 |
| Schenectady, N.Y. | 21 | 18 | 2 | 1 | - | - | - | Little Rock, Ark. | 62 | 33 | 12 | 11 | 4 | 2 | 3 |
| Scranton, Pa.§ | 29 | 22 | 6 | 1 | - | - | 10 | New Orleans, La. | U | U | U | U | U | U | U |
| Syracuse, N.Y. | 105 | 84 | 14 | 4 | 1 | 2 | 10 | San Antonio, Tex. | 219 | 160 | 38 | 14 | 6 | 1 | 17 |
| Trenton, N.J. | 24 | 17 | 6 | - | - | 1 | - | Shreveport, La. | 73 | 59 | 8 | 3 | 3 | - | 7 |
| Utica, N.Y. | 17 | 14 | 1 | 1 | 1 | - | 2 | Tulsa, Okla. | 125 | 91 | 26 | 3 | 5 | - | 8 |
| Yonkers, N.Y. | U | U | U | U | U | U | U | MOUNTAIN | 952 | 625 | 178 | 95 | 31 | 23 | 61 |
| E.N. CENTRAL | 1,693 | 1,154 | 345 | 107 | 50 | 37 | 123 | Albuquerque, N.M. | 67 | 46 | 8 | 11 | 2 | - | 5 |
| Akron, Ohio | 42 | 32 | 7 | 3 | - | - | 3 | Boise, Idaho | 43 | 28 | 9 | 4 | 2 | - | 4 |
| Canton, Ohio | 42 | 31 | 10 | 1 | - | - | 3 | Colo. Springs, Colo. | 62 | 48 | 7 | 6 | - | 1 | 3 |
| Chicago, Ill. | U | U | U | U | U | U | U | Denver, Colo. | 114 | 71 | 24 | 12 | 4 | 3 | 11 |
| Cincinnati, Ohio | 104 | 79 | 14 | 5 | 3 | 6 | 6 | Las Vegas, Nev. | 248 | 164 | 55 | 19 | 5 | 5 | 11 |
| Cleveland, Ohio | 119 | 65 | 39 | 9 | 2 | 4 | 5 | Ogden, Utah | 25 | 20 | 3 | 1 | 1 | - | - |
| Columbus, Ohio | 185 | 123 | 34 | 21 | 3 | 4 | 11 | Phoenix, Ariz. | 140 | 78 | 31 | 20 | 7 | 4 | 8 |
| Dayton, Ohio | 117 | 87 | 23 | 2 | 5 | - | 7 | Pueblo, Colo. | 21 | 18 | 2 | - | 1 | - | 1 |
| Detroit, Mich. | 186 | 90 | 56 | 22 | 11 | 7 | 17 | Salt Lake City, Utah | 112 | 72 | 22 | 8 | 3 | 7 | 12 |
| Evansville, Ind. | 54 | 39 | 9 | 4 | 1 | 1 | 4 | Tucson, Ariz. | 120 | 80 | 17 | 14 | 6 | 3 | 6 |
| Fort Wayne, Ind. | 64 | 52 | 10 | 1 | - | 1 | 10 | PACIFIC | 1,849 | 1,304 | 355 | 128 | 34 | 25 | 128 |
| Gary, Ind. | 17 | 8 | 4 | 3 | 2 | - | 1 | Berkeley, Calif. | 12 | 8 | 3 | - | 1 | - | 1 |
| Grand Rapids, Mich. | 99 | 64 | 11 | 5 | 12 | 7 | 9 | Fresno, Calif. | 159 | 116 | 31 | 7 | 3 | 2 | 6 |
| Indianapolis, Ind. | 171 | 112 | 38 | 14 | 4 | 3 | 10 | Glendale, Calif. | 19 | 16 | 3 | - | - | - | - |
| Lansing, Mich. | 45 | 32 | 10 | 2 | 1 | - | - | Honolulu, Hawaii | 57 | 39 | 14 | 1 | 1 | 2 | 4 |
| Milwaukee, Wis. | 115 | 85 | 23 | 5 | 1 | 1 | 16 | Long Beach, Calif. | 62 | 46 | 12 | 4 | - | - | 8 |
| Peoria, Ill. | 46 | 31 | 12 | 2 | 1 | - | 3 | Los Angeles, Calif. | 447 | 301 | 84 | 46 | 12 | 4 | 28 |
| Rockford, Ill. | 54 | 41 | 8 | 1 | 1 | 3 | 4 | Pasadena, Calif. | 33 | 24 | 6 | 2 | - | 1 | 6 |
| South Bend, Ind. | 53 | 42 | 9 | - | 2 | - | 7 | Portland, Oreg. | 148 | 113 | 23 | 6 | 1 | 5 | 4 |
| Toledo, Ohio | 101 | 78 | 15 | 7 | - | 1 | 4 | Sacramento, Calif. | 188 | 140 | 29 | 11 | 5 | 3 | 17 |
| Youngstown, Ohio | 79 | 63 | 13 | - | 1 | 2 | 3 | San Diego, Calif. | 166 | 110 | 42 | 7 | 3 | 3 | 20 |
| W.N. CENTRAL | 653 | 452 | 115 | 52 | 19 | 15 | 32 | San Francisco, Calif. | 102 | 70 | 23 | 7 | 1 | 1 | 10 |
| Des Moines, Iowa | 49 | 34 | 11 | 2 | 1 | 1 | 4 | San Jose, Calif. | 161 | 121 | 28 | 7 | 3 | 2 | 7 |
| Duluth, Minn. | 30 | 21 | 6 | 1 | 2 | - | 4 | Santa Cruz, Calif. | 28 | 19 | 4 | 5 | - | - | 3 |
| Kansas City, Kans. | 29 | 21 | 1 | 7 | - | - | 3 | Seattle, Wash. | 112 | 76 | 20 | 12 | 2 | 2 | 10 |
| Kansas City, Mo. | U | U | U | U | U | U | U | Spokane, Wash. | 49 | 36 | 7 | 5 | 1 | - | 2 |
| Lincoln, Nebr. | 41 | 32 | 8 | 1 | - | - | 3 | Tacoma, Wash. | 106 | 69 | 26 | 8 | 1 | - | 2 |
| Minneapolis, Minn. | 147 | 104 | 24 | 13 | 2 | 4 | 4 | TOTAL | 11,316† | 7,654 | 2,247 | 845 | 348 | 216 | 751 |
| Omaha, Nebr. | 80 | 57 | 13 | 6 | 2 | 2 | 4 | | | | | | | | |
| St. Louis, Mo. | 108 | 55 | 26 | 17 | 8 | 2 | - | | | | | | | | |
| St. Paul, Minn. | 81 | 62 | 11 | 3 | 1 | 4 | 6 | | | | | | | | |
| Wichita, Kans. | 88 | 66 | 15 | 2 | 3 | 2 | 4 | | | | | | | | |

U: Unavailable. --:No reported cases.

*Mortality data in this table are voluntarily reported from 122 cities in the United States, most of which have populations of ≥100,000. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included.

†Pneumonia and influenza.

‡Because of changes in reporting methods in this Pennsylvania city, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks.

§Total includes unknown ages.

Contributors to the Production of the *MMWR* (Weekly)

Weekly Notifiable Disease Morbidity Data and 122 Cities Mortality Data

Samuel L. Groseclose, D.V.M., M.P.H.

State Support Team

Robert Fagan
Jose Aponte
Gerald Jones
David Nitschke
Scott Noldy
Jim Vaughan
Carol A. Worsham

CDC Operations Team

Carol M. Knowles
Deborah A. Adams
Willie J. Anderson
Patsy A. Hall
Mechele Hester
Felicia J. Perry
Pearl Sharp

Informatics

T. Demetri Vacalis, Ph.D.

Michele D. Renshaw

Erica R. Shaver

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/Publications/mmwr>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (888) 232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

| | | |
|---------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------|
| Director, Centers for Disease Control and Prevention Jeffrey P. Koplan, M.D., M.P.H. | Director, Epidemiology Program Office Stephen B. Thacker, M.D., M.Sc. | Writers-Editors, <i>MMWR</i> (Weekly) Jill Crane David C. Johnson |
| Deputy Director for Science and Public Health, Centers for Disease Control and Prevention David W. Fleming, M.D. | Editor, <i>MMWR</i> Series John W. Ward, M.D. Acting Managing Editor, <i>MMWR</i> (Weekly) Teresa F. Rutledge | Desktop Publishing Lynda G. Cupell Morie M. Higgins |

☆U.S. Government Printing Office: 2001-633-173/48247 Region IV
