



# MMWR

## Morbidity and Mortality Weekly Report

Weekly

June 1, 2007 / Vol. 56 / No. 21


### Multistate Outbreak of *Salmonella* Serotype Tennessee Infections Associated with Peanut Butter — United States, 2006–2007

In November 2006, public health officials at CDC and state health departments detected a substantial increase in the reported incidence of isolates of *Salmonella* serotype Tennessee. In a multistate case-control study conducted during February 5–13, 2007, illness was strongly associated with consumption of either of two brands (Peter Pan or Great Value) of peanut butter produced at the same plant. Based on these findings, the plant ceased production and recalled both products on February 14, 2007. The outbreak strain of *Salmonella* Tennessee subsequently was isolated from several opened and unopened jars of Peter Pan and Great Value peanut butter and from two environmental samples obtained from the plant. New case reports decreased substantially after the product recall (Figure 1). As of May 22, 2007, a total of 628 persons infected with an outbreak strain of *Salmonella* serotype Tennessee had been reported from 47 states since August 1,

2006 (Figure 2). Local and state public health officials in multiple states, with assistance from CDC and the Food and Drug Administration (FDA), are continuing to investigate this outbreak caused by peanut butter, a new food source for salmonellosis in the United States. All remaining jars of Peter Pan or Great Value peanut butter with a product code beginning with 2111 should be discarded.


Public health officials in PulseNet (the molecular subtyping network for foodborne disease surveillance) and OutbreakNet

**FIGURE 1.** Number of confirmed cases (n = 481)\* of *Salmonella* Tennessee infection associated with consumption of peanut butter, by week of symptom onset — United States, August 1, 2006–April 23, 2007



\* Cases with outbreak-associated pulsed-field gel electrophoresis pattern and for which date of symptom onset was available.

**FIGURE 2.** Number of confirmed cases (N = 628)\* of *Salmonella* Tennessee infection associated with consumption of peanut butter, by state — United States, August 1, 2006–May 22, 2007



\*Cases with outbreak-associated pulsed-field gel electrophoresis pattern.

#### INSIDE

- 524 Sunburn Prevalence Among Adults — United States, 1999, 2003, and 2004
- 529 Characteristics and Health of Caregivers and Care Recipients — North Carolina, 2005
- 532 *Acanthamoeba* Keratitis — Multiple States, 2005–2007
- 535 QuickStats

The *MMWR* series of publications is published by the Coordinating Center for Health Information and Service, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

**Suggested Citation:** Centers for Disease Control and Prevention. [Article title]. *MMWR* 2007;56:[inclusive page numbers].

#### Centers for Disease Control and Prevention

Julie L. Gerberding, MD, MPH  
*Director*

Tanja Popovic, MD, PhD  
*Chief Science Officer*

James W. Stephens, PhD  
*(Acting) Associate Director for Science*

Steven L. Solomon, MD  
*Director, Coordinating Center for Health Information and Service*

Jay M. Bernhardt, PhD, MPH  
*Director, National Center for Health Marketing*

B. Kathleen Skipper, MA  
*(Acting) Director, Division of Health Information Dissemination (Proposed)*

#### Editorial and Production Staff

Frederic E. Shaw, MD, JD  
*Editor, MMWR Series*

Myron G. Schultz, DVM, MD  
*(Acting) Deputy Editor, MMWR Series*

Suzanne M. Hewitt, MPA  
*Managing Editor, MMWR Series*

Douglas W. Weatherwax  
*Lead Technical Writer-Editor*

Catherine H. Bricker, MS  
Jude C. Rutledge  
*Writers-Editors*

Beverly J. Holland  
*Lead Visual Information Specialist*

Lynda G. Cupell  
Malbea A. LaPete  
*Visual Information Specialists*

Quang M. Doan, MBA  
Erica R. Shaver  
*Information Technology Specialists*

#### Editorial Board

William L. Roper, MD, MPH, Chapel Hill, NC, Chairman

Virginia A. Caine, MD, Indianapolis, IN

David W. Fleming, MD, Seattle, WA

William E. Halperin, MD, DrPH, MPH, Newark, NJ

Margaret A. Hamburg, MD, Washington, DC

King K. Holmes, MD, PhD, Seattle, WA

Deborah Holtzman, PhD, Atlanta, GA

John K. Iglehart, Bethesda, MD

Dennis G. Maki, MD, Madison, WI

Sue Mallonee, MPH, Oklahoma City, OK

Stanley A. Plotkin, MD, Doylestown, PA

Patricia Quinlisk, MD, MPH, Des Moines, IA

Patrick L. Remington, MD, MPH, Madison, WI

Barbara K. Rimer, DrPH, Chapel Hill, NC

John V. Rullan, MD, MPH, San Juan, PR

Anne Schuchat, MD, Atlanta, GA

Dixie E. Snider, MD, MPH, Atlanta, GA

John W. Ward, MD, Atlanta, GA

(the network of public health epidemiologists who investigate foodborne illnesses nationwide) have been investigating this outbreak and attempting to identify the mechanism of initial contamination. The investigation began in November 2006, when public health officials in PulseNet noted a substantial increase in the number of isolates of the outbreak strain of *Salmonella* serotype Tennessee; throughout 2005 and most of 2006, these isolates were reported to PulseNet at a rate of one to five per month, whereas in October 2006, 30 isolates were reported. Pulsed-field gel electrophoresis (PFGE) patterns of *Salmonella* Tennessee strains isolated from patients were uploaded from state health department databases to CDC databases. Three closely related patterns\* were determined to be associated with this outbreak.

A case was defined as infection with *Salmonella* Tennessee with a PFGE pattern matching one of the three outbreak patterns in a person residing in the United States with symptom onset on or after August 1, 2006 (or, if onset date unknown, *Salmonella* Tennessee isolated on or after August 1, 2006). The median age of patients was 52 years (range: 2 months–95 years); 73% were female. Symptoms of infection included diarrhea (72%), abdominal cramps (65%), fever (43%), and dysuria (45%). Symptom onset dates were known for 481 of 628 patients and ranged from August 1, 2006 to April 23, 2007 (Figure 1). Twenty percent of patients were hospitalized; no deaths were attributed to *Salmonella* infection. Sixty-one percent of isolates were from stool specimens, 35% from urine specimens, and 4% from other specimens.

The initial investigation indicated that cases were not clustered geographically, and patient interviews conducted during November–December 2006 by state and local officials from OutbreakNet did not reveal a common food exposure. Officials in multiple states then interviewed 26 patients in January 2007 using a standard food-consumption survey instrument of approximately 200 items. Interviews indicated that 48% of the patients had eaten turkey (excluding delicatessen-sliced turkey) and 85% had eaten peanut butter during the week before illness onset, higher proportions than would be expected from food-consumption surveys of the U.S. population (1).

In February 2007, a case-control study with 65 patients and 124 controls was conducted to identify the food item associated with illness; the majority of interviews were completed by state and local health departments and were coordinated by CDC. For the study, a case was defined as infection with the outbreak strain of *Salmonella* Tennessee in a person aged  $\geq 18$  years with a history of diarrhea. Controls were well adults

\* CDC PulseNet patterns JNXX01.0010, JNXX01.0011, and JNXX01.0026.

from the patient's community who were matched by geographic location. Controls were identified using a reverse online telephone directory that when given an address provided telephone numbers for residences in the same extended neighborhood as the patients. The median ages for the patients and controls were 53 and 58 years, respectively. Patients were more likely than controls to have eaten peanut butter (81% versus 65%, matched odds ratio [mOR] = 1.9, 95% confidence interval [CI] = 0.8–5.2), to have eaten peanut butter more than once a week (66% versus 40%, mOR = 3.5, CI = 1.4–9.9), and to have eaten either Peter Pan or Great Value peanut butter (67% versus 13%, mOR = 10.9, CI = 3.8–43.0). Neither the consumption of other peanut butter brands nor consumption of turkey products was associated with illness.

Epidemiologic data suggesting Peter Pan brands of peanut butter as the possible source of the outbreak were provided to FDA officials on February 13, 2007. The following day, FDA issued a health alert to consumers indicating that they should not eat Peter Pan or Great Value peanut butter with a product code beginning with 2111, both of which were manufactured in a single facility in Georgia operated by ConAgra Foods. ConAgra Foods voluntarily recalled the products, destroyed existing products in their possession, and temporarily halted production pending further investigation.

New case reports decreased substantially after the February 14 recall (Figure 1). Investigation of the cases is ongoing to determine whether persons are still eating peanut butter from contaminated lots.

Subsequent laboratory testing of leftover peanut butter from patients was performed at state public health laboratories and CDC. *Salmonella* Tennessee with a PFGE pattern matching one of the outbreak strains was isolated from 21 opened and unopened peanut butter jars with production dates ranging from July 2006 to December 2006. These jars were collected from patients in 13 states (Arkansas, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Minnesota, New York, Oklahoma, Pennsylvania, South Carolina, and Tennessee); two of the PFGE strains were isolated from these peanut butter samples. FDA isolated *Salmonella* Tennessee from 13 unopened jars of Peter Pan and Great Value peanut butter with production dates ranging from August 2006 to January 2007 and from two plant environmental samples. Peanut butter from the Georgia plant was exported to 70 countries. No confirmed cases linked to this outbreak have been reported from other countries, although several possibly related cases have been investigated.

The source of the peanut butter contamination is unknown. FDA is investigating the plant operations, including heating temperatures, to determine the mechanism.

**Reported by:** *Salmonella Tennessee Outbreak Investigation Team. Local and state health departments. Div of Foodborne, Bacterial, and Mycotic Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases, CDC.*

**Editorial Note:** Approximately 2,500 *Salmonella* serotypes can cause salmonellosis, an illness characterized by diarrhea, fever, and abdominal cramps, typically 12–72 hours after infection (2). *Salmonella* Tennessee infections are rare, and the source of most of these infections is unknown. An average of 52 *Salmonella* Tennessee cases were reported to the National *Salmonella* Surveillance System<sup>†</sup> each year during 1995–2004, representing 0.1% of all reported *Salmonella* strains (3). Only one other outbreak of *Salmonella* Tennessee infection with an identified food source, contaminated powdered milk, has been reported to CDC (4). In addition to causing gastrointestinal symptoms, certain serotypes, including *Salmonella* Tennessee, are more likely than other serotypes to infect the urinary tract. The percentage of patient *Salmonella* Tennessee isolates from urine specimens increased from 15% during 1995–2004 to 27% during 2005–2006. Because urinary tract infections are more common among females, the high proportion of isolates from urine in this outbreak might explain the high percentage of identified cases among females (3,5).

This is the first reported outbreak of a foodborne illness caused by peanut butter consumption in the United States. Outside the United States, one outbreak implicating peanut butter, caused by *Salmonella* serotype Mbandaka, was reported from Australia in 1996 (6). In addition, an outbreak of *Salmonella* serotype Agona infection in four countries was associated with consumption of a peanut-butter-coated snack produced in Israel (7,8).

Peanuts can become contaminated with salmonellae during growth, harvest, or storage, and the organisms are able to survive high temperatures in a high-fat, low-water-activity environment (9). Peanut butter provides such an environment, and although it typically undergoes heat treatment to temperatures >158°F (>70°C), such heating might not always eliminate salmonellae (10). In addition, after heat treatment, peanut butter that is being processed might be contaminated by salmonellae that are introduced into the production environment on raw peanuts or another source (e.g., animals in the production plant, salmonellae brought into the plant on containers or humans from the outside environment, or other ingredients used to make peanut butter).

<sup>†</sup> The National *Salmonella* Surveillance System collects information on serotypes of *Salmonella* isolates reported through the Public Health Laboratory Information System, an electronic reporting system. Additional information is available at <http://www.cdc.gov/ncidod/dbmd/phlisdata/salmonella.htm>.

This outbreak demonstrates the potential for widespread illness from a broadly distributed contaminated product, one that has not been previously implicated in a foodborne illness outbreak in the United States. In addition, the outbreak demonstrates that processed food can become contaminated even when the production process includes a heat-treatment step, underscoring the need for effective preventive controls in food-processing plants to prevent contamination.

Certain consumers might still be eating peanut butter from contaminated lots. All remaining jars of Peter Pan and Great Value peanut butter with a product code beginning with 2111 should be discarded.

#### References

1. CDC. Foodborne diseases active surveillance network (FoodNet): population survey atlas of exposures, 2002. Atlanta, GA: CDC; 2004. Available at <http://www.cdc.gov/foodnet/surveys/pop/2002/2002atlas.pdf>.
2. Brenner FW, Villar RG, Angulo FJ, Tauxe RV, Swaminathan B. *Salmonella* nomenclature. *J Clin Microbiol* 2000;38:2465–7.
3. CDC. *Salmonella* surveillance summary, 2004. Atlanta, GA: US Department of Health and Human Services, CDC; 2006. Available at <http://www.cdc.gov/ncidod/dbmd/phlisdata/salmonella.htm>.
4. CDC. *Salmonella* serotype Tennessee in powdered milk products and infant formula—Canada and the United States, 1993. *MMWR* 1993; 42:501–19.
5. Sivapalasingam, S, Hoekstra RM, McQuiston JR, et al. *Salmonella* bacteriuria: an increasing entity in elderly women in the United States. *Epidemiol Infect* 2004;132:897–902.
6. Scheil W, Cameron S, Dalton C, Murray C, Wilson D. A South Australian *Salmonella* Mbandaka outbreak investigation using a database to select controls. *Aust N Z J Public Health* 1998;22:536–9.
7. Killalea D, Ward LR, de Roberts D, et al. International epidemiological and microbiological study of outbreak of *Salmonella agona* infection from a ready-to-eat savoury snack—I: England and Wales and the United States. *BMJ* 1996;313:1105–7.
8. Shohat T, Green MS, Marom D, et al. International epidemiological and microbiological study of outbreak of *Salmonella agona* infection from a ready-to-eat savoury snack—II: Israel. *BMJ* 1996;313:1107–9.
9. Mattick KL, Jorgensen F, Legan JD, Lappin-Scott HM, Humphrey TJ. Habituation of *Salmonella* spp. at reduced water activity and its effect on heat tolerance. *Appl Environ Microbiol* 2001;66:4921–5.
10. Shachar D, Yaron S. Heat tolerance of *Salmonella enterica* serovars Agona, Enteritidis, and Typhimurium in peanut butter. *J Food Protect* 2006;69:2687–91.

## Sunburn Prevalence Among Adults — United States, 1999, 2003, and 2004

Episodic acute overexposure to ultraviolet (UV) radiation (i.e., sunburn) is an important risk factor for two types of skin cancer: basal cell carcinoma and melanoma. Melanoma is the most lethal type of skin cancer. In 2003, a total of 45,625 new cases of melanoma were diagnosed in the United States, and 7,818 persons died from the disease (1). A meta-analysis of 57 studies indicated that the relative risk for melanoma

among persons with sunburn history compared with those without sunburn history was 2.03 (95% confidence interval [CI] = 1.73–2.37) (2). Monitoring sunburn prevalence with population-based surveys allows an estimate of compliance with sun-protection behaviors, assessments of risk for developing skin cancer, and measurement of the success of prevention programs (3). To evaluate trends in sunburn prevalence among U.S. adults, CDC analyzed cross-sectional data from the 1999, 2003, and 2004 Behavioral Risk Factor Surveillance System (BRFSS) surveys. This report describes the results of that analysis, which indicated that sunburn prevalence among all adults increased from 31.8% in 1999 to 33.7% in 2004. Further research is needed to determine which interventions will best improve sun-protection behaviors among the public.

BRFSS is a state-based, random-digit-dialed telephone survey of the noninstitutionalized, U.S. civilian population aged ≥18 years. Questions are administered in English and Spanish, as necessary. Two questions related to self-reported sunburn experiences were used in the core section (i.e., the survey questions administered in all states) of the 1999, 2003, and 2004 questionnaires. The first question was presented as follows: “The next question is about sunburns, including any time that even a small part of your skin was red for more than 12 hours. Have you had a sunburn within the past 12 months?” Persons who responded “yes” were then asked, “Including times when even a small part of your skin was red for more than 12 hours, how many sunburns have you had within the past 12 months?” Excluded from the analysis were respondents who had missing answers, refused to answer, or answered “don’t know” on the questionnaire. In 1999, a total of 156,095 persons responded, of whom 2,778 (1.8%) were excluded; in 2003, a total of 256,457 persons responded, of whom 3,478 (1.4%) were excluded; and in 2004, a total of 296,027 persons responded, of whom 906 (0.3%) were excluded. The median state response rate, based on Council of American Survey and Research Organizations guidelines, was 55.2% in 1999, 53.2% in 2003, and 52.7% in 2004. Information on each respondent’s age, race/ethnicity, and sex and other demographic data also were collected in BRFSS.

Weighted prevalence of sunburn by race/ethnicity, sex, and state was calculated for each year and for all survey years combined. Data were weighted to the sex, racial/ethnic, and age distribution of the adult population of each state using intercensal estimates. To allow comparison among survey years, stratified BRFSS data were age adjusted to the 2000 U.S. standard population using six age groups: 18–24 years, 25–34 years, 35–44 years, 45–54 years, 55–64 years, and ≥65 years. Differences in prevalence were considered statistically significant if CIs did not overlap.

Estimated sunburn prevalence among all adults ranged from 31.8% in 1999 to 33.7% in 2004 (Table 1). Men had a higher prevalence of sunburn than women in all three survey years (35.8% versus 28.0% in 1999, 37.0% versus 30.2% in 2003, and 37.0% versus 30.3% in 2004). Sunburn prevalence increased from 1999 to 2004 among non-Hispanic white women (from 35.3% to 39.6%) and non-Hispanic white men (from 44.1% to 46.9%) (Table 1). Sunburn also was reported among racial/ethnic groups traditionally considered at lower risk for sunburn or skin cancer, such as Hispanic blacks (12.4% among men and 9.5% among women in 2004), Asians/Pacific Islanders (16.2% among men and 16.1% among women in 2004), and American Indians/Alaska Natives (30.4% among men and 21.5% among women in 2004). Non-Hispanic blacks had low prevalence of sunburn (5.8% among men and women in 2004) (Table 1).

Among adults who reported sunburn during the preceding year, 20.7% reported four or more sunburns (all survey years combined). Non-Hispanics whites and American Indians/Alaska Natives had the highest proportion of respondents with four or more sunburns during the preceding year (21.2% and 19.6%, respectively) (Table 2).

In 2004, a total of 20 states\* reported a statistically significant increase in sunburn prevalence among whites, compared with 1999; four states (Indiana, Iowa, Kentucky, and Louisi-

ana) reported a significant decrease in sunburn prevalence. The lowest reported sunburn prevalence among whites during any of the three survey years was 25.7% (Arizona, 1999), and the highest was 51.3% (Utah, 2003) (Table 3 and Figure).

**Reported by:** *M Saraiya, MD, Div of Cancer Prevention and Control; L Balluz, ScD, XJ Wen, MD, Div of Adult and Community Health, National Center for Chronic Disease Prevention and Health Promotion; DA Joseph, MD, EIS Officer, CDC.*

**Editorial Note:** For all three survey years, approximately one third of the U.S. adult population had at least one sunburn during the preceding year. Of those who had at least one sunburn during the preceding year, two thirds had more than one sunburn. These findings are consistent with previous similar analyses and suggest that a substantial segment of the adult population is not consistently practicing sun-protection behaviors (3). CDC recommends the following sun-protection behaviors: wearing a wide-brimmed hat, covering up while in the sun, seeking shade, wearing wrap-around sunglasses, avoiding the sun during the hours of 10 a.m. to 4 p.m., and using sunscreen with a sun protection factor [SPF] of 15 or higher.†

Sunburn prevalence in racial/ethnic minority groups has not been well characterized previously. Skin-cancer prevention messages traditionally have been targeted toward white audiences because whites are at least 10 times more likely to

\*Alaska, Arizona, Connecticut, Florida, Idaho, Kansas, Maine, Massachusetts, Minnesota, Montana, Nebraska, New Jersey, New York, North Dakota, Oklahoma, Pennsylvania, South Carolina, Tennessee, Vermont, and Washington.

† Additional information available at [http://www.cdc.gov/cancer/skin/basic\\_info/howto.htm](http://www.cdc.gov/cancer/skin/basic_info/howto.htm).

**TABLE 1. Estimated percentage of respondents aged  $\geq 18$  years who reported having had at least one sunburn during the preceding year, by sex and race/ethnicity — Behavioral Risk Factor Surveillance System, United States, 1999, 2003, and 2004\***

Characteristic	1999		2003		2004	
	%	(95% CI) <sup>†</sup>	%	(95% CI)	%	(95% CI)
<b>Men</b>						
White, non-Hispanic	44.1	(43.5–44.8)	46.5 <sup>§</sup>	(45.9–47.1)	46.9 <sup>§</sup>	(46.4–47.5)
White, Hispanic	21.6	(19.4–23.8)	24.8	(22.0–27.7)	20.1	(18.8–23.3)
Black, non-Hispanic	5.3	(4.3–6.4)	4.6	(3.7–5.6)	5.8	(4.7–7.0)
Black, Hispanic	14.3	(10.1–19.8)	12.3	(8.4–17.8)	12.4	(8.4–17.5)
Asian/Pacific Islander	18.0	(14.5–22.1)	18.6	(15.5–22.1)	16.2	(13.6–19.2)
American Indian/Alaska Native	27.4	(23.0–32.4)	25.9	(22.1–30.0)	30.4	(26.3–34.8)
<b>Total</b>	<b>35.8</b>	<b>(35.2–36.3)</b>	<b>37.0<sup>§</sup></b>	<b>(36.5–37.6)</b>	<b>37.0<sup>§</sup></b>	<b>(36.4–37.6)</b>
<b>Women</b>						
White, non-Hispanic	35.3	(34.8–35.8)	38.7 <sup>§</sup>	(38.3–39.2)	39.6 <sup>§</sup>	(39.1–40.0)
White, Hispanic	17.2	(15.6–19.0)	19.7	(17.9–21.6)	17.2	(15.8–18.8)
Black, non-Hispanic	5.1	(4.5–5.9)	5.7	(5.1–6.4)	5.8	(5.2–6.5)
Black, Hispanic	8.3	(5.7–12.0)	13.5	(8.7–20.3)	9.5	(6.7–13.2)
Asian/Pacific Islander	11.0	(8.9–13.6)	14.4	(12.3–16.8)	16.1	(12.9–19.9)
American Indian/Alaska Native	23.5	(19.7–27.8)	23.3	(19.9–27.0)	21.5	(18.6–24.7)
<b>Total</b>	<b>28.0</b>	<b>(27.5–28.4)</b>	<b>30.3<sup>§</sup></b>	<b>(29.9–30.7)</b>	<b>30.3<sup>§</sup></b>	<b>(29.9–30.7)</b>
<b>Total</b>	<b>31.8</b>	<b>(31.4–32.2)</b>	<b>33.6<sup>§</sup></b>	<b>(33.2–33.9)</b>	<b>33.7<sup>§</sup></b>	<b>(33.4–34.1)</b>

\* Age adjusted to the 2000 U.S. standard population.

† Confidence interval.

§ Statistically significant difference compared with 1999. Differences were considered statistically significant if CIs did not overlap.

**TABLE 2. Estimated percentage of respondents aged  $\geq 18$  years who reported having had at least one sunburn during the preceding year, by number of sunburns and race/ethnicity — Behavioral Risk Factor Surveillance System, United States, 1999, 2003, and 2004 (all survey years combined)\***

Race/Ethnicity <sup>†</sup>	No. of sunburns							
	1		2		3		$\geq 4$	
	%	(95% CI) <sup>§</sup>	%	(95% CI)	%	(95% CI)	%	(95% CI)
White, non-Hispanic	35.6	(35.2–36.2)	28.3	(27.9–28.7)	14.9	(14.6–15.2)	21.2	(20.8–21.5)
White, Hispanic	45.6	(42.2–49.1)	23.4	(20.8–26.1)	11.9	(10.3–13.7)	19.1	(17.0–21.5)
Black, non-Hispanic	50.4	(46.4–54.4)	25.5	(21.5–28.0)	11.8	(9.7–14.3)	12.3	(10.3–14.6)
Asian/Pacific Islander	53.3	(48.1–58.5)	22.1	(18.6–25.9)	9.1	(7.2–11.4)	15.5	(12.0–19.8)
American Indian/Alaska Native	44.2	(39.8–48.6)	22.5	(19.2–26.1)	13.8	(10.9–17.4)	19.6	(16.7–22.9)
<b>Total</b>	<b>36.9</b>	<b>(36.5–37.4)</b>	<b>27.8</b>	<b>(27.4–28.1)</b>	<b>14.6</b>	<b>(14.3–14.8)</b>	<b>20.7</b>	<b>(20.4–21.1)</b>

\* Age adjusted to the 2000 U.S. standard population.

<sup>†</sup> Hispanic blacks excluded from analysis because of small sample size.

<sup>§</sup> Confidence interval.

develop melanoma than racial/ethnic minorities, although racial/ethnic minority populations are more likely to have more advanced disease diagnosed and to have lower 5-year survival rates (4,5). However, the findings in this report indicate that substantial portions of the Hispanic, Asian/Pacific Islander, and American Indian/Alaska Native populations acquired sunburns. Racial/ethnic groups with darker skin (higher melanin content) have a lower incidence of skin cancer, which is attributed, in part, to the inherent sun protection provided by melanin (5). However, race/ethnicity is a poor proxy for skin cancer risk because persons in racial/ethnic minority groups might have individual risk factors for skin cancer (e.g., lighter skin color; skin that burns, freckles, or reddens easily in the sun; or personal or family history of skin cancer) and might not benefit from the protective effects of melanin. In a study of adolescents aged 11–18 years, 21.7% of black respondents who reported having had a sunburn reported severe sunburn with blisters or peeling after 1 hour of exposure to sun during the summer (6).

The results of this analysis also indicated that men had a higher prevalence of sunburn than women in most of the racial/ethnic groups surveyed. This finding might be attributed to different sun-protection behaviors or different sun-exposure conditions between men and women, (e.g., differences in leisure or work activities). In addition, women might be more concerned about the cosmetic effect of long-term sun exposure (e.g., wrinkling of the skin and the appearance of age spots) and thus might be more likely to avoid sun exposure, use makeup with sunscreen, or practice sun-protection behaviors (7). In 2003, white men had a higher melanoma incidence and mortality than white women (22.7 versus 15.1 and 4.4 versus 2.0 per 100,000 population, respectively) (1).

None of the states with sunburn prevalence among whites greater than 45% were traditional “sunbelt” states. Persons living in the northern states might use fewer precautions dur-

ing the first sunny days after winter or might travel to other locations where they acquire sunburns (3). Previous analyses have demonstrated that states with lower UV radiation (i.e., those in higher latitudes) have had more rapid increases in melanoma incidence than states with higher UV radiation (8).

The findings in this report are subject to at least six limitations. First, the BRFSS survey is a telephone survey, and results obtained might not be generalizable to U.S. adults without landline telephones. Second, responses are self-reported and therefore subject to recall bias. Third, the BRFSS survey does not contain questions regarding skin type or sun-protection behaviors, so this information cannot be correlated to sunburns. Fourth, this analysis was a cross-sectional study, and individual sunburn patterns could not be followed over time. Fifth, the source of UV exposure (sun or artificial source) was not provided. Finally, the U.S. states in which respondents actually acquired their sunburns were unknown.

Sunburn prevalence among U.S. adults increased from 1999 to 2004. Several reasons might account for these increases. For example, the public might be receiving conflicting or confusing messages about what constitutes the best sun-protection behaviors. A review of 20 Internet sites about skin cancer prevention revealed inconsistent advice regarding a safe amount of sun exposure, times of day to avoid the sun, how many sunburns increase the risk for skin cancer, and the best types of clothing to use for sun protection (9). That review noted that only three recommendations were common to all 20 Internet sites: wearing broad-brimmed hats, wearing sunglasses, and using sunscreen with an SPF of 15 or higher (9). In addition, certain segments of the public might view the purported benefits of sun exposure (e.g., tanned skin or elevated mood) as outweighing the risk for skin cancer or might not be concerned about the risks of overexposure to the sun (9).

**TABLE 3. Estimated percentage of white\* respondents aged ≥18 years who reported having had at least one sunburn during the preceding year, by state/area — Behavioral Risk Factor Surveillance System, United States, 1999, 2003, and 2004†**

State/Area	1999		2003		2004	
	%	(95% CI)§	%	(95% CI)	%	(95% CI)
Alabama	39.6	(37.0–42.1)	39.7	(37.5–41.9)	39.6	(37.4–41.8)
Alaska	27.8	(24.7–30.9)	32.6	(29.6–35.5)	34.1¶	(31.1–37.0)
Arizona	25.7	(22.1–29.2)	39.5¶	(36.5–42.5)	42.1¶	(40.0–45.2)
Arkansas	42.0	(40.0–44.0)	41.8	(40.0–43.6)	42.9	(41.1–44.8)
California	34.1	(32.5–35.8)	36.8	(35.0–38.6)	34.8	(33.0–36.6)
Colorado	45.8	(43.4–48.1)	46.6	(44.9–48.4)	45.1	(43.3–46.8)
Connecticut	33.3	(30.1–35.6)	40.7¶	(39.1–42.3)	43.1¶	(41.4–44.7)
Delaware	43.6	(40.8–46.5)	33.5¶	(31.3–35.7)	41.4	(39.1–43.6)
District of Columbia	42.8	(37.5–48.1)	41.3	(38.1–44.5)	40.1	(37.4–42.8)
Florida	33.4	(31.7–35.1)	36.8	(34.4–39.2)	37.7¶	(35.7–39.6)
Georgia	36.3	(33.8–38.8)	40.7¶	(38.9–42.5)	39.2	(37.1–41.3)
Hawaii	44.8	(39.8–49.8)	42.3	(39.6–44.9)	—**	—**
Idaho	45.4	(43.8–46.9)	49.3¶	(47.7–50.9)	48.5¶	(47.0–50.1)
Illinois	44.0	(40.1–47.1)	40.8	(38.5–43.1)	41.7	(39.8–43.7)
Indiana	48.1	(44.8–51.4)	44.9	(43.4–46.4)	43.3¶	(41.8–44.6)
Iowa	49.0	(47.1–50.8)	43.5¶	(41.5–45.1)	43.6¶	(42.0–45.2)
Kansas	34.2	(32.5–35.9)	41.3¶	(39.6–43.0)	41.4¶	(40.2–42.6)
Kentucky	30.7	(29.2–32.2)	28.0	(26.3–29.7)	27.0¶	(25.2–28.8)
Louisiana	35.2	(32.3–38.1)	31.0	(29.2–32.7)	30.5¶	(29.1–31.9)
Maine	37.0	(34.3–39.5)	42.8¶	(40.6–44.9)	42.6¶	(40.6–44.5)
Maryland	41.4	(39.2–43.6)	41.4	(39.4–43.4)	43.9	(41.8–46.0)
Massachusetts	35.2	(33.4–36.9)	41.0¶	(39.5–42.6)	42.6¶	(41.1–44.2)
Michigan	47.6	(45.5–49.8)	47.9	(45.9–49.8)	45.6	(43.9–47.3)
Minnesota	40.0	(38.6–41.4)	49.2¶	(47.5–50.9)	48.7¶	(46.9–50.4)
Mississippi	39.9	(37.4–42.4)	42.4	(40.4–44.3)	40.5	(38.6–42.4)
Missouri	42.9	(40.8–45.0)	45.9	(43.7–48.1)	45.2	(43.2–47.2)
Montana	38.8	(36.3–41.2)	47.6¶	(45.4–48.1)	44.1¶	(42.2–45.9)
Nebraska	43.1	(41.1–45.1)	46.3	(44.7–47.8)	46.9¶	(45.5–48.3)
Nevada	40.9	(37.8–44.1)	39.2	(36.6–41.2)	38.3	(35.4–41.4)
New Hampshire	41.7	(38.8–44.6)	42.8	(41.2–44.4)	43.8	(42.2–45.5)
New Jersey	32.8	(30.5–35.1)	39.0¶	(37.8–40.3)	40.2¶	(38.9–41.5)
New Mexico	38.6	(36.6–40.7)	38.8	(37.0–40.6)	41.3	(39.6–43.0)
New York	30.3	(28.1–32.5)	39.7¶	(38.0–41.4)	40.2¶	(38.5–42.0)
North Carolina	30.9	(28.5–33.4)	28.1	(26.4–29.8)	28.1	(27.0–29.3)
North Dakota	38.3	(36.0–40.6)	45.6¶	(43.6–47.6)	46.4¶	(44.5–48.3)
Ohio	39.1	(36.1–42.1)	42.4	(40.3–44.5)	43.4	(41.0–45.9)
Oklahoma	30.2	(28.2–32.3)	41.9¶	(40.4–43.3)	41.5¶	(40.0–43.0)
Oregon	41.3	(38.7–43.8)	42.1	(40.3–43.9)	43.6	(41.9–45.2)
Pennsylvania	36.5	(34.8–38.4)	43.8¶	(41.9–45.7)	42.7¶	(41.1–44.2)
Rhode Island	35.0	(33.4–36.8)	38.5	(36.6–40.5)	38.7	(36.7–40.7)
South Carolina	32.7	(30.7–34.7)	42.8¶	(41.2–44.4)	41.6¶	(40.1–43.2)
South Dakota	45.9	(44.2–47.5)	47.2	(45.7–48.8)	46.1	(44.6–47.7)
Tennessee	26.6	(24.7–28.5)	30.7	(28.4–33.0)	32.6¶	(30.4–34.7)
Texas	36.6	(34.8–38.3)	38.1	(36.6–39.7)	37.7	(36.0–39.3)
Utah	46.1	(43.9–48.3)	51.3¶	(49.3–53.1)	49.9	(48.3–51.5)
Vermont	39.1	(37.3–40.9)	45.7¶	(44.0–47.4)	47.1¶	(45.8–48.5)
Virginia	40.5	(37.8–43.2)	41.5	(39.5–43.4)	42.9	(40.8–44.9)
Washington	39.6	(37.6–41.6)	38.3	(37.3–39.2)	43.6¶	(42.7–44.6)
West Virginia	34.8	(32.8–36.8)	41.5¶	(39.5–43.4)	38.0	(36.1–40.0)
Wisconsin	51.4	(49.1–53.7)	49.3	(47.5–51.2)	48.6	(46.9–50.4)
Wyoming	48.4	(46.4–50.5)	49.5	(47.8–51.2)	48.3	(46.6–50.0)
Guam	—**	—**	50.6	(41.3–59.8)	—**	—**
Puerto Rico	12.8	(11.2–14.7)	11.8	(10.0–13.6)	14.2	(12.4–16.0)
U.S. Virgin Islands	—**	—**	46.5	(38.1–54.8)	50.1	(45.0–55.3)
<b>United States</b>	<b>37.0</b>	<b>(36.5–37.4)</b>	<b>40.0¶</b>	<b>(39.6–40.4)</b>	<b>39.9¶</b>	<b>(39.5–40.3)</b>

\* Includes Hispanic whites.


† Age adjusted to the 2000 U.S. standard population.

§ Confidence interval.

¶ Statistically significant difference compared with 1999. Differences were considered statistically significant if CIs did not overlap.

\*\* Did not participate in survey.

**FIGURE. Estimated percentage of white\* respondents aged  $\geq 18$  years who reported having had at least one sunburn during the preceding year, by state/area — Behavioral Risk Factor Surveillance System, 1999, 2003, and 2004†**



\* Includes Hispanic whites.

† Age adjusted to the 2000 U.S. standard population.

§ Puerto Rico.

¶ U.S. Virgin Islands.

Further research is needed to determine which public health interventions will improve sun-protection behaviors. The *Guide to Community Preventive Services* review of interventions to prevent skin cancer found sufficient evidence to warrant recommending educational and policy interventions for children in primary schools and for adults in recreational and tourism settings. However, evidence to warrant recommending other interventions, such as mass media campaigns and interventions in secondary schools, was insufficient (10). Future research on interventions to improve sun-protection behaviors should assess 1) the incorporation of messages that emphasize that sun-protection behaviors are readily implemented, and 2) the standardization of messages to eliminate confusing or conflicting advice. Public health messages specific to various racial/ethnic groups, certain subgroups (e.g., persons with lighter skin and men), and other populations at high risk (e.g., those with a personal or family history of skin cancer) need to be developed to emphasize that these populations also are susceptible to the harmful effects of the sun. Greater knowledge of interventions that will increase the proportion of persons who practice sun-protection behaviors is important for reducing sunburn prevalence and ultimately reducing skin cancer incidence and mortality. Continued surveillance of sunburn prevalence is necessary for evaluating the effectiveness of both current and new interventions.

#### References

1. US Cancer Statistics Working Group. United States cancer statistics: 1999–2003 incidence and mortality web based report. Atlanta, GA: US Department of Health and Human Services, CDC, National Cancer Institute; 2006. Available at <http://www.cdc.gov/uscs>.
2. Gandini S, Sera F, Cattaruzza MS, et al. Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure. *Eur J Cancer* 2005;41:45–60.
3. Saraiya M, Hall I, Uhler RJ. Sunburn prevalence among adults in the United States, 1999. *Am J Prev Med* 2002;23:91–7.
4. Cockburn MG, Zadnick J, Deapen D. Developing epidemic of melanoma in the Hispanic population of California. *Cancer* 2006;106:1162–8.
5. Gloster HM, Neal K. Skin cancer in skin of color. *J Am Acad Dermatol* 2006;55:741–60.
6. Davis KJ, Cokkinides VE, Weinstock MA, O'Connell MC, Wingo PA. Summer sunburn and sun exposure among US youths ages 11 to 18: national prevalence and associated factors. *Pediatrics* 2002;110:27–35.
7. Abroms L, Jorgensen CM, Southwell BG, Geller AC, Emmons KM. Gender differences in young adults' beliefs about sunscreen use. *Health Educ Behav* 2003;30:29–43.
8. Jemal A, Devesa SS, Fears TR, Hartge P. Cancer surveillance series: changing patterns of cutaneous malignant melanoma mortality rates among whites in the United States. *J Natl Cancer Inst* 2000;92:811–8.
9. Hillhouse J, Turrissi R. Skin cancer risk behaviors: a conceptual framework for complex behavioral change. *Arch Dermatol* 2005;141:1028–31.
10. CDC. Guide to community preventive services. Atlanta, GA: US Department of Health and Human Services, CDC; 2003. Available at [www.thecommunityguide.org/cancer](http://www.thecommunityguide.org/cancer).



## Characteristics and Health of Caregivers and Care Recipients — North Carolina, 2005

Approximately 53.4 million caregivers in the United States provide an estimated \$257–\$389 billion worth of unpaid care annually to persons of all ages with disabilities and chronic illness (1,2). The health of caregivers and their ability to continue their contributions have emerged as public health concerns (3). A 2004 study indicated that those persons who provided the most intense caregiving reported substantially poorer health than noncaregivers or those with modest caregiving responsibilities (2). A *Healthy People 2010* objective calls for public health surveillance and health promotion programs for persons with disabilities and caregivers in every state and the District of Columbia (objective 6-13) (4). Although limited caregiver surveys have provided data at the national level, data have not been available at the state level to characterize the health of caregivers or health effects of caregiving. Such information could be useful to states for planning and policy decisions and the development and implementation of interventions to promote caregivers' health. To analyze the characteristics and health of caregivers and care recipients and to assess the effects of caregiving, data were analyzed from a caregiver module that was piloted in North Carolina in the 2005 Behavioral Risk Factor Surveillance System (BRFSS) survey. This report summarizes the results of that analysis, which determined that caregivers provided an average of 20.1 hours of care per week, and 72.2% of caregivers lived in the same household as (24.9%) or within 20 minutes of (47.3%) the care recipient. Caregivers were more likely to be women (59.5%) than men and averaged more days when their mental health was not good when compared with noncaregivers (4.3 days versus 3.0 days, of the preceding 30 days). Public health initiatives should be designed to promote the health and well-being of both care recipients and caregivers.

BRFSS is a state-based, random-digit-dialed telephone survey of the noninstitutionalized, U.S. civilian population aged  $\geq 18$  years. Questions related to caregiving were administered as part of the national BRFSS core survey (i.e., the survey questions administered in all states) in 2000 (5) but were not repeated. To improve caregiver surveillance, CDC provided funds to the University of Florida in 2004 to develop a caregiver module to collect data on the characteristics and health of caregivers. Module development was based on previous research (2) and influenced by key national stakeholders convened by AARP (formerly known as the American Association of Retired Persons) in February 2005. North Carolina was selected as the site for piloting the module because of the state's large BRFSS sample size and administrative capacity.

The following caregiver screening question was administered in the North Carolina 2005 BRFSS survey: "People may provide regular care or assistance to someone who has a long-term illness or disability. During the past month, did you provide any such care or assistance to a family member or friend?" Persons who responded "yes" to the question were classified as caregivers and completed the caregiver module by responding to questions about themselves and their primary care recipient. If the caregiver had more than one care recipient, the caregiver was asked to answer module questions in reference to the recipient who required the most care. Questions in the caregiver module related to the age and sex of the care recipient; relationship of the care recipient to the caregiver (e.g., parent, spouse, or child); care recipient's major diagnosis (e.g., heart disease, cancer, stroke, or diabetes); care recipient's functional limitations (e.g., moving around or self-care) as defined by the *International Classification of Functioning, Disability, and Health* (6); duration of caregiving; hours per week of caregiving; difficulties for the caregiver that were created by caregiving (e.g., stress, lack of time, and health problems); and travel time to reach the care recipient. Comparisons were tested using *t* tests for differences in means of continuous variables and chi-square or chi-square for trend tests among categorical variables.

Among the 5,859 survey respondents interviewed during May–August 2005, a total of 895 (weighted prevalence: 15.5%; 95% confidence interval [CI] = 14.2%–16.9%) indicated they were caregivers. A greater percentage of caregivers (59.5%) were women than men (40.5%) (Table 1). A greater percentage of caregivers (21.2%) than noncaregivers (15.8%) were non-Hispanic blacks, but a smaller percentage of caregivers (2.3%) than noncaregivers (10.3%) were Hispanic (Table 1). On average, caregivers reported more days (4.3 days out of 30 days) that their mental health was not good than noncaregivers (3.0 days), although the number of days that physical health was reported not good was similar for caregivers (3.2 days) and noncaregivers (3.5 days).

Most care recipients (67.2%) were female and older than the general population; 64.3% of care recipients were aged  $\geq 65$  years, and 82.8% were cared for by a relative (Table 2). The major diagnoses of care recipients specified by caregivers were heart disease (12.8%), cancer (11.7%), stroke (9.1%), diabetes (9.0%), dementia (8.8%), arthritis/rheumatism (5.1%), lung disease/emphysema (3.0%), cerebral palsy (2.6%), and hypertension (2.4%). When asked to identify the functional limitations of their care recipients that required the most help, caregivers named moving around (41.7%); self-care (e.g., eating, dressing, bathing, and toileting) (41.0%); learning, memory, and confusion (17.0%); and anxiety or depression (16.4%) (Table 2). On average, caregivers had pro-

**TABLE 1. Characteristics of caregivers compared with noncaregivers\* — Behavioral Risk Factor Surveillance System, North Carolina, 2005**

Characteristic	Caregiver (n = 895)		Noncaregiver (n = 4,964)	
	%	(95% CI†)	%	(95% CI)
<b>Age group (yrs)</b>				
18–34	26.9	(21.8–32.8)	32.1	(29.7–33.9)
35–44	19.3	(16.2–22.9)	21.0	(19.6–22.6)
45–54	21.4	(18.2–24.9)	17.0	(15.7–18.4)
55–64	17.4	(14.7–20.6)	13.1	(12.4–14.5)
≥65	15.0	(12.5–17.8)	16.7	(15.6–17.9)
<b>Race/Ethnicity</b>				
White, non-Hispanic	71.5	(66.1–76.3)	69.6	(68.4–72.2)
Black, non-Hispanic§	21.2	(16.6–26.6)	15.8	(13.8–16.4)
Other, non-Hispanic	5.0	(3.1–8.0)	4.2	(3.5–5.0)
Hispanic§	2.3	(1.4–3.9)	10.3	(8.9–12.2)
<b>Sex</b>				
Men	40.5	(35.6–45.7)	48.8	(45.8–50.7)
Women§	59.5	(54.3–64.4)	51.1	(49.3–53.2)
<b>Marital status</b>				
Married/Coupled	64.8	(59.6–69.7)	63.9	(62.4–66.2)
Divorced/Separated	11.2	(9.2–13.6)	11.7	(10.6–12.6)
Widowed	5.1	(3.8–6.9)	7.1	(6.4–7.8)
Never married	18.8	(14.0–24.8)	17.3	(15.3–19.0)
<b>Education¶</b>				
No formal/Eighth grade or less	2.9	(1.6–5.2)	7.4	(6.2–8.7)
Some high school	9.9	(6.2–15.5)	10.4	(9.1–11.6)
High school graduate	28.3	(24.4–32.6)	29.3	(27.4–31.0)
Beyond high school	58.8	(53.8–63.7)	52.9	(51.2–55.1)
<b>Annual income</b>				
<\$25,000	30.3	(26.0–35.0)	34.0	(31.8–36.0)
\$25,000–\$34,999	17.4	(14.0–21.4)	14.0	(12.6–15.5)
\$35,000–\$49,999	16.3	(11.8–22.1)	15.0	(13.8–16.5)
\$50,000–\$74,999	15.9	(13.0–19.4)	16.1	(14.8–17.7)
≥\$75,000	20.1	(16.7–24.0)	21.0	(19.2–22.5)

\* Percentages are weighted according to state population estimates; groupings do not all add to 100.0% because of rounding.

† Confidence interval.

§ Significant difference between caregivers and noncaregivers by chi-square test;  $p < 0.05$ .

¶ Significant difference between caregivers and noncaregivers across categories of education (rather than between education levels);  $p < 0.05$ .

vided care for 42.5 months, with 26.4% providing care for >5 years (Table 3). Although caregivers averaged 20.1 hours per week of care, 13.6% provided ≥40 hours per week. When asked to name the one or two greatest difficulties they experienced from caregiving, 29.9% of caregivers cited stress, 27.9% cited not enough time for themselves or their families, and 12.0% indicated that caregiving had created a financial burden (Table 3). In addition, 3.5% of caregivers said caregiving created or aggravated health problems. In response to a separate question, 3.7% (CI = 2.5%–5.4%) reported sustaining an injury while caregiving. Nearly half (47.3%) of caregivers lived within 20 minutes of the care recipient; 24.9% resided in the same household (Table 3).

**TABLE 2. Characteristics of care recipients\* — Behavioral Risk Factor Surveillance System, North Carolina, 2005**

Characteristic	Care recipients (n = 895)	
	%	(95% CI†)
<b>Age group (yrs)</b>		
0–5	1.6	(0.7–3.8)
6–17	5.5	(3.6–8.3)
18–29	3.1	(1.8–5.3)
30–49	10.5	(7.9–13.9)
50–64	15.1	(12.4–18.2)
65–74	19.2	(15.9–22.9)
75–84	25.5	(22.0–29.3)
≥85	19.6	(15.2–24.8)
<b>Sex</b>		
Male	32.8	(28.7–37.3)
Female	67.2	(62.7–71.3)
<b>Relationship to caregiver</b>		
Relative other than spouse	72.0	(66.9–76.6)
Nonrelative	16.0	(11.8–21.4)
Spouse	10.8	(8.6–13.5)
Paid caregiver	1.2	(0.6–2.4)
<b>Major diagnosis</b>		
Heart disease	12.8	(10.3–15.8)
Cancer	11.7	(9.3–14.6)
Stroke	9.1	(6.8–12.0)
Diabetes	9.0	(6.5–12.4)
Dementia	8.8	(6.5–11.7)
Arthritis/Rheumatism	5.1	(3.6–7.3)
Lung disease/Emphysema	3.0	(1.8–4.9)
Cerebral palsy	2.6	(1.2–5.3)
Hypertension	2.4	(1.3–4.5)
Other disease or condition	35.5	(31.4–39.9)

\* Percentages are weighted according to state population estimates; groupings do not all add to 100.0% because of rounding.

† Confidence interval.

**Reported by:** B Neugaard, PhD, EM Andresen, PhD, EL DeFries, MPH, Univ of Florida. RC Talley, PhD, JE Crews, DPA, Div of Human Development and Disability, National Center on Birth Defects and Developmental Disabilities, CDC.

**Editorial Note:** These findings from the piloting of the BRFSS caregiver module reveal that an estimated 15.5% of adults in North Carolina were caregivers in 2005, compared with an estimated 21% of adults nationally, according to a study published in 2004 (2). In North Carolina, caregivers were more likely to be women than men, and one fourth of caregivers had been providing care for >5 years. Care recipients were most likely to be aged ≥65 years, and more than half had a major diagnosis of a chronic disabling condition, such as heart disease, cancer, stroke, diabetes, and dementia.

Recent data on the prevalence of disability indicated that variations occur from state to state (from 11% to 26% of the population) and persons with disabilities are likely to report overall poorer health (9). The extent to which the prevalence of caregiving mirrors these variations in disability is unknown and represents an area for future research.

**TABLE 3. Characteristics of caregivers\* — Behavioral Risk Factor Surveillance System, North Carolina, 2005**

Characteristic	Caregiver (n = 895)	
	%	(95% CI†)
<b>Duration of caregiving</b>		
≤3 mos	22.1	(18.5–26.3)
4–12 mos	22.0	(18.6–25.8)
13–24 mos	17.6	(13.0–23.5)
25 mos–5 yrs	11.8	(9.4–14.8)
>5 yrs	26.4	(22.7–30.4)
<b>Amount of caregiving (hrs per week)</b>		
≤8	52.1	(47.1–57.2)
9–19	18.1	(14.8–22.0)
20–39	16.1	(13.0–19.8)
≥40	13.6	(10.9–16.9)
<b>Greatest difficulties resulting from caregiving§</b>		
Creates stress	29.9	(26.1–34.0)
Not enough time for self or family	27.9	(22.4–34.7)
Financial burden	12.0	(9.7–14.9)
Interferes with work	6.9	(4.9–9.7)
Affects family relationships	5.6	(4.1–7.6)
Causes or aggravates health problems	3.5	(2.4–5.0)
Creates other difficulties	4.0	(2.5–6.2)
<b>Functional limitations for which caregiver provides the most help¶</b>		
Moving around	41.7	(37.2–46.3)
Self-care (e.g., eating, dressing, bathing, and toileting)	41.0	(36.2–45.9)
Learning, remembering, and confusion	17.0	(14.0–20.4)
Feeling anxious or depressed	16.4	(13.6–19.6)
Communicating with others	8.7	(6.8–11.1)
Seeing or hearing	7.0	(5.2–9.4)
Getting along with others	6.1	(4.4–8.5)
<b>Travel time to care recipient</b>		
Same house	24.9	(21.2–29.0)
<20 min	47.3	(42.5–52.1)
20–59 min	18.4	(13.9–23.9)
1–2 hrs	2.9	(1.9–4.3)
>2 hrs	6.6	(4.9–8.9)

\* Percentages are weighted according to state population estimates; groupings do not all add to 100.0% because of rounding.

† Confidence interval.

§ Respondents were asked to name one or two.

¶ Respondents were asked to name the top one or two activity limitations of the care recipient as defined by the World Health Organization's *International Classification of Functioning, Disability, and Health*. Available at <http://www.who.int/classifications/icf/en> (6).

The intensity and duration of caregiving has the potential to affect overall health among caregivers (7,8). Programs should be developed to address both the physical and mental health needs of caregivers and to relieve some of the difficulties they report (e.g., not enough time for self or family and financial burdens). Because 29.9% of caregivers indicate that stress, which can precipitate long-term physical or mental health problems, is one of the greatest difficulties resulting from caregiving, policies and preventive support should remove or mediate stressors. Also, given that 3.7% of caregivers report

injuries, training and materials should be considered to prevent these injuries. In addition, because BRFSS data are collected only from adults aged ≥18 years, they do not reflect younger caregivers; alternative forms of national and state assessment should be developed to analyze the health needs and caregiving patterns for populations aged <18 years.

The caregiver module should be implemented in additional states to create national and state profiles of caregiving and caregiver health effects. By adopting the caregiver module, state officials, including policy makers and program planners in services for aging populations, children and youths, and persons with disabilities, will be able to develop and monitor data-driven state plans to support caregivers and care recipients.

The findings in this report are subject to at least five limitations. First, BRFSS is a telephone-based survey and excludes households without landline telephones or with cellular telephones only. Second, data are self-reported and subject to recall bias; therefore, prevalence estimates might be underestimated or overestimated. Third, no question specifically asked whether caregivers were paid or unpaid; the 1.2% of caregivers recorded as paid represents only those who provided that information without being asked, and therefore likely underestimate the proportion of caregivers who were paid. Further research might determine whether differences exist in the characteristics of paid and unpaid caregivers and the implications of these differences. Fourth, assessment of the greatest difficulties resulting from caregiving was asked only of caregivers, and no comparison can be made between caregivers and noncaregivers with regard to these difficulties (e.g., stress and not enough time for self or family). Finally, information on the relationship between the caregiver and the care recipient was limited (i.e., spouse, relative other than spouse, non-relative, and paid). Future surveys will more fully describe this relationship (e.g., parent, sibling, or child).

The data presented in this report are the first CDC-sponsored state-level data to assess the characteristics and health of caregivers and care recipients and their caregiving situations. CDC plans to support further testing to develop an optional BRFSS caregiver module for adoption by more states in 2009. Participation by all states would enable CDC to report caregiver health status, using population-based data, for the first time nationally and by state. Additionally, states would have the surveillance results needed to create benchmarks, document and prioritize caregiver needs among their residents, and plan interventions to address those needs.

#### Acknowledgments

The report is based, in part, on contributions by the North Carolina BRFSS team and AARP.

## References

1. Arno PS, Levine C, Memmott MM. The economic value of informal caregiving. *Health Aff (Millwood)* 1999;18:182–8.
2. National Alliance for Caregiving and AARP. Caregiving in the U.S. Washington, DC: National Alliance for Caregiving and AARP; 2004. Available at <http://www.caregiving.org/data/04finalreport.pdf>.
3. Talley RC, Crews JE. Framing the public health of caregiving. *Am J Public Health* 2007;97:224–8.
4. US Department of Health and Human Services. Healthy people 2010 (conference ed, in 2 vols). Washington, DC: US Department of Health and Human Services; 2000. Available at <http://www.health.gov/healthy-people>.
5. Behavioral Risk Factor Surveillance System. 2000 Behavioral Risk Factor Surveillance System questionnaire. Available at [http://www.cdc.gov/brfss/technical\\_infodata/surveydata/2000/qcoremod\\_00.rtf](http://www.cdc.gov/brfss/technical_infodata/surveydata/2000/qcoremod_00.rtf).
6. World Health Organization. International classification of functioning, disability and health. Geneva, Switzerland: World Health Organization; 2001. Available at <http://www.who.int/classifications/icf/en>.
7. Schulz R, Beach SR. Caregiving as a risk factor for mortality: the Caregiver Health Effects Study. *JAMA* 1999;282:2215–9.
8. Ekwall AK, Sivberg B, Hallberg IR. Older caregivers' coping strategies and sense of coherence in relation to quality of life. *J Adv Nurs* 2007;57:584–96.
9. CDC. Disability and health state chartbook 2006: profiles of health for adults with disabilities. Atlanta, GA: US Department of Health and Human Services, CDC; 2006. Available at <http://www.cdc.gov/ncbddd/dh/chartbook>.

## **Acanthamoeba Keratitis — Multiple States, 2005–2007**

On May 26, 2007, this report was posted as an MMWR Dispatch on the MMWR website (<http://www.cdc.gov/mmwr>).

In May 2006, the Illinois Department of Public Health (IDPH) informed CDC about a possible increase in *Acanthamoeba* keratitis (AK) at an ophthalmology center in Illinois during the preceding 3 years. The University of Illinois at Chicago (UIC) was investigating this possible increase. In October 2006, IDPH updated CDC about the ongoing UIC investigation. At that time, CDC informally contacted multiple ophthalmology centers in the United States to assess whether the potential increase in cases extended beyond Illinois. Responses from the ophthalmology centers were inconclusive. In January 2007, CDC initiated a retrospective survey of 22 ophthalmology centers nationwide to assess whether cases were increasing throughout the United States. In March 2007, data received from 13 centers demonstrated an increase in culture-confirmed cases of AK with wide geographic distribution. The increase in cases had begun in 2004 and continued to the present. On March 16, 2007, CDC initiated a multistate investigation to look for risk factors associated with this increase in AK cases. This report summarizes recent preliminary results of that investigation, which, indicated an association with AK in soft contact lens wearers who used

Advanced Medical Optics (Santa Ana, California) Complete<sup>®</sup> MoisturePlus<sup>™</sup> (AMOCMP) multipurpose cleaning solution. CDC and the Food and Drug Administration (FDA) are taking steps to notify the public and the medical and public health communities of this preliminary association. The manufacturer has undertaken a voluntary recall of the product.

AK, a rare but potentially blinding infection of the cornea, is caused by a ubiquitous, free-living amoeba (*Acanthamoeba*) that is found commonly in the environment, including water (e.g., tap and recreational water), soil, sewage systems, cooling towers, and heating/ventilation/air conditioning (HVAC) systems. AK primarily affects otherwise healthy persons who wear contact lenses; an estimated 85% of U.S. cases occur in contact lens wearers (including wearers who follow recommended contact lens-care practices) (1). Persons who improperly store, handle, or disinfect their lenses (e.g., by using tap water or homemade solutions for cleaning); swim, use hot tubs, or shower while wearing lenses; come in contact with contaminated water; have minor damage to their corneas; or have previous corneal trauma are at increased risk for infection (2). Based on an analysis of cases reported to CDC during 1985–1987, the incidence of AK in the United States has been estimated at one to two cases per million contact lens users (3,4). An estimated 30 million persons in the United States wear soft contact lenses (5).

Initial case finding for this investigation was facilitated through postings on the *Epidemic Information Exchange (Epi-X)*, on ophthalmology/optometry/infection control listservs and websites, and through queries of clinical microbiology laboratories. As of May 24, 2007, a total of 138 patients with onset of symptoms on or after January 1, 2005, and positive *Acanthamoeba* cultures from corneal specimens had been reported to CDC by public health authorities and ophthalmologists from 35 states and Puerto Rico. Standardized telephone interviews of patients, ophthalmologists, and primary eye-care providers are being conducted by state and local health officials and CDC. Laboratory testing of clinical specimens, contact lenses, bottles of solution, and contact lens cases received from AK patients, including typing of *Acanthamoeba* spp. isolates, is ongoing. An initial analysis was conducted using data from the first 46 completed patient interviews.

Among the 46 culture-confirmed patients who were interviewed, the median age was 40 years (range: 15–77 years); six (13%) were aged <18 years. Twenty-seven (59%) were female. Of the 37 of these patients for whom clinical data were available, medical therapy was unsuccessful for nine (24%), and they were required or expected to undergo corneal transplantation. Of the 46 patients, 39 (85%) wore soft contact lenses, three (7%) wore rigid lenses, and four (9%) reported no con-

tact lens use. Among the 42 contact lens users, 16 (38%) reported swimming while wearing contact lenses and 35 (83%) reported showering while wearing contact lenses during the month before symptom onset.

Among the 39 soft contact lens users, 36 reported using one or more specific types of contact lens solution, 21 of these (58%) reported any use of AMOCMP in the month before symptom onset, 20 (56%) reported using AMOCMP as their primary solution, and 14 (39%) reported using AMOCMP as their exclusive solution. Exposure data from the 36 patients who wore soft contact lenses and used any type of contact lens solution were compared with exposure data from controls who were interviewed as part of the 2006 CDC *Fusarium* keratitis outbreak investigation (6). These controls, who were selected as geographically matched controls for the *Fusarium* keratitis cases, represented a sample of adult soft contact lens wearers from different U.S. states who were asked about product use and behaviors during March 2006 (6).

The 14 AK soft contact lens-wearing case-patients with symptom onset dates before April 1, 2006 (the period most comparable to *Fusarium* controls), who reported use of a single solution were compared with 115 controls from the *Fusarium* investigation who reported using a single solution. The results indicated that four (29%) of the 14 AK case-patients had used AMOCMP, compared with six (5%) of the 115 *Fusarium* controls (odds ratio: 7.3 [95% confidence interval (CI) = 1.7–30.1]). In a separate comparison, 36 soft contact lens-wearing AK case-patients with symptom onset dates before May 24, 2007, who reported use of one or more solutions were compared with 124 *Fusarium* controls who reported using one or more solutions. The results indicated that 21 (58%) of the 36 AK case-patients had used AMOCMP, compared with eight (6%) of the 124 *Fusarium* controls (odds ratio: 20.3; [CI = 7.6–53.9]). AMOCMP lot numbers were available for 10 patients who reported using the solution; no single lot number was repeated, suggesting that AMOCMP was not intrinsically contaminated. Analysis of the reported use of other brands of contact lens solution did not reveal any statistically significant associations.

The AK investigation by CDC, state and local health departments, FDA, and other partners, is continuing, and interviews of the remaining patients with culture-confirmed AK, their treating ophthalmologists, and their primary eye-care providers are ongoing. Although the results of initial analyses are preliminary, they suggest that use of AMOCMP increases the risk for AK. Additional studies will provide a more definitive assessment of the risk associated with use of AMOCMP. However, based on the preliminary findings, persons who wear soft contact lenses and who use AMOCMP should 1) stop using the product immediately and discard all

remaining solution, including partially used or unopened bottles; 2) choose an alternative contact lens solution; 3) discard current lens storage container; 4) discard their current pair of soft lenses; 5) see a health-care provider if they experience any signs of eye infection, including eye pain or redness, blurred vision, sensitivity to light, sensation of something in the eye, or excessive tearing.

Contact lens users with questions regarding which solutions are best for them should consult their eye-care provider. Patients should also consult their eye-care provider if they have any of the following symptoms: eye pain or redness, blurred vision, sensitivity to light, sensation of something in the eye, and/or excessive tearing. AK symptoms, which can last several weeks to months, vary among patients. Early in the infection, symptoms can be similar to the symptoms of other more common eye infections; however, AK can result in vision loss or blindness if untreated.

All contact lens wearers should follow established guidelines to help reduce the risk for eye infections, including AK (Box). Primary-care clinicians evaluating contact lens users with symptoms of eye pain or redness, tearing, decreased visual acuity, discharge, sensitivity to light, or foreign body sensation should consider the diagnosis of AK and refer

#### **BOX. Guidelines for contact lens users to help reduce their risk for eye infections**

- Visit your eye-care provider for regular eye examinations.
- Wear and replace contact lenses according to the schedule prescribed by your eye-care provider.
- Remove contact lenses before any activity involving contact with water, including showering, using a hot tub, or swimming.
- Wash hands with soap and water and dry before handling contact lenses.
- Clean contact lenses according to the manufacturer's guidelines and instructions from your eye-care provider.
  - Use fresh cleaning or disinfecting solution each time lenses are cleaned and stored. Never reuse or top off old solution.
  - Never use saline solution and rewetting drops to disinfect lenses. Neither solution is an effective or approved disinfectant.
- Store reusable lenses in the proper storage case.
  - Rinse storage cases with sterile contact lens solution (never use tap water) and leave open to dry after each use.
  - Replace storage cases at least once every 3 months.

patients to an ophthalmologist, if appropriate. Diagnosis of AK requires a high degree of suspicion, especially in a contact lens wearer with a recent diagnosis of another form of keratitis, such as herpes simplex virus keratitis, who is not responding to therapy. Diagnosis of AK is based on clinical presentation and isolation of organisms from corneal culture or detection of trophozoites and/or cysts on histopathology. However, a negative culture does not necessarily rule out *Acanthamoeba* infection. Confocal microscopy and polymerase chain reaction assays to detect *Acanthamoeba* can also assist with diagnosis. Early diagnosis can greatly improve treatment efficacy.

Clinicians should consider obtaining clinical specimens (e.g., corneal scrapings) for culture before initiating treatment. Clinicians or microbiology laboratories should report cases of AK to state and local health departments or directly to CDC at telephone, 770-488-7775. *Acanthamoeba* isolates should be submitted to state laboratories according to instructions provided by local and state public health laboratories. Public inquiries should be made via telephone 800-CDC-INFO. Further information regarding *Acanthamoeba* infections is available at <http://www.cdc.gov/ncidod/dpd/parasites/acanthamoeba/index.htm>

**Reported by:** K Bryant, J Bugante Los Angeles County Health Dept; T Chang, DVM, S Chen, MPH, J Rosenberg, MD, California Dept of Health Svcs. R Hammond, PhD, K McConnell, MPH, R Sanderson, MA, Florida Dept of Health. J Elm, MS, M Nakata, C Wakida, Hawaii Dept of Health. C Austin, DVM, J Bestudik, MG Bordson, C Conover, MD, Illinois Dept of Public Health. L Granzow, MPH, Indiana Dept of Health. A Pelletier, MD, V Rea, MPH, Maine Dept of Health and Human Svcs. A Chu, MHS, E Luckman, MPH, Maryland Dept of Health and Mental Hygiene. K Signs, DVM, Michigan Dept of Community Health. J Harper, MS, Minnesota Dept of Health. T Damrow, PhD, E Mosher, Montana Dept of Public Health and Human Svcs. K Kruger, North Dakota Dept of Health. E Saheli, MPH, Ohio Dept of Health. M Cassidy, J Hatch, Oregon Public Health Div, Dept Human Svcs. A Weltman, MD, Pennsylvania Dept of Health.

EJ Garcia Rivera, MD, Y Garcia, MPH, Puerto Rico Dept of Health. MA Kainer, MD, Tennessee Dept of Health. J Archer, MS, Wisconsin Dept of Health and Family Svcs. C Joslin, OD, Univ of Illinois Chicago. P Cernoch, Methodist Hospital of Houston; D Jones MD, M Hamill MD, A Matoba MD, S Pflugfelder MD, K Wilhelmus, MD, Baylor College of Medicine, Texas. S Beavers, MD, T Chen, MD, K Christian, DVM, M Cooper, MD, D Dufficy, DVM, M Gershman, MD, M Glenshaw, PhD, A Hall, DVM, S Holzbauer, DVM, A Huang, MD, A Langer, DVM, Z Moore, MD, AS Patel, PhD, LR Carpenter, DVM, J Schaffzin, MD, J Su, MD, I Trevino, DVM, T Weiser, MD, P Wiersma, MD, S Lorick, DO, JR Verani, MD, EIS officers, CDC.

### Acknowledgments

The findings in this report are based, in part, on contributions by M Bonhomme, N Pressly, M Robboy, OD, J Saviola, OD, E Woo, Food and Drug Admin. MJ Beach, PhD, C Braden, MD, S Brim, MA, D Chang, MD, F Chow, A daSilva, PhD, AJ Deokar, MPH, R Greco Kone, MPH, S Johnston, MS, AS Kusano, MS, B Park, Y Qvarnstrom, PhD, MD, S Persaud, S Roy, MD, G Visvesvara PhD, D Wagner, K Wannemuehler, MS, JS Yoder, MPH, National Center for Zoonotic, Vector-Borne, and Enteric Diseases, CDC.


### References

1. Stehr-Green JK, Bailey TM, Brandt FH, Carr JH, Bond WW, Visvesvara GS. *Acanthamoeba* keratitis in soft contact lens wearers: a case-control study. *JAMA* 1987;258:57-60.
2. Parmar DN, Awwad ST, Petroll WM, Bowman RW, McCulley JP, Cavanagh HD. Tandem scanning confocal corneal microscopy in the diagnosis of suspected *Acanthamoeba* keratitis. *Ophthalmology* 2006;113:538-47.
3. Schaumberg DA, Snow KK, Dana MR. The epidemic of *Acanthamoeba* keratitis: where do we stand? *Cornea* 1998;17:3-10.
4. Stehr-Green JK, Bailey TM, Visvesvara GS. The epidemiology of *Acanthamoeba* keratitis in the United States. *Am J Ophthalmol* 1989;107:331-6.
5. US Environmental Protection Agency. Do you wear contact lenses? There's something you should know. Available at <http://www.epa.gov/waterscience/acanthamoeba>.
6. Chang DC, Grant GB, O'Donnell K, et al. Multistate outbreak of *Fusarium* keratitis associated with use of a contact lens solution. *JAMA* 2006;296:953-63.

# QuickStats

FROM THE NATIONAL CENTER FOR HEALTH STATISTICS

## Estimated Percentage\* of Persons Who Delayed or Did Not Receive Medical Care During the Preceding Year Because of Cost, by Respondent-Assessed Health Status† — National Health Interview Survey, United States, 2005



\* Estimates are age adjusted using the 2000 projected U.S. population as the standard population and using five age groups: 0–11 years, 12–17 years, 18–44 years, 45–64 years, and  $\geq 65$  years. Estimates are based on household interviews of a sample of the civilian, noninstitutionalized U.S. population.

† Based on responses to the following questions: “During the past 12 months, has [person] delayed seeking medical care because of worry about the cost?” and “During the past 12 months was there any time when [person] needed medical care but did not get it because [person] could not afford it?” Both questions exclude dental care. Respondents were asked to answer regarding themselves and other family members living in the same household. Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor.

§ 95% confidence interval.

In 2005, approximately 7% of persons (21.7 million) delayed medical care during the preceding year because of worry about the cost, and another 5% (15.2 million) did not receive needed medical care because they could not afford it. Persons whose health was assessed as fair or poor were four to five times as likely as persons whose health was assessed as excellent or very good to delay or not receive needed medical care because of cost.

**SOURCE:** Adams PF, Dey, AN, Vickerie JL. Summary health statistics for the U.S. population: National Health Interview Survey, 2005. Vital Health Stat 2007;10(233). Available at [http://www.cdc.gov/nchs/data/series/sr\\_10/sr10\\_233.pdf](http://www.cdc.gov/nchs/data/series/sr_10/sr10_233.pdf).

**TABLE I. Provisional cases of infrequently reported notifiable diseases (<1,000 cases reported during the preceding year) — United States, week ending May 26, 2007 (21st Week)\***

Disease	Current week	Cum 2007	5-year weekly average†	Total cases reported for previous years					States reporting cases during current week (No.)
				2006	2005	2004	2003	2002	
Anthrax	—	—	—	1	—	—	—	2	
Botulism:									
foodborne	—	2	0	20	19	16	20	28	
infant	1	28	2	98	85	87	76	69	PA (1)
other (wound & unspecified)	2	7	0	47	31	30	33	21	WA (1), CA (1)
Brucellosis	1	47	2	118	120	114	104	125	CA (1)
Chancroid	—	10	1	33	17	30	54	67	
Cholera	—	—	0	8	8	5	2	2	
Cyclosporiasis§	1	24	17	136	543	171	75	156	MI (1)
Diphtheria	—	—	—	—	—	—	1	1	
Domestic arboviral diseases§¶:									
California serogroup	—	—	0	63	80	112	108	164	
eastern equine	—	—	0	7	21	6	14	10	
Powassan	—	—	—	1	1	1	—	1	
St. Louis	—	—	0	9	13	12	41	28	
western equine	—	—	—	—	—	—	—	—	
Ehrlichiosis§:									
human granulocytic	4	25	9	689	786	537	362	511	MN (4)
human monocytic	2	53	5	556	506	338	321	216	NC (2)
human (other & unspecified)	—	19	2	238	112	59	44	23	
<i>Haemophilus influenzae</i> **,									
invasive disease (age <5 yrs):									
serotype b	—	5	1	24	9	19	32	34	
nonserotype b	—	33	2	140	135	135	117	144	
unknown serotype	5	106	4	218	217	177	227	153	GA (2), CO (1), UT (1), AK (1)
Hansen disease§	—	19	2	67	87	105	95	96	
Hantavirus pulmonary syndrome§	—	6	1	37	26	24	26	19	
Hemolytic uremic syndrome, postdiarrheal§	1	37	4	284	221	200	178	216	CA (1)
Hepatitis C viral, acute	4	248	20	806	652	713	1,102	1,835	MN (1), UT (1), WA (1), CA (1)
HIV infection, pediatric (age <13 yrs)††	—	—	4	52	380	436	504	420	
Influenza-associated pediatric mortality§§§	1	61	0	41	45	—	N	N	MN (1)
Listeriosis	4	187	11	858	896	753	696	665	OH (1), KS (1), NC (1), CA (1)
Measles¶¶	1	13	2	73	66	37	56	44	FL (1)
Meningococcal disease, invasive***:									
A, C, Y, & W-135	3	107	5	272	297	—	—	—	CT (1), OK (1), CO (1)
serogroup B	—	41	3	170	156	—	—	—	
other serogroup	—	9	0	28	27	—	—	—	
unknown serogroup	13	299	15	682	765	—	—	—	PA (1), OH (1), MN (1), UT (1), OR (1), CA (8)
Mumps	7	384	59	6,585	314	258	231	270	OH (1), MN (1), KS (1), NC (1), GA (1), FL (1), OK (1)
Novel influenza A virus infections	—	—	—	N	N	N	N	N	
Plague	—	—	0	17	8	3	1	2	
Poliomyelitis, paralytic	—	—	—	—	1	—	—	—	
Poliovirus infection, nonparalytic§	—	—	—	N	N	N	N	N	
Psittacosis§	—	3	0	19	16	12	12	18	
Q fever§	1	66	3	176	136	70	71	61	MO (1)
Rabies, human	—	—	—	3	2	7	2	3	
Rubella†††	—	8	0	12	11	10	7	18	
Rubella, congenital syndrome	—	—	—	1	1	—	1	1	
SARS-CoV§§§§	—	—	0	—	—	—	8	N	
Smallpox§	—	—	—	—	—	—	—	—	
Streptococcal toxic-shock syndrome§	2	34	3	125	129	132	161	118	OH (1), KY (1)
Syphilis, congenital (age <1 yr)	3	74	8	379	329	353	413	412	NC (2), WA (1)
Tetanus	1	4	1	38	27	34	20	25	MO (1)
Toxic-shock syndrome (staphylococcal)§	—	30	2	98	90	95	133	109	
Trichinellosis	—	1	0	13	16	5	6	14	
Tularemia	2	6	3	98	154	134	129	90	MO (1), OK (1)
Typhoid fever	1	101	5	333	324	322	356	321	CA (1)
Vancomycin-intermediate <i>Staphylococcus aureus</i> §	—	3	—	6	2	—	N	N	
Vancomycin-resistant <i>Staphylococcus aureus</i> §	—	—	0	1	3	1	N	N	
Vibriosis (non-cholera <i>Vibrio</i> species infections)§	2	65	0	N	N	N	N	N	FL (1), CA (1)
Yellow fever	—	—	—	—	—	—	—	1	

—: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts.

\* Incidence data for reporting years 2006 and 2007 are provisional, whereas data for 2002, 2003, 2004, and 2005 are finalized.

† Calculated by summing the incidence counts for the current week, the 2 weeks preceding the current week, and the 2 weeks following the current week, for a total of 5 preceding years. Additional information is available at <http://www.cdc.gov/epo/dphsi/phs/files/5yearweeklyaverage.pdf>.

§ Not notifiable in all states. Data from states where the condition is not notifiable are excluded from this table, except in 2007 for the domestic arboviral diseases and influenza-associated pediatric mortality, and in 2003 for SARS-CoV. Reporting exceptions are available at <http://www.cdc.gov/epo/dphsi/phs/infdis.htm>.

¶ Includes both neuroinvasive and non-neuroinvasive. Updated weekly from reports to the Division of Vector-Borne Infectious Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases (ArboNET Surveillance). Data for West Nile virus are available in Table II.

\*\* Data for *H. influenzae* (all ages, all serotypes) are available in Table II.

†† Updated monthly from reports to the Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention. Implementation of HIV reporting influences the number of cases reported. Updates of pediatric HIV data have been temporarily suspended until upgrading of the national HIV/AIDS surveillance data management system is completed. Data for HIV/AIDS, when available, are displayed in Table IV, which appears quarterly.

§§ Updated weekly from reports to the Influenza Division, National Center for Immunization and Respiratory Diseases. A total of 62 cases were reported for the 2006–07 flu season.

¶¶ The one measles case reported for the current week was indigenous.

\*\*\* Data for meningococcal disease (all serogroups) are available in Table II.

††† No rubella cases were reported for the current week.

§§§ Updated weekly from reports to the Division of Viral and Rickettsial Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases.



TABLE II. Provisional cases of selected notifiable diseases, United States, weeks ending May 26, 2007, and May 27, 2006 (21st Week)\*

Reporting area	Chlamydia†					Coccidioidomycosis					Cryptosporidiosis				
	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006
		Med	Max				Med	Max				Med	Max		
<b>United States</b>	8,574	19,843	25,557	368,532	405,901	108	153	658	3,228	3,520	32	71	321	966	1,068
<b>New England</b>	523	665	1,357	13,254	12,525	—	0	0	—	—	—	5	38	61	98
Connecticut	—	204	829	3,286	3,084	N	0	0	N	N	—	0	11	11	38
Maine§	—	47	73	970	862	—	0	0	—	—	—	0	6	9	13
Massachusetts	387	302	600	6,514	5,997	—	0	0	—	—	—	2	29	18	38
New Hampshire	33	38	69	756	733	—	0	0	—	—	—	1	5	10	6
Rhode Island§	72	64	108	1,378	1,341	—	0	0	—	—	—	0	5	5	1
Vermont§	31	20	45	350	508	N	0	0	N	N	—	1	4	8	2
<b>Mid. Atlantic</b>	483	2,571	4,271	54,213	49,713	—	0	0	—	—	1	10	33	118	175
New Jersey	—	377	541	5,132	7,708	N	0	0	N	N	—	0	1	—	9
New York (Upstate)	—	509	2,745	9,903	9,141	N	0	0	N	N	—	3	13	43	36
New York City	—	753	1,523	16,763	16,982	N	0	0	N	N	—	2	10	22	53
Pennsylvania	483	832	1,776	22,415	15,882	N	0	0	N	N	1	3	18	53	77
<b>E.N. Central</b>	1,435	3,190	6,223	67,350	69,328	2	1	3	13	16	6	15	110	219	238
Illinois	655	982	1,290	18,565	22,281	—	0	0	—	—	—	2	22	19	31
Indiana	—	385	644	8,059	8,329	—	0	0	—	—	—	1	18	16	19
Michigan	603	749	1,225	14,983	12,629	1	1	3	10	12	1	3	10	54	37
Ohio	75	643	3,647	18,435	17,297	1	0	2	3	4	5	5	33	74	79
Wisconsin	102	372	528	7,308	8,792	N	0	0	N	N	—	4	53	56	72
<b>W.N. Central</b>	57	1,188	1,445	18,686	24,870	—	0	54	3	—	6	11	77	146	159
Iowa	—	160	238	3,106	3,428	N	0	0	N	N	—	2	28	25	15
Kansas	—	148	316	3,178	3,342	N	0	0	N	N	2	1	8	23	21
Minnesota	—	243	314	3,744	5,272	—	0	54	—	—	4	2	25	38	60
Missouri	—	434	628	5,220	8,997	—	0	1	3	—	—	2	21	29	33
Nebraska§	—	104	185	1,991	2,010	N	0	0	N	N	—	1	16	6	12
North Dakota	—	28	64	446	757	N	0	0	N	N	—	0	1	1	1
South Dakota	57	49	84	1,001	1,064	N	0	0	N	N	—	1	7	24	17
<b>S. Atlantic</b>	1,399	3,566	7,072	56,080	77,421	—	0	1	1	2	13	18	71	246	241
Delaware	24	69	111	1,354	1,442	N	0	0	N	N	—	0	3	2	—
District of Columbia	97	79	161	2,219	1,221	—	0	0	—	—	—	0	2	3	7
Florida	—	921	1,187	3,300	19,068	N	0	0	N	N	12	8	32	127	98
Georgia	—	675	3,822	7,608	13,694	N	0	0	N	N	—	4	18	45	70
Maryland§	—	392	669	6,594	8,201	—	0	1	1	2	—	0	2	11	6
North Carolina	479	634	1,207	12,643	14,575	—	0	0	—	—	1	1	11	25	29
South Carolina§	353	425	2,105	11,106	8,620	N	0	0	N	N	—	1	14	14	12
Virginia§	432	495	685	10,187	9,393	N	0	0	N	N	—	1	5	16	17
West Virginia	14	55	85	1,069	1,207	N	0	0	N	N	—	0	3	3	2
<b>E.S. Central</b>	765	1,409	2,044	25,713	30,823	—	0	0	—	—	—	3	14	47	40
Alabama§	70	368	539	2,079	9,833	N	0	0	N	N	—	0	11	17	14
Kentucky	174	130	691	3,118	3,810	N	0	0	N	N	—	1	3	15	10
Mississippi	—	422	959	8,916	6,998	N	0	0	N	N	—	0	8	8	5
Tennessee§	521	531	700	11,600	10,182	N	0	0	N	N	—	1	5	7	11
<b>W.S. Central</b>	1,627	2,158	3,028	43,866	45,844	—	0	1	—	—	3	5	45	36	46
Arkansas§	171	161	337	3,324	3,230	N	0	0	N	N	—	0	3	2	6
Louisiana	—	315	610	5,225	6,888	—	0	1	—	—	—	1	9	14	—
Oklahoma	260	257	472	5,190	4,749	N	0	0	N	N	3	0	9	15	12
Texas§	1,196	1,457	1,911	30,127	30,977	N	0	0	N	N	—	2	36	5	28
<b>Mountain</b>	198	1,334	2,025	20,832	26,307	48	100	293	2,173	2,496	3	4	40	66	41
Arizona	8	463	993	6,497	7,989	48	99	293	2,125	2,426	—	0	5	13	4
Colorado	112	306	416	3,730	6,408	N	0	0	N	N	1	1	7	21	9
Idaho§	—	44	253	1,263	1,329	N	0	0	N	N	—	0	5	4	4
Montana§	—	52	144	945	901	N	0	0	N	N	—	0	26	4	6
Nevada§	67	167	397	3,439	3,027	—	1	3	17	31	1	0	3	4	3
New Mexico§	—	166	324	2,591	4,048	—	0	3	7	8	—	1	6	11	9
Utah	—	97	200	1,886	2,010	—	1	4	24	29	1	0	3	2	6
Wyoming§	11	27	45	481	595	—	0	0	—	2	—	0	11	7	—
<b>Pacific</b>	2,087	3,362	4,362	68,538	69,070	58	53	311	1,038	1,006	—	1	5	27	30
Alaska	72	88	157	1,730	1,672	N	0	0	N	N	—	0	1	—	1
California	1,565	2,656	3,627	53,759	53,745	58	53	311	1,038	1,006	—	0	0	—	—
Hawaii	—	105	130	1,994	2,337	N	0	0	N	N	—	0	1	—	—
Oregon§	133	161	394	3,818	3,915	N	0	0	N	N	—	1	5	27	29
Washington	317	344	621	7,237	7,401	N	0	0	N	N	—	0	0	—	—
American Samoa	U	0	21	U	U	U	0	0	U	U	U	0	0	U	U
C.N.M.I.	U	—	—	U	U	U	—	—	U	U	U	—	—	U	U
Guam	—	—	—	—	44	—	—	—	—	—	—	—	—	—	—
Puerto Rico	229	122	234	3,041	1,999	N	0	0	N	N	N	0	0	N	N
U.S. Virgin Islands	U	3	10	U	U	U	0	0	U	U	U	0	0	U	U

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

\* Incidence data for reporting years 2006 and 2007 are provisional. Data for HIV/AIDS, AIDS, and TB, when available, are displayed in Table IV, which appears quarterly.

† Chlamydia refers to genital infections caused by *Chlamydia trachomatis*.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

**TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending May 26, 2007, and May 27, 2006 (21st Week)\***

Reporting area	Giardiasis					Gonorrhea					Haemophilus influenzae, invasive All ages, all serotypes†				
	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006
		Med	Max				Med	Max				Med	Max		
<b>United States</b>	122	310	1,477	5,088	6,068	2,958	6,865	8,969	116,556	137,450	22	47	174	948	972
<b>New England</b>	—	26	67	359	563	73	109	259	2,098	2,147	—	3	18	67	82
Connecticut	—	5	25	86	94	—	43	204	689	779	—	0	6	20	17
Maine <sup>§</sup>	—	4	14	50	33	—	2	8	41	50	—	0	4	6	7
Massachusetts	—	12	39	157	343	61	46	96	1,090	1,002	—	2	8	36	49
New Hampshire	—	0	9	4	2	4	2	8	62	95	—	0	3	4	2
Rhode Island <sup>§</sup>	—	0	17	22	35	6	10	19	196	198	—	0	10	1	2
Vermont <sup>§</sup>	—	3	12	40	56	2	1	5	20	23	—	0	1	—	5
<b>Mid. Atlantic</b>	10	63	120	876	1,193	144	683	1,537	14,256	12,996	2	10	26	200	200
New Jersey	—	7	17	36	180	—	104	155	1,483	2,138	—	1	5	17	37
New York (Upstate)	—	25	101	332	375	—	119	1,035	2,299	2,401	—	3	14	57	53
New York City	2	16	32	282	380	—	177	376	3,624	4,005	—	2	6	41	38
Pennsylvania	8	14	34	226	258	144	249	608	6,850	4,452	2	3	10	85	72
<b>E.N. Central</b>	20	44	100	709	976	542	1,297	2,581	26,709	27,637	4	7	15	101	170
Illinois	—	11	30	103	234	243	352	485	6,609	8,144	—	1	6	11	54
Indiana	N	0	0	N	N	—	157	292	3,265	3,576	3	1	10	20	32
Michigan	4	14	38	241	267	225	300	880	6,145	5,042	—	0	5	12	18
Ohio	16	15	32	272	291	26	328	1,563	8,115	8,006	1	2	6	51	34
Wisconsin	—	9	27	93	184	48	131	181	2,575	2,869	—	1	4	7	32
<b>W.N. Central</b>	6	22	539	333	667	2	385	516	5,601	7,465	1	3	23	61	49
Iowa	1	5	16	70	88	—	40	63	695	708	—	0	1	1	—
Kansas	1	3	11	45	62	—	43	89	903	922	1	0	2	6	9
Minnesota	—	0	514	12	279	—	66	87	1,006	1,216	—	1	17	22	23
Missouri	4	9	28	149	164	—	195	269	2,354	3,958	—	1	5	25	13
Nebraska <sup>§</sup>	—	2	9	32	36	—	27	57	512	486	—	0	2	6	3
North Dakota	—	0	4	5	6	—	2	6	24	43	—	0	2	1	1
South Dakota	—	1	6	20	32	2	6	15	107	132	—	0	0	—	—
<b>S. Atlantic</b>	25	54	103	934	872	852	1,598	3,282	22,190	33,005	4	11	34	250	240
Delaware	—	1	4	11	10	10	27	44	543	591	—	0	3	5	1
District of Columbia	—	1	7	28	23	26	37	63	898	729	—	0	2	2	1
Florida	13	24	44	457	355	—	425	549	1,564	8,894	—	3	8	76	78
Georgia	12	12	28	176	213	—	339	2,068	3,159	6,264	3	2	7	52	57
Maryland <sup>§</sup>	—	4	12	85	55	—	130	189	2,155	2,864	—	2	5	43	30
North Carolina	—	0	0	—	—	349	328	676	6,427	6,930	1	0	9	33	15
North Carolina <sup>§</sup>	—	1	8	25	42	310	176	1,026	4,681	3,936	—	1	4	24	20
Virginia <sup>§</sup>	—	9	28	140	164	147	125	238	2,472	2,471	—	1	7	7	28
West Virginia	—	0	21	12	10	10	18	44	291	326	—	0	6	8	10
<b>E.S. Central</b>	—	9	34	163	142	299	551	879	9,277	12,159	—	2	9	47	56
Alabama <sup>§</sup>	—	3	22	82	73	36	170	271	996	4,490	—	0	3	10	12
Kentucky	N	0	0	N	N	61	50	268	1,107	1,317	—	0	1	2	4
Mississippi	N	0	0	N	N	—	158	434	3,206	2,615	—	0	1	—	5
Tennessee <sup>§</sup>	—	5	12	81	69	202	194	240	3,968	3,737	—	1	6	35	35
<b>W.S. Central</b>	5	7	53	113	59	551	941	1,490	17,584	19,564	5	2	30	49	32
Arkansas <sup>§</sup>	3	3	13	50	29	65	80	142	1,582	1,793	—	0	2	3	2
Louisiana	—	1	6	22	1	—	193	366	3,169	4,093	—	0	3	4	1
Oklahoma	2	2	40	41	29	84	94	236	2,026	1,722	5	1	27	39	27
Texas <sup>§</sup>	N	0	0	N	N	402	560	938	10,807	11,956	—	0	2	3	2
<b>Mountain</b>	29	30	67	501	552	58	281	456	3,886	5,729	5	4	11	127	102
Arizona	3	3	11	68	55	3	104	220	1,338	1,989	—	2	6	56	38
Colorado	10	9	26	171	182	40	67	93	875	1,456	3	1	4	27	30
Idaho <sup>§</sup>	—	3	12	39	60	—	2	20	84	82	—	0	1	4	3
Montana <sup>§</sup>	—	2	11	30	26	—	3	20	38	58	—	0	0	—	—
Nevada <sup>§</sup>	7	2	9	43	40	14	48	135	808	1,050	—	0	2	6	6
New Mexico <sup>§</sup>	—	1	6	32	23	—	30	64	443	669	—	0	4	14	15
Utah	9	6	27	106	159	—	16	28	276	367	2	0	3	19	10
Wyoming <sup>§</sup>	—	1	4	12	7	1	2	5	24	58	—	0	1	1	—
<b>Pacific</b>	27	57	558	1,100	1,044	437	765	935	14,955	16,748	1	2	16	46	41
Alaska	—	1	17	20	17	5	10	27	167	220	1	0	2	5	4
California	17	43	93	770	855	331	638	804	12,640	13,838	—	0	10	—	10
Hawaii	—	1	4	25	21	—	14	26	249	422	—	0	2	2	8
Oregon <sup>§</sup>	4	9	14	150	151	21	26	46	440	568	—	1	6	39	19
Washington	6	0	449	135	—	80	72	142	1,459	1,700	—	0	5	—	—
American Samoa	U	0	0	U	U	U	0	4	U	U	U	0	0	U	U
C.N.M.I.	U	—	—	U	U	U	—	—	U	U	U	—	—	U	U
Guam	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Puerto Rico	—	6	19	69	52	5	6	16	140	130	—	0	2	1	1
U.S. Virgin Islands	U	0	0	U	U	U	0	3	U	U	U	0	0	U	U

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

\* Incidence data for reporting years 2006 and 2007 are provisional.

† Data for *H. influenzae* (age <5 yrs for serotype b, nonserotype b, and unknown serotype) are available in Table I.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending May 26, 2007, and May 27, 2006 (21st Week)\*

Reporting area	Hepatitis (viral, acute), by type <sup>†</sup>										Legionellosis				
	A					B									
	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006
	Med	Max				Med	Max				Med	Max			
<b>United States</b>	17	56	175	957	1,512	16	79	390	1,458	1,623	10	53	114	507	532
<b>New England</b>	2	2	21	24	107	—	2	5	28	53	1	3	16	23	27
Connecticut	2	0	3	7	13	—	0	5	14	23	1	0	9	4	5
Maine <sup>§</sup>	—	0	2	—	5	—	0	2	1	10	—	0	2	—	3
Massachusetts	—	1	4	8	79	—	0	1	2	15	—	1	11	13	16
New Hampshire	—	0	15	5	2	—	0	2	5	—	—	0	2	—	2
Rhode Island <sup>§</sup>	—	0	2	3	2	—	0	4	5	4	—	0	6	5	—
Vermont <sup>§</sup>	—	0	2	1	6	—	0	1	1	1	—	0	2	1	1
<b>Mid. Atlantic</b>	2	7	18	123	139	—	9	19	173	204	—	15	57	126	146
New Jersey	—	1	4	21	40	—	2	6	30	63	—	2	11	12	20
New York (Upstate)	—	2	12	31	31	—	1	14	36	26	—	5	30	40	49
New York City	—	2	10	46	44	—	2	6	37	46	—	3	24	19	20
Pennsylvania	2	1	4	25	24	—	3	7	70	69	—	5	19	55	57
<b>E.N. Central</b>	2	6	17	89	125	1	9	23	171	188	1	11	31	99	108
Illinois	—	1	7	18	29	—	2	5	39	65	—	1	13	1	23
Indiana	—	0	7	5	10	—	0	21	14	14	—	1	6	6	3
Michigan	1	2	8	32	40	1	2	8	47	60	—	3	10	39	21
Ohio	1	1	4	27	33	—	3	10	65	44	1	4	19	49	47
Wisconsin	—	0	4	7	13	—	0	3	6	5	—	0	3	4	14
<b>W.N. Central</b>	—	2	17	62	57	—	2	14	54	55	2	1	16	16	18
Iowa	—	0	3	11	4	—	0	3	9	8	—	0	3	2	2
Kansas	—	0	1	2	19	—	0	2	4	6	—	0	3	—	1
Minnesota	—	0	17	33	3	—	0	13	4	6	2	0	11	4	—
Missouri	—	1	2	10	18	—	1	5	32	31	—	0	2	8	9
Nebraska <sup>§</sup>	—	0	2	4	8	—	0	3	3	3	—	0	2	1	4
North Dakota	—	0	0	—	—	—	0	0	—	—	—	0	0	—	—
South Dakota	—	0	2	2	5	—	0	1	2	1	—	0	1	1	2
<b>S. Atlantic</b>	2	9	28	172	204	8	21	55	386	484	2	8	24	122	124
Delaware	—	0	1	1	8	—	0	3	6	21	—	0	2	1	2
District of Columbia	—	0	5	14	2	—	0	2	1	4	—	0	5	1	4
Florida	—	3	13	57	72	7	7	14	141	172	2	2	9	57	61
Georgia	2	1	4	27	20	1	3	10	43	75	—	1	3	10	5
Maryland <sup>§</sup>	—	1	7	26	29	—	2	7	35	76	—	2	8	24	20
North Carolina	—	0	11	7	40	—	1	16	56	68	—	0	5	13	14
South Carolina <sup>§</sup>	—	0	3	4	10	—	2	5	29	28	—	0	2	5	3
Virginia <sup>§</sup>	—	1	5	34	22	—	2	5	55	15	—	1	4	8	14
West Virginia	—	0	3	2	1	—	0	23	20	25	—	0	4	3	1
<b>E.S. Central</b>	—	2	7	31	52	1	6	20	102	139	1	2	9	29	23
Alabama <sup>§</sup>	—	0	2	7	3	—	2	10	40	36	—	0	2	3	5
Kentucky	—	0	2	5	23	1	1	3	6	34	1	1	6	13	6
Mississippi	—	0	4	4	4	—	0	8	8	17	—	0	2	—	1
Tennessee <sup>§</sup>	—	1	5	15	22	—	3	7	48	52	—	1	7	13	11
<b>W.S. Central</b>	—	6	19	63	130	1	19	159	257	250	—	1	15	26	11
Arkansas <sup>§</sup>	—	0	2	4	31	—	1	7	7	27	—	0	1	1	1
Louisiana	—	0	4	8	3	—	1	5	17	11	—	0	2	1	—
Oklahoma	—	0	3	3	3	1	1	41	14	1	—	0	6	—	1
Texas <sup>§</sup>	—	5	15	48	93	—	15	108	219	211	—	1	12	24	9
<b>Mountain</b>	5	5	17	127	128	2	3	9	89	58	—	2	8	31	37
Arizona	3	3	14	104	70	—	0	5	38	4	—	0	4	10	13
Colorado	2	1	3	11	21	—	1	2	15	15	—	0	2	6	5
Idaho <sup>§</sup>	—	0	1	2	6	—	0	2	4	6	—	0	3	1	4
Montana <sup>§</sup>	—	0	3	1	4	—	0	0	—	—	—	0	1	1	1
Nevada <sup>§</sup>	—	0	2	6	7	2	1	5	19	16	—	0	2	3	4
New Mexico <sup>§</sup>	—	0	2	1	9	—	0	2	4	8	—	0	2	2	1
Utah	—	0	1	2	10	—	0	4	9	9	—	0	2	6	9
Wyoming <sup>§</sup>	—	0	1	—	1	—	0	1	—	—	—	0	1	2	—
<b>Pacific</b>	4	14	92	266	570	3	10	105	198	192	3	1	11	35	38
Alaska	—	0	1	2	1	1	0	3	4	1	—	0	1	—	—
California	4	13	40	241	543	1	8	31	146	157	2	1	11	27	38
Hawaii	—	0	2	2	6	—	0	1	—	4	—	0	0	—	—
Oregon <sup>§</sup>	—	1	3	11	20	—	2	5	29	30	—	0	1	1	—
Washington	—	0	52	10	—	1	0	74	19	—	1	0	2	7	—
American Samoa	U	0	0	U	U	U	0	0	U	U	U	0	0	U	U
C.N.M.I.	U	—	—	U	U	U	—	—	U	U	U	—	—	U	U
Guam	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Puerto Rico	—	1	10	25	20	—	1	9	20	19	—	0	2	2	1
U.S. Virgin Islands	U	0	0	U	U	U	0	0	U	U	U	0	0	U	U

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: Not reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

\* Incidence data for reporting years 2006 and 2007 are provisional.

<sup>†</sup> Data for acute hepatitis C, viral are available in Table I.

<sup>§</sup> Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

**TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending May 26, 2007, and May 27, 2006 (21st Week)\***

Reporting area	Lyme disease					Malaria					Meningococcal disease, invasive† All serogroups				
	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006
		Med	Max				Med	Max				Med	Max		
<b>United States</b>	36	255	1,116	2,221	2,729	8	23	78	295	461	16	19	79	456	551
<b>New England</b>	14	37	350	139	334	—	1	7	12	25	1	1	3	20	22
Connecticut	14	9	227	62	73	—	0	3	—	1	1	0	2	4	6
Maine§	—	1	38	18	34	—	0	1	3	2	—	0	3	3	2
Massachusetts	—	1	112	2	201	—	0	4	8	20	—	0	3	10	13
New Hampshire	—	6	97	46	16	—	0	3	1	1	—	0	2	—	—
Rhode Island§	—	0	93	—	1	—	0	1	—	—	—	0	1	1	—
Vermont§	—	1	15	11	9	—	0	0	—	1	—	0	1	2	1
<b>Mid. Atlantic</b>	13	142	552	1,106	1,631	1	5	18	65	112	1	2	8	54	90
New Jersey	—	26	190	102	462	—	0	7	—	33	—	0	2	1	9
New York (Upstate)	—	52	392	348	669	—	1	7	16	9	—	1	2	14	18
New York City	—	3	23	6	22	—	3	9	40	58	—	1	4	16	33
Pennsylvania	13	39	223	650	478	1	1	4	9	12	1	0	5	23	30
<b>E.N. Central</b>	—	6	162	25	254	2	3	10	36	54	1	3	8	61	85
Illinois	—	1	16	4	12	—	1	6	10	20	—	0	3	13	25
Indiana	—	0	3	1	2	—	0	2	1	6	—	0	4	14	10
Michigan	—	1	5	7	3	—	0	2	7	8	—	0	3	13	14
Ohio	—	0	5	3	15	2	0	2	11	14	1	1	3	15	24
Wisconsin	—	5	154	10	222	—	0	3	7	6	—	0	2	6	12
<b>W.N. Central</b>	7	5	188	63	75	—	1	12	19	20	1	1	5	30	33
Iowa	—	1	8	8	25	—	0	1	2	1	—	0	3	7	9
Kansas	1	0	3	7	1	—	0	2	1	—	—	0	1	1	1
Minnesota	6	2	188	41	46	—	0	12	11	14	1	0	3	9	7
Missouri	—	0	3	7	—	—	0	1	2	3	—	0	3	8	10
Nebraska§	—	0	2	—	3	—	0	1	2	—	—	0	1	2	5
North Dakota	—	0	0	—	—	—	0	0	—	1	—	0	1	2	1
South Dakota	—	0	1	—	—	—	0	1	1	1	—	0	1	1	—
<b>S. Atlantic</b>	—	44	134	808	404	2	5	14	70	118	—	3	11	65	95
Delaware	—	8	28	174	148	—	0	1	2	3	—	0	1	—	3
District of Columbia	—	0	7	6	7	—	0	2	3	—	—	0	1	—	—
Florida	—	0	3	13	8	—	1	4	17	19	—	1	7	25	37
Georgia	—	0	1	—	1	—	1	5	6	43	—	0	3	7	9
Maryland§	—	23	106	478	213	—	1	4	20	20	—	0	2	14	6
North Carolina	—	0	4	6	9	2	0	4	7	11	—	0	6	6	15
South Carolina§	—	0	2	5	3	—	0	2	1	4	—	0	2	6	11
Virginia§	—	7	36	122	15	—	1	4	13	17	—	0	2	7	11
West Virginia	—	0	14	4	—	—	0	1	1	1	—	0	2	—	3
<b>E.S. Central</b>	—	1	4	11	2	1	0	3	12	9	—	1	4	25	21
Alabama§	—	0	3	2	1	—	0	2	1	4	—	0	2	6	4
Kentucky	—	0	2	—	—	1	0	1	3	1	—	0	2	4	5
Mississippi	—	0	1	—	—	—	0	1	1	2	—	0	4	4	3
Tennessee§	—	0	3	9	1	—	0	2	7	2	—	0	2	11	9
<b>W.S. Central</b>	—	1	6	18	5	—	1	7	13	28	1	1	13	40	35
Arkansas§	—	0	0	—	—	—	0	2	—	1	—	0	2	5	5
Louisiana	—	0	1	2	—	—	0	2	11	1	—	0	4	11	5
Oklahoma	—	0	0	—	—	—	0	3	1	2	1	0	4	11	8
Texas§	—	1	6	16	5	—	1	6	1	24	—	0	9	13	17
<b>Mountain</b>	—	0	3	8	4	1	1	6	20	23	2	1	5	39	36
Arizona	—	0	1	—	3	—	0	3	4	7	—	0	3	10	10
Colorado	—	0	0	—	—	—	0	2	9	7	1	0	2	14	13
Idaho§	—	0	2	2	—	—	0	1	—	—	—	0	1	2	1
Montana§	—	0	1	1	—	—	0	1	1	1	—	0	1	1	2
Nevada§	—	0	2	5	—	—	0	1	1	—	—	0	1	3	3
New Mexico§	—	0	1	—	1	—	0	1	—	1	—	0	1	1	1
Utah	—	0	1	—	—	1	0	2	5	7	1	0	2	7	4
Wyoming§	—	0	1	—	—	—	0	0	—	—	—	0	2	1	2
<b>Pacific</b>	2	2	16	43	20	1	3	45	48	72	9	4	48	122	134
Alaska	—	0	1	2	—	—	0	4	2	8	—	0	1	1	2
California	2	2	8	41	20	1	2	6	33	56	8	3	10	90	105
Hawaii	N	0	0	N	N	—	0	1	2	2	—	0	1	2	4
Oregon§	—	0	1	—	—	—	0	3	8	6	1	0	3	15	23
Washington	—	0	8	—	—	—	0	43	3	—	—	0	43	14	—
American Samoa	U	0	0	U	U	U	0	0	U	U	U	0	0	—	—
C.N.M.I.	U	—	—	U	U	U	—	—	U	U	U	—	—	—	—
Guam	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Puerto Rico	N	0	0	N	N	—	0	1	1	—	—	0	1	5	4
U.S. Virgin Islands	U	0	0	U	U	U	0	0	U	U	U	0	0	—	—

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: Not reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

\* Incidence data for reporting years 2006 and 2007 are provisional.

† Data for meningococcal disease, invasive caused by serogroups A, C, Y, & W-135; serogroup B; other serogroup; and unknown serogroup are available in Table I.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending May 26, 2007, and May 27, 2006 (21st Week)\*

Reporting area	Pertussis					Rabies, animal					Rocky Mountain spotted fever				
	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006
		Med	Max				Med	Max				Med	Max		
<b>United States</b>	67	254	1,377	2,875	5,601	30	90	168	1,357	1,990	37	23	156	272	458
<b>New England</b>	—	37	78	451	885	5	11	25	190	202	—	0	9	—	4
Connecticut	—	2	10	18	26	5	4	14	66	51	—	0	0	—	—
Maine†	—	2	15	32	23	—	2	8	29	33	N	0	0	N	N
Massachusetts	—	28	45	369	663	—	0	7	—	86	—	0	1	—	4
New Hampshire	—	2	21	16	84	—	1	5	14	—	—	0	1	—	—
Rhode Island†	—	0	31	—	21	—	0	3	15	6	—	0	9	—	—
Vermont†	—	1	9	16	68	—	2	10	66	26	—	0	0	—	—
<b>Mid. Atlantic</b>	2	32	159	427	675	—	10	21	121	154	—	1	5	16	16
New Jersey	—	3	12	46	139	—	0	0	—	—	—	0	2	—	7
New York (Upstate)	—	19	150	258	241	—	0	0	—	—	—	0	2	—	—
New York City	—	1	6	—	31	—	1	5	24	3	—	0	3	6	4
Pennsylvania	2	9	20	123	264	—	9	20	97	151	—	0	3	10	5
<b>E.N. Central</b>	26	41	80	586	794	2	1	18	31	23	—	1	9	6	19
Illinois	—	9	23	62	201	—	0	7	3	6	—	0	4	1	11
Indiana	—	2	44	11	74	1	0	2	4	2	—	0	1	1	1
Michigan	1	10	39	109	150	1	0	5	7	15	—	0	1	1	—
Ohio	25	13	56	332	268	—	0	12	17	—	—	0	4	3	6
Wisconsin	—	3	17	72	101	—	0	0	—	—	—	0	0	—	1
<b>W.N. Central</b>	6	17	139	176	601	8	6	20	83	91	5	4	13	53	34
Iowa	—	4	16	52	154	—	1	7	9	12	—	0	1	—	1
Kansas	2	3	14	64	128	4	2	6	49	31	—	0	1	—	—
Minnesota	—	0	119	—	75	2	0	6	6	11	—	0	2	—	1
Missouri	4	3	10	35	167	2	1	6	8	9	5	3	12	52	30
Nebraska†	—	1	4	7	61	—	0	0	—	—	—	0	5	1	2
North Dakota	—	0	9	4	4	—	0	7	6	6	—	0	0	—	—
South Dakota	—	0	4	14	12	—	0	3	5	22	—	0	0	—	—
<b>S. Atlantic</b>	16	18	163	379	406	9	39	62	726	939	32	11	67	139	311
Delaware	—	0	1	2	2	—	0	0	—	—	—	0	3	4	7
District of Columbia	—	0	2	2	3	—	0	0	—	—	—	0	1	1	—
Florida	1	4	18	100	87	—	0	24	52	176	—	0	4	6	7
Georgia	—	0	3	5	9	—	4	9	46	99	—	0	5	3	9
Maryland†	—	2	7	48	74	—	5	10	93	156	—	1	6	16	15
North Carolina	15	1	112	145	77	9	11	21	197	152	32	4	61	90	254
South Carolina†	—	3	11	33	62	—	3	11	46	54	—	0	5	6	5
Virginia†	—	2	17	37	86	—	12	31	260	259	—	2	12	12	13
West Virginia	—	0	19	7	6	—	1	8	32	43	—	0	2	1	1
<b>E.S. Central</b>	—	6	24	77	112	—	4	13	60	100	—	5	27	54	58
Alabama†	—	1	17	23	25	—	0	8	—	33	—	1	9	12	14
Kentucky	—	0	5	2	17	—	0	4	8	6	—	0	1	1	—
Mississippi	—	0	9	9	15	—	0	1	—	4	—	0	1	—	—
Tennessee†	—	3	11	43	55	—	2	8	52	57	—	4	22	41	44
<b>W.S. Central</b>	—	17	152	172	254	—	15	34	30	343	—	1	114	3	9
Arkansas†	—	2	17	36	26	—	0	5	10	15	—	0	53	—	6
Louisiana	—	0	2	6	8	—	0	0	—	—	—	0	1	—	—
Oklahoma	—	0	9	1	2	—	0	7	20	24	—	0	55	—	1
Texas†	—	13	134	129	218	—	14	34	—	304	—	0	6	3	2
<b>Mountain</b>	15	29	63	494	1,330	1	2	28	33	60	—	0	4	1	6
Arizona	4	6	16	125	300	—	2	10	29	48	—	0	2	—	2
Colorado	6	7	18	135	476	—	0	0	—	—	—	0	1	—	1
Idaho†	—	1	7	18	30	—	0	24	—	—	—	0	3	1	—
Montana†	—	1	8	21	44	—	0	2	—	5	—	0	2	—	—
Nevada†	—	0	9	3	35	—	0	1	—	—	—	0	0	—	—
New Mexico†	—	2	8	13	37	—	0	1	1	5	—	0	1	—	2
Utah	5	10	48	167	377	1	0	1	2	1	—	0	0	—	—
Wyoming†	—	1	8	12	31	—	0	2	1	1	—	0	1	—	1
<b>Pacific</b>	2	25	546	113	544	5	4	13	83	78	—	0	1	—	1
Alaska	1	1	8	11	31	3	0	6	33	13	N	0	0	N	N
California	—	22	225	—	397	2	3	12	50	63	—	0	0	—	—
Hawaii	—	0	5	9	51	N	0	0	N	N	N	0	0	N	N
Oregon†	—	1	11	41	65	—	0	4	—	2	—	0	1	—	1
Washington	1	0	376	52	—	—	0	0	—	—	N	0	0	N	N
American Samoa	U	0	0	U	U	U	0	0	U	U	U	0	0	U	U
C.N.M.I.	U	—	—	U	U	U	—	—	U	U	U	—	—	U	U
Guam	—	—	—	—	—	—	—	—	—	—	N	—	—	N	N
Puerto Rico	—	0	1	—	—	—	1	6	19	45	N	0	0	N	N
U.S. Virgin Islands	U	0	0	U	U	U	0	0	U	U	U	0	0	U	U

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

\* Incidence data for reporting years 2006 and 2007 are provisional.

† Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

**TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending May 26, 2007, and May 27, 2006 (21st Week)\***

Reporting area	Salmonellosis					Shiga toxin-producing <i>E. coli</i> (STEC) <sup>†</sup>					Shigellosis				
	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006
		Med	Max				Med	Max				Med	Max		
<b>United States</b>	277	835	1,842	10,709	11,899	25	72	295	823	793	202	268	749	4,326	3,904
<b>New England</b>	1	39	121	561	1,142	—	4	22	50	114	—	4	21	74	191
Connecticut	—	0	107	107	503	—	0	9	9	75	—	0	11	11	67
Maine <sup>§</sup>	—	2	14	35	33	—	1	8	12	4	—	0	5	8	2
Massachusetts	—	24	87	335	548	—	2	13	21	30	—	3	18	50	115
New Hampshire	1	4	26	37	18	—	0	4	4	2	—	0	2	3	—
Rhode Island <sup>§</sup>	—	2	15	28	29	—	0	2	1	1	—	0	3	1	5
Vermont <sup>§</sup>	—	1	6	19	11	—	0	4	3	2	—	0	2	1	2
<b>Mid. Atlantic</b>	11	96	189	1,383	1,412	—	8	61	82	99	1	13	48	173	335
New Jersey	—	19	50	54	278	—	1	16	1	28	—	2	34	13	115
New York (Upstate)	—	28	93	431	301	—	3	14	36	33	—	3	43	39	87
New York City	1	23	45	368	387	—	0	4	8	14	1	5	12	94	97
Pennsylvania	10	31	66	530	446	—	3	47	37	24	—	1	6	27	36
<b>E.N. Central</b>	39	97	203	1,428	1,711	2	9	63	99	118	17	25	75	275	400
Illinois	—	29	65	290	486	—	1	8	12	14	—	9	53	35	135
Indiana	5	15	55	195	195	1	1	8	10	14	—	2	17	24	53
Michigan	7	18	35	274	301	—	1	6	19	25	—	2	5	14	77
Ohio	27	23	56	397	420	1	3	18	42	34	17	4	23	145	57
Wisconsin	—	17	32	272	309	—	2	41	16	31	—	4	14	57	78
<b>W.N. Central</b>	28	49	109	873	761	1	12	45	120	116	39	44	85	892	472
Iowa	—	8	26	125	132	—	2	38	19	22	—	2	14	20	16
Kansas	10	7	20	144	114	1	0	4	12	4	—	1	11	13	32
Minnesota	—	12	60	209	176	—	3	26	51	38	—	5	24	93	30
Missouri	18	16	35	279	213	—	3	13	23	37	39	14	78	742	321
Nebraska <sup>§</sup>	—	3	10	58	73	—	1	11	14	11	—	1	14	7	32
North Dakota	—	0	5	11	6	—	0	0	—	—	—	0	18	4	4
South Dakota	—	2	11	47	47	—	0	5	1	4	—	6	24	13	37
<b>S. Atlantic</b>	93	227	403	2,906	2,786	4	13	32	186	136	113	74	150	1,529	921
Delaware	—	2	10	29	27	—	0	3	6	1	—	0	2	4	—
District of Columbia	—	1	4	14	23	—	0	1	—	—	—	0	5	4	3
Florida	68	95	176	1,297	1,227	4	2	8	56	29	62	37	76	1,006	403
Georgia	19	29	76	423	403	—	2	7	20	23	51	25	62	413	327
Maryland <sup>§</sup>	—	14	32	201	132	—	3	9	34	12	—	1	10	25	19
North Carolina	6	29	130	444	453	—	2	11	25	28	—	1	14	25	82
South Carolina <sup>§</sup>	—	18	47	231	243	—	0	3	4	3	—	0	4	23	65
Virginia <sup>§</sup>	—	20	58	229	245	—	3	11	40	40	—	2	9	28	22
West Virginia	—	1	31	38	33	—	0	5	1	—	—	0	2	1	—
<b>E.S. Central</b>	5	51	139	672	669	—	4	21	37	55	3	12	84	323	265
Alabama <sup>§</sup>	—	11	70	199	224	—	0	5	8	6	—	6	66	141	67
Kentucky	5	9	23	153	119	—	1	12	12	13	3	2	15	41	130
Mississippi	—	12	86	86	139	—	0	3	—	1	—	1	71	71	31
Tennessee <sup>§</sup>	—	17	32	234	187	—	2	9	17	35	—	3	14	70	37
<b>W.S. Central</b>	13	84	186	410	1,038	3	4	53	49	41	8	38	245	399	489
Arkansas <sup>§</sup>	8	13	45	134	276	—	1	7	10	8	2	2	10	41	29
Louisiana	—	14	42	120	138	—	0	1	—	—	—	3	24	68	9
Oklahoma	5	10	103	120	86	3	0	17	11	4	6	2	60	28	32
Texas <sup>§</sup>	—	44	107	36	538	—	2	48	28	29	—	30	174	262	419
<b>Mountain</b>	33	50	88	843	835	4	8	34	104	89	5	22	84	260	301
Arizona	8	17	44	300	244	4	2	9	42	24	2	10	37	129	161
Colorado	10	12	30	233	246	—	1	8	19	23	—	3	15	45	45
Idaho <sup>§</sup>	—	3	9	38	48	—	1	8	6	13	—	0	3	4	6
Montana <sup>§</sup>	—	2	10	31	40	—	0	0	—	—	—	0	13	11	2
Nevada <sup>§</sup>	7	4	20	69	54	—	0	5	8	11	1	1	20	13	29
New Mexico <sup>§</sup>	—	4	15	56	73	—	1	5	11	7	—	2	15	33	37
Utah	8	4	14	93	105	—	2	14	18	10	2	1	4	8	18
Wyoming <sup>§</sup>	—	1	4	23	25	—	0	3	—	1	—	0	19	17	3
<b>Pacific</b>	54	105	890	1,633	1,545	11	3	164	96	25	16	33	256	401	530
Alaska	3	1	5	32	33	N	0	0	N	N	—	0	2	6	4
California	44	89	260	1,254	1,272	4	0	8	56	N	15	28	84	321	449
Hawaii	—	5	16	75	87	1	0	3	6	4	—	1	3	13	17
Oregon <sup>§</sup>	—	7	17	90	153	—	1	9	12	21	—	1	6	19	60
Washington	7	0	625	182	—	6	0	162	22	—	1	0	170	42	—
American Samoa	U	0	0	U	U	U	0	0	U	U	U	0	0	U	U
C.N.M.I.	U	—	—	U	U	U	—	—	U	U	U	—	—	U	U
Guam	—	—	—	—	—	N	—	—	N	N	—	—	—	—	—
Puerto Rico	—	15	66	250	127	—	0	0	—	—	—	0	6	11	9
U.S. Virgin Islands	U	0	0	U	U	U	0	0	U	U	U	0	0	U	U

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

\* Incidence data for reporting years 2006 and 2007 are provisional.

† Includes *E. coli* O157:H7; Shiga toxin-positive, serogroup non-O157; and Shiga toxin-positive, not serogrouped.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending May 26, 2007, and May 27, 2006 (21st Week)\*

Reporting area	Streptococcal disease, invasive, group A					<i>Streptococcus pneumoniae</i> , invasive disease† Age <5 years				
	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006
		Med	Max				Med	Max		
<b>United States</b>	44	93	247	2,251	2,756	9	28	103	663	647
<b>New England</b>	—	6	28	167	220	—	2	11	54	77
Connecticut	—	0	17	35	55	—	0	6	—	19
Maine§	—	0	2	8	9	—	0	1	1	—
Massachusetts	—	4	10	95	140	—	2	6	42	57
New Hampshire	—	0	9	18	6	—	0	4	6	1
Rhode Island§	—	0	12	—	4	—	0	3	3	—
Vermont§	—	0	2	11	6	—	0	1	2	—
<b>Mid. Atlantic</b>	6	16	39	416	523	—	3	19	53	89
New Jersey	—	1	6	28	94	—	0	4	—	31
New York (Upstate)	—	5	26	149	154	—	2	14	53	49
New York City	—	3	11	96	96	—	0	3	—	9
Pennsylvania	6	6	11	143	179	N	0	0	N	N
<b>E.N. Central</b>	11	15	29	400	593	—	6	14	98	170
Illinois	—	4	10	81	184	—	1	6	9	46
Indiana	4	2	12	57	63	—	0	10	10	21
Michigan	1	4	10	105	121	—	1	4	40	42
Ohio	6	4	14	138	154	—	1	7	35	34
Wisconsin	—	1	6	19	71	—	0	2	4	27
<b>W.N. Central</b>	4	5	32	186	177	5	2	9	60	52
Iowa	—	0	0	—	—	—	0	0	—	—
Kansas	1	1	3	24	36	—	0	3	1	11
Minnesota	—	0	29	86	78	5	1	6	40	25
Missouri	3	2	6	52	32	—	0	3	14	10
Nebraska§	—	0	2	11	18	—	0	2	4	4
North Dakota	—	0	2	9	6	—	0	1	1	2
South Dakota	—	0	2	4	7	—	0	0	—	—
<b>S. Atlantic</b>	13	20	48	504	531	1	3	12	127	33
Delaware	—	0	2	3	5	—	0	0	—	—
District of Columbia	—	0	3	7	7	—	0	1	—	—
Florida	7	5	16	127	125	—	0	5	31	—
Georgia	6	5	11	100	135	1	0	4	39	—
Maryland§	—	4	8	89	72	—	1	6	36	25
North Carolina	—	0	26	56	67	—	0	0	—	—
South Carolina§	—	1	7	45	40	—	0	3	11	—
Virginia§	—	2	11	67	66	—	0	3	8	—
West Virginia	—	0	5	10	14	—	0	4	2	8
<b>E.S. Central</b>	—	4	11	90	114	—	0	6	42	9
Alabama§	N	0	0	N	N	N	0	0	N	N
Kentucky	—	1	4	24	28	—	0	0	—	—
Mississippi	N	0	0	N	N	—	0	2	2	9
Tennessee§	—	3	7	66	86	—	0	6	40	—
<b>W.S. Central</b>	—	6	80	138	192	1	4	39	110	87
Arkansas§	—	0	2	12	17	1	0	2	7	14
Louisiana	—	0	2	4	2	—	0	4	24	2
Oklahoma	—	2	21	41	56	—	1	12	25	20
Texas§	—	3	56	81	117	—	1	24	54	51
<b>Mountain</b>	9	11	23	297	364	1	4	12	102	117
Arizona	2	5	11	117	196	—	2	7	57	69
Colorado	4	3	9	90	58	1	1	4	30	27
Idaho§	—	0	1	6	6	—	0	1	2	1
Montana§	N	0	0	N	N	N	0	0	N	N
Nevada§	—	0	1	2	1	—	0	1	1	—
New Mexico§	—	1	6	25	67	—	0	4	12	20
Utah	3	1	7	54	34	—	0	0	—	—
Wyoming§	—	0	1	3	2	—	0	0	—	—
<b>Pacific</b>	1	3	9	53	42	1	0	4	17	13
Alaska	1	0	2	15	N	1	0	2	15	—
California	N	0	0	N	N	N	0	0	N	N
Hawaii	—	2	9	38	42	—	0	2	2	13
Oregon§	N	0	0	N	N	N	0	0	N	N
Washington	N	0	0	N	N	N	0	0	N	N
American Samoa	U	0	0	U	U	U	0	0	U	U
C.N.M.I.	U	—	—	U	U	U	—	—	U	U
Guam	—	—	—	—	—	N	—	—	N	N
Puerto Rico	—	0	0	—	—	N	0	0	N	N
U.S. Virgin Islands	U	0	0	U	U	U	0	0	U	U

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

\* Incidence data for reporting years 2006 and 2007 are provisional.

† Includes cases of invasive pneumococcal disease, in children aged <5 years, caused by *S. pneumoniae*, which is susceptible or for which susceptibility testing is not available (NNDS event code 11717).

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

**TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending May 26, 2007, and May 27, 2006 (21st Week)\***

Reporting area	<i>Streptococcus pneumoniae</i> , invasive disease, drug resistant†										Syphilis, primary and secondary				
	All ages				Age <5 years										
	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Current week	Previous 52 weeks		Cum 2007	Cum 2006
		Med	Max				Med	Max				Med	Max		
<b>United States</b>	27	46	254	1,147	1,308	2	7	31	168	163	72	183	315	3,223	3,600
<b>New England</b>	—	1	12	24	73	—	0	3	4	2	4	4	13	85	76
Connecticut	—	0	5	—	57	—	0	0	—	—	—	0	10	10	17
Maine§	—	0	2	5	4	—	0	2	1	1	—	0	1	1	4
Massachusetts	—	0	0	—	—	—	0	0	—	—	3	3	7	54	42
New Hampshire	—	0	0	—	—	—	0	0	—	—	1	0	2	10	5
Rhode Island§	—	0	4	8	4	—	0	1	1	—	—	0	5	9	6
Vermont§	—	0	2	11	8	—	0	1	2	1	—	0	1	1	2
<b>Mid. Atlantic</b>	—	3	8	74	75	—	0	5	17	10	1	23	44	592	465
New Jersey	—	0	0	—	—	—	0	0	—	—	—	3	8	57	71
New York (Upstate)	—	1	5	25	21	—	0	4	7	4	—	3	14	46	61
New York City	—	0	0	—	—	—	0	0	—	—	—	15	35	396	232
Pennsylvania	—	2	6	49	54	—	0	2	10	6	1	5	12	93	101
<b>E.N. Central</b>	9	10	40	284	284	1	1	7	35	47	7	15	32	276	357
Illinois	—	0	3	3	14	—	0	1	1	3	4	6	13	106	196
Indiana	3	2	31	66	68	—	0	5	6	13	—	2	5	18	31
Michigan	—	0	1	1	14	—	0	1	—	2	1	2	10	46	33
Ohio	6	5	38	214	188	1	1	5	28	29	2	4	9	82	81
Wisconsin	N	0	0	N	N	—	0	0	—	—	—	1	4	24	16
<b>W.N. Central</b>	—	1	124	88	21	—	0	15	7	1	—	5	14	54	106
Iowa	—	0	0	—	—	—	0	0	—	—	—	0	3	3	7
Kansas	—	0	10	46	—	—	0	2	2	—	—	0	3	8	10
Minnesota	—	0	123	—	—	—	0	15	—	—	—	1	5	21	22
Missouri	—	1	6	35	21	—	0	2	3	1	—	2	8	21	64
Nebraska§	—	0	1	2	—	—	0	0	—	—	—	0	2	1	2
North Dakota	—	0	0	—	—	—	0	0	—	—	—	0	0	—	1
South Dakota	—	0	3	5	—	—	0	1	2	—	—	0	3	—	—
<b>S. Atlantic</b>	18	21	59	517	691	1	3	8	74	60	15	40	185	541	787
Delaware	—	0	1	4	—	—	0	1	1	—	—	0	3	5	12
District of Columbia	—	0	2	5	17	—	0	0	—	2	3	2	11	62	47
Florida	14	11	29	302	312	1	2	8	66	57	—	12	23	68	293
Georgia	4	6	21	173	295	—	0	1	—	1	—	4	153	20	91
Maryland§	—	0	1	1	—	—	0	0	—	—	—	5	15	116	136
North Carolina	—	0	0	—	—	—	0	0	—	—	2	5	23	140	117
South Carolina§	—	0	0	—	—	—	0	0	—	—	1	1	10	43	34
Virginia§	N	0	0	N	N	—	0	0	—	—	9	4	17	84	56
West Virginia	—	1	17	32	67	—	0	1	7	—	—	0	2	3	1
<b>E.S. Central</b>	—	2	9	73	98	—	0	3	15	16	16	14	29	294	232
Alabama§	N	0	0	N	N	—	0	0	—	—	11	5	17	95	101
Kentucky	—	0	2	15	23	—	0	1	1	3	2	1	7	32	32
Mississippi	—	0	0	—	—	—	0	0	—	—	—	2	9	47	24
Tennessee§	—	2	8	58	75	—	0	3	14	13	3	6	13	120	75
<b>W.S. Central</b>	—	1	9	61	11	—	0	2	8	3	24	29	56	598	562
Arkansas§	—	0	3	1	5	—	0	0	—	2	6	1	7	43	33
Louisiana	—	1	3	22	6	—	0	1	2	1	—	6	30	126	79
Oklahoma	—	0	8	38	—	—	0	2	6	—	3	1	5	31	32
Texas§	—	0	0	—	—	—	0	0	—	—	15	21	31	398	418
<b>Mountain</b>	—	1	5	26	55	—	0	5	8	24	3	8	27	107	195
Arizona	—	0	0	—	—	—	0	0	—	—	—	2	16	29	81
Colorado	—	0	0	—	—	—	0	0	—	—	—	1	5	12	34
Idaho§	N	0	0	N	N	—	0	0	—	—	—	0	1	1	2
Montana§	—	0	0	—	—	—	0	0	—	—	—	0	1	1	1
Nevada§	—	0	3	15	13	—	0	2	5	—	3	2	12	36	47
New Mexico§	—	0	0	—	—	—	0	0	—	—	—	1	7	24	26
Utah	—	0	5	8	24	—	0	4	2	16	—	0	2	3	4
Wyoming§	—	0	3	3	18	—	0	1	1	8	—	0	1	1	—
<b>Pacific</b>	—	0	0	—	—	—	0	0	—	—	2	38	57	676	820
Alaska	—	0	0	—	—	—	0	0	—	—	—	0	2	4	5
California	N	0	0	N	N	—	0	0	—	—	1	35	54	614	718
Hawaii	—	0	0	—	—	—	0	0	—	—	—	0	1	2	10
Oregon§	N	0	0	N	N	—	0	0	—	—	—	0	6	8	7
Washington	N	0	0	N	N	—	0	0	—	—	1	2	11	48	80
American Samoa	U	0	0	U	U	U	0	1	U	U	U	0	0	U	U
C.N.M.I.	U	—	—	U	U	U	—	—	U	U	U	—	—	U	U
Guam	N	—	—	N	N	—	—	—	—	—	—	—	—	—	—
Puerto Rico	N	0	0	N	N	—	0	0	—	—	4	3	11	56	61
U.S. Virgin Islands	U	0	0	U	U	U	0	0	U	U	U	0	0	U	U

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

\* Incidence data for reporting years 2006 and 2007 are provisional.

† Includes cases of invasive pneumococcal disease caused by drug-resistant *S. pneumoniae* (DRSP) (NNDSS event code 11720).

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).



TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending May 26, 2007, and May 27, 2006 (21st Week)\*

Reporting area	Varicella (chickenpox)					West Nile virus disease†									
	Current week	Previous 52 weeks		Cum 2007	Cum 2006	Neuroinvasive			Non-neuroinvasive§						
		Med	Max			Current week	Previous 52 weeks	Cum 2007	Cum 2006	Current week	Previous 52 weeks	Cum 2007	Cum 2006		
<b>United States</b>	437	808	1,579	19,159	25,322	—	0	178	—	10	—	1	399	—	5
<b>New England</b>	1	31	215	310	2,183	—	0	3	—	—	—	0	2	—	—
Connecticut	—	10	76	1	865	—	0	3	—	—	—	0	1	—	—
Maine¶	—	1	17	—	146	—	0	0	—	—	—	0	0	—	—
Massachusetts	—	0	95	—	776	—	0	1	—	—	—	0	1	—	—
New Hampshire	1	6	43	123	63	—	0	0	—	—	—	0	0	—	—
Rhode Island¶	—	0	0	—	—	—	0	0	—	—	—	0	0	—	—
Vermont¶	—	9	66	186	333	—	0	0	—	—	—	0	0	—	—
<b>Mid. Atlantic</b>	91	106	195	2,369	2,648	—	0	11	—	—	—	0	4	—	—
New Jersey	N	0	0	N	N	—	0	2	—	—	—	0	1	—	—
New York (Upstate)	N	0	0	N	N	—	0	5	—	—	—	0	1	—	—
New York City	—	0	0	—	—	—	0	4	—	—	—	0	2	—	—
Pennsylvania	91	106	195	2,369	2,648	—	0	2	—	—	—	0	1	—	—
<b>E.N. Central</b>	208	218	568	5,637	8,885	—	0	43	—	1	—	0	33	—	—
Illinois	—	2	11	71	69	—	0	23	—	1	—	0	23	—	—
Indiana	—	0	0	—	—	—	0	7	—	—	—	0	12	—	—
Michigan	72	88	258	2,220	2,581	—	0	11	—	—	—	0	2	—	—
Ohio	132	118	449	2,873	5,561	—	0	11	—	—	—	0	3	—	—
Wisconsin	4	15	57	473	674	—	0	2	—	—	—	0	2	—	—
<b>W.N. Central</b>	58	32	136	1,083	1,092	—	0	36	—	—	—	0	79	—	1
Iowa	N	0	0	N	N	—	0	3	—	—	—	0	4	—	1
Kansas	4	9	52	403	215	—	0	3	—	—	—	0	3	—	—
Minnesota	—	0	0	—	—	—	0	6	—	—	—	0	7	—	—
Missouri	54	16	78	546	828	—	0	14	—	—	—	0	2	—	—
Nebraska¶	N	0	0	N	N	—	0	9	—	—	—	0	38	—	—
North Dakota	—	0	60	84	18	—	0	5	—	—	—	0	28	—	—
South Dakota	—	1	15	50	31	—	0	7	—	—	—	0	22	—	—
<b>S. Atlantic</b>	41	85	224	2,185	2,457	—	0	2	—	—	—	0	7	—	—
Delaware	—	0	6	12	40	—	0	0	—	—	—	0	0	—	—
District of Columbia	—	0	8	8	18	—	0	0	—	—	—	0	1	—	—
Florida	19	0	89	655	N	—	0	1	—	—	—	0	0	—	—
Georgia	N	0	0	N	N	—	0	1	—	—	—	0	4	—	—
Maryland¶	N	0	0	N	N	—	0	2	—	—	—	0	2	—	—
North Carolina	—	0	0	—	—	—	0	1	—	—	—	0	0	—	—
South Carolina¶	—	18	72	572	710	—	0	1	—	—	—	0	0	—	—
Virginia¶	—	19	176	331	811	—	0	0	—	—	—	0	2	—	—
West Virginia	22	25	52	607	878	—	0	1	—	—	—	0	0	—	—
<b>E.S. Central</b>	—	6	43	246	50	—	0	15	—	3	—	0	16	—	—
Alabama¶	—	6	43	244	50	—	0	2	—	—	—	0	0	—	—
Kentucky	N	0	0	N	N	—	0	2	—	—	—	0	1	—	—
Mississippi	—	0	2	2	—	—	0	10	—	3	—	0	16	—	—
Tennessee¶	N	0	0	N	N	—	0	4	—	—	—	0	2	—	—
<b>W.S. Central</b>	1	200	979	5,749	6,340	—	0	58	—	4	—	0	26	—	2
Arkansas¶	1	9	105	178	422	—	0	4	—	—	—	0	2	—	—
Louisiana	—	1	11	46	46	—	0	13	—	—	—	0	9	—	1
Oklahoma	—	0	0	—	—	—	0	6	—	—	—	0	4	—	—
Texas¶	—	172	873	5,525	5,872	—	0	38	—	4	—	0	16	—	1
<b>Mountain</b>	37	56	129	1,558	1,667	—	0	61	—	2	—	0	228	—	2
Arizona	—	0	0	—	—	—	0	9	—	—	—	0	15	—	—
Colorado	28	22	62	612	863	—	0	10	—	2	—	0	51	—	1
Idaho¶	N	0	0	N	N	—	0	30	—	—	—	0	157	—	1
Montana¶	—	0	26	194	N	—	0	3	—	—	—	0	8	—	—
Nevada¶	—	0	3	1	8	—	0	9	—	—	—	0	16	—	—
New Mexico¶	—	4	35	216	283	—	0	1	—	—	—	0	1	—	—
Utah	9	17	73	522	491	—	0	8	—	—	—	0	17	—	—
Wyoming¶	—	0	11	13	22	—	0	7	—	—	—	0	10	—	—
<b>Pacific</b>	—	0	9	22	—	—	0	15	—	—	—	0	51	—	—
Alaska	—	0	9	22	N	—	0	0	—	—	—	0	0	—	—
California	—	0	0	—	N	—	0	15	—	—	—	0	37	—	—
Hawaii	—	0	0	—	—	—	0	0	—	—	—	0	0	—	—
Oregon¶	N	0	0	N	N	—	0	2	—	—	—	0	14	—	—
Washington	N	0	0	N	N	—	0	0	—	—	—	0	2	—	—
American Samoa	U	0	0	U	U	U	0	0	U	U	U	0	0	U	U
C.N.M.I.	U	—	—	U	U	U	—	—	U	U	U	—	—	U	U
Guam	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Puerto Rico	1	12	26	305	247	—	0	0	—	—	—	0	0	—	—
U.S. Virgin Islands	U	0	0	U	U	U	0	0	U	U	U	0	0	U	U

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

† Incidence data for reporting years 2006 and 2007 are provisional.

¶ Updated weekly from reports to the Division of Vector-Borne Infectious Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases (ArboNET Surveillance). Data for California serogroup, eastern equine, Powassan, St. Louis, and western equine diseases are available in Table I.

§ Not notifiable in all states. Data from states where the condition is not notifiable are excluded from this table, except in 2007 for the domestic arboviral diseases and influenza-associated pediatric mortality, and in 2003 for SARS-CoV. Reporting exceptions are available at <http://www.cdc.gov/epo/dphsi/phs/infdis.htm>.

¶ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE III. Deaths in 122 U.S. cities,\* week ending May 26, 2007 (21st Week)

Reporting Area	All causes, by age (years)							Reporting Area	All causes, by age (years)						
	All Ages	≥65	45-64	25-44	1-24	<1	P&I <sup>†</sup> Total		All Ages	≥65	45-64	25-44	1-24	<1	P&I <sup>†</sup> Total
<b>New England</b>	521	387	95	21	8	10	47	<b>S. Atlantic</b>	1,077	641	273	98	36	29	60
Boston, MA	148	106	28	7	4	3	14	Atlanta, GA	26	7	7	8	4	—	1
Bridgeport, CT	33	24	7	1	1	—	2	Baltimore, MD	151	82	35	18	9	7	10
Cambridge, MA	11	11	—	—	—	—	—	Charlotte, NC	112	68	29	11	2	2	13
Fall River, MA	20	18	—	2	—	—	5	Jacksonville, FL	175	98	50	20	4	3	1
Hartford, CT	32	22	8	2	—	—	3	Miami, FL	85	55	20	5	5	—	8
Lowell, MA	26	19	5	—	1	1	4	Norfolk, VA	52	35	10	2	—	5	1
Lynn, MA	6	5	1	—	—	—	1	Richmond, VA	48	29	14	4	1	—	4
New Bedford, MA	20	16	4	—	—	—	1	Savannah, GA	40	26	11	2	1	—	1
New Haven, CT	30	18	4	4	1	3	7	St. Petersburg, FL	57	35	12	3	4	3	3
Providence, RI	58	50	7	—	—	1	—	Tampa, FL	201	131	48	12	4	6	15
Somerville, MA	2	1	—	—	—	1	—	Washington, D.C.	119	69	34	11	2	3	2
Springfield, MA	48	31	13	2	1	1	5	Wilmington, DE	11	6	3	2	—	—	1
Waterbury, CT	23	21	2	—	—	—	2	<b>E.S. Central</b>	890	593	195	62	20	20	76
Worcester, MA	64	45	16	3	—	—	3	Birmingham, AL	161	108	39	12	—	2	10
<b>Mid. Atlantic</b>	1,945	1,333	432	122	34	23	100	Chattanooga, TN	85	57	20	3	2	3	6
Albany, NY	33	17	9	4	1	2	1	Knoxville, TN	94	61	20	8	3	2	9
Allentown, PA	27	24	1	2	—	—	2	Lexington, KY	55	39	13	1	1	1	3
Buffalo, NY	102	69	25	6	—	2	4	Memphis, TN	180	124	38	11	2	5	23
Camden, NJ	12	7	1	3	—	1	—	Mobile, AL	124	76	26	13	9	—	5
Elizabeth, NJ	18	13	4	1	—	—	3	Montgomery, AL	44	31	11	2	—	—	5
Erie, PA	45	38	7	—	—	—	2	Nashville, TN	147	97	28	12	3	7	15
Jersey City, NJ	31	21	6	2	1	1	5	<b>W.S. Central</b>	1,226	808	280	82	25	31	70
New York City, NY	1,011	692	235	58	17	8	39	Austin, TX	106	65	27	12	—	2	11
Newark, NJ	33	12	16	5	—	—	3	Baton Rouge, LA	45	23	16	5	—	1	1
Paterson, NJ	16	8	1	6	1	—	2	Corpus Christi, TX	58	48	7	2	—	1	4
Philadelphia, PA	245	154	60	20	6	5	11	Dallas, TX	U	U	U	U	U	U	U
Pittsburgh, PA <sup>‡</sup>	27	20	6	1	—	—	3	El Paso, TX	46	28	12	4	—	2	2
Reading, PA	35	30	3	1	1	—	2	Fort Worth, TX	110	77	24	4	1	4	7
Rochester, NY	141	110	22	5	1	3	13	Houston, TX	368	225	90	35	11	7	8
Schenectady, NY	24	20	2	1	1	—	2	Little Rock, AR	82	54	17	3	2	6	2
Scranton, PA	20	15	4	—	1	—	—	New Orleans, LA <sup>¶</sup>	U	U	U	U	U	U	U
Syracuse, NY	77	54	16	4	3	—	7	San Antonio, TX	209	142	42	12	8	5	17
Trenton, NJ	25	13	8	2	1	1	1	Shreveport, LA	73	48	20	3	—	2	8
Utica, NY	8	5	2	1	—	—	—	Tulsa, OK	129	98	25	2	3	1	10
Yonkers, NY	15	11	4	—	—	—	—	<b>Mountain</b>	654	422	152	48	12	20	39
<b>E.N. Central</b>	1,995	1,298	458	133	48	58	123	Albuquerque, NM	U	U	U	U	U	U	U
Akron, OH	51	30	16	2	2	1	1	Boise, ID	48	27	12	6	1	2	3
Canton, OH	32	24	4	3	—	1	6	Colorado Springs, CO	77	57	13	4	1	2	1
Chicago, IL	261	152	60	30	7	12	14	Denver, CO	94	55	25	7	3	4	9
Cincinnati, OH	94	51	24	9	5	5	7	Las Vegas, NV	255	159	57	24	6	9	16
Cleveland, OH	235	164	57	8	2	4	9	Ogden, UT	30	25	4	—	—	1	4
Columbus, OH	216	140	52	13	8	3	11	Phoenix, AZ	U	U	U	U	U	U	U
Dayton, OH	126	93	24	5	1	3	7	Pueblo, CO	36	30	6	—	—	—	3
Detroit, MI	149	85	38	18	3	5	16	Salt Lake City, UT	114	69	35	7	1	2	3
Evansville, IN	48	33	11	1	2	1	4	Tucson, AZ	U	U	U	U	U	U	U
Fort Wayne, IN	55	39	10	4	2	—	1	<b>Pacific</b>	1,083	786	196	59	20	22	81
Gary, IN	16	7	4	3	2	—	—	Berkeley, CA	13	11	1	1	—	—	2
Grand Rapids, MI	42	28	12	1	1	—	3	Fresno, CA	U	U	U	U	U	U	U
Indianapolis, IN	206	129	40	20	7	10	10	Glendale, CA	U	U	U	U	U	U	U
Lansing, MI	50	41	8	—	—	1	2	Honolulu, HI	69	51	13	2	1	2	6
Milwaukee, WI	102	59	26	8	3	6	8	Long Beach, CA	50	35	6	6	2	1	5
Peoria, IL	45	37	5	1	2	—	4	Los Angeles, CA	U	U	U	U	U	U	U
Rockford, IL	53	31	17	3	—	2	6	Pasadena, CA	17	12	2	2	—	1	2
South Bend, IN	52	40	11	1	—	—	3	Portland, OR	123	85	22	7	2	7	12
Toledo, OH	100	68	27	2	—	3	6	Sacramento, CA	181	135	38	6	1	1	11
Youngstown, OH	62	47	12	1	1	1	5	San Diego, CA	143	101	26	10	3	3	17
<b>W.N. Central</b>	662	420	157	42	13	30	42	San Francisco, CA	U	U	U	U	U	U	U
Des Moines, IA	95	70	17	8	—	—	8	San Jose, CA	157	116	28	6	3	4	6
Duluth, MN	19	14	4	—	1	—	1	Santa Cruz, CA	28	20	3	3	2	—	2
Kansas City, KS	20	12	6	1	—	1	2	Seattle, WA	120	84	21	11	3	1	8
Kansas City, MO	86	60	18	2	4	2	4	Spokane, WA	60	47	11	2	—	—	5
Lincoln, NE	46	33	8	3	1	1	3	Tacoma, WA	122	89	25	3	3	2	5
Minneapolis, MN	70	36	21	4	2	7	4	<b>Total</b>	10,053**	6,688	2,238	667	216	243	638
Omaha, NE	79	50	20	3	—	6	6								
St. Louis, MO	111	54	33	13	3	8	9								
St. Paul, MN	37	29	7	1	—	—	2								
Wichita, KS	99	62	23	7	2	5	3								

U: Unavailable. —:No reported cases.

\* Mortality data in this table are voluntarily reported from 122 cities in the United States, most of which have populations of ≥100,000. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included.


† Pneumonia and influenza.

‡ Because of changes in reporting methods in this Pennsylvania city, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks.

¶ Because of Hurricane Katrina, weekly reporting of deaths has been temporarily disrupted.

\*\* Total includes unknown ages.

**FIGURE I. Selected notifiable disease reports, United States, comparison of provisional 4-week totals May 26, 2007, with historical data**



\* Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

**Notifiable Disease Data Team and 122 Cities Mortality Data Team**  
 Patsy A. Hall  
 Deborah A. Adams      Rosaline Dhara  
 Willie J. Anderson      Vernitta Love  
 Lenee Blanton      Pearl C. Sharp

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format. To receive an electronic copy each week, send an e-mail message to [listserv@listserv.cdc.gov](mailto:listserv@listserv.cdc.gov). The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's Internet server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/publications/mmwr>. Paper copy subscriptions are available through the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Data are compiled in the National Center for Public Health Informatics, Division of Integrated Surveillance Systems and Services. Address all inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop E-90, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333 or to [www.mmwrq@cdc.gov](mailto:www.mmwrq@cdc.gov).

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.