



MMWR™

Morbidity and Mortality Weekly Report

www.cdc.gov/mmwr

Weekly

September 11, 2009 / Vol. 58 / No. 35

Oseltamivir-Resistant 2009 Pandemic Influenza A (H1N1) Virus Infection in Two Summer Campers Receiving Prophylaxis – North Carolina, 2009

Initial testing of the 2009 pandemic influenza A (H1N1) virus found it susceptible to neuraminidase inhibitors (oseltamivir and zanamivir) and resistant to adamantanes (amantadine and rimantadine) (1). Neuraminidase inhibitors have been used widely for treatment and chemoprophylaxis of 2009 pandemic influenza A (H1N1); however, sporadic cases of oseltamivir-resistant 2009 pandemic influenza A (H1N1) virus infection have been reported worldwide (2), including nine U.S. cases identified as of September 4.* On July 14, CDC was contacted by a physician at a summer camp in North Carolina regarding two cases of influenza-like illness (ILI) in adolescent girls receiving oseltamivir chemoprophylaxis during an ILI outbreak that had begun June 18. The two girls stayed in the same cabin, and both received oseltamivir during a mass chemoprophylaxis program in which approximately 600 campers and staff members received oseltamivir or zanamivir. On July 20 and July 22, the North Carolina State Laboratory of Public Health confirmed pandemic H1N1 virus infection in respiratory specimens from both girls. On August 14 and August 19, CDC detected the H275Y mutation (N1 numbering) in neuraminidase from both specimens by pyrosequencing (3,4). The H275Y mutation is associated with resistance to oseltamivir; zanamivir susceptibility is retained. A second mutation (I223V) in neuraminidase also was detected in both specimens. This is the first report of oseltamivir resistance in pandemic H1N1 cases with an epidemiologic link. Health-care providers should be aware that antiviral resistance can develop during chemoprophylaxis or treatment with subtherapeutic dosages and should follow published recommendations for antiviral medications (5).

* Additional information available at <http://www.cdc.gov/flu/weekly/fluactivity.htm>.

The summer camp offered two 4-week sessions. The first session was conducted from June 14 to July 10, followed by a weekend break, July 11–12, before the start of the second session on July 13. Approximately 650 campers and 350 staff members participated in the first session, and 350 campers and 300 staff members participated in the second session. An outbreak of ILI began on June 18, soon after the start of the first session; the last case was diagnosed on July 22. All ill persons were grouped in isolation until 7 days after symptom onset and until well. All but one of the 61 ill campers and four ill staff members received treatment with either oseltamivir or zanamivir. Also, beginning on June 18, medical staff members conducted a mass program of antiviral chemoprophylaxis in which prophylactic oseltamivir or zanamivir was administered to all persons who had an ill sibling at the camp and to all persons who lived in a cabin with an ill person. Chemoprophylaxis was administered daily by camp staff members to ensure compliance. Over the two sessions, a total of 418 campers and 189 staff members received 10 days of chemoprophylaxis with either oseltamivir (75 mg or appropriate weight-based dosing, once daily) or zanamivir (two 5 mg inhalations, once daily). The camp medical staff continued the program until July 24.

INSIDE

- 972 Receipt of Influenza Vaccine During Pregnancy Among Women With Live Births – Georgia and Rhode Island, 2004–2007
- 975 National Laboratory Inventories for Wild Poliovirus Containment – Western Pacific Region, 2008
- 978 Notice to Readers

The *MMWR* series of publications is published by the Coordinating Center for Health Information and Service, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

Suggested Citation: Centers for Disease Control and Prevention. [Article title]. *MMWR* 2009;58:[inclusive page numbers].

Centers for Disease Control and Prevention

Thomas R. Frieden, MD, MPH

Director

Tanja Popovic, MD, PhD

Chief Science Officer

James W. Stephens, PhD

Associate Director for Science

Steven L. Solomon, MD

Director, Coordinating Center for Health Information and Service

Jay M. Bernhardt, PhD, MPH

Director, National Center for Health Marketing

Katherine L. Daniel, PhD

Deputy Director, National Center for Health Marketing

Editorial and Production Staff

Frederic E. Shaw, MD, JD

Editor, MMWR Series

Christine G. Casey, MD

Deputy Editor, MMWR Series

Robert A. Gunn, MD, MPH

Associate Editor, MMWR Series

Teresa F. Rutledge

Managing Editor, MMWR Series

Douglas W. Weatherwax

Lead Technical Writer-Editor

Donald G. Meadows, MA

Jude C. Rutledge

Writers-Editors

Martha F. Boyd

Lead Visual Information Specialist

Malbea A. LaPete

Stephen R. Spriggs

Terraye M. Starr

Visual Information Specialists

Kim L. Bright

Quang M. Doan, MBA

Phyllis H. King

Information Technology Specialists

Editorial Board

William L. Roper, MD, MPH, Chapel Hill, NC, Chairman

Virginia A. Caine, MD, Indianapolis, IN

Jonathan E. Fielding, MD, MPH, MBA, Los Angeles, CA

David W. Fleming, MD, Seattle, WA

William E. Halperin, MD, DrPH, MPH, Newark, NJ

King K. Holmes, MD, PhD, Seattle, WA

Deborah Holtzman, PhD, Atlanta, GA

John K. Iglehart, Bethesda, MD

Dennis G. Maki, MD, Madison, WI

Sue Mallonee, MPH, Oklahoma City, OK

Patricia Quinlisk, MD, MPH, Des Moines, IA

Patrick L. Remington, MD, MPH, Madison, WI

Barbara K. Rimer, DrPH, Chapel Hill, NC

John V. Rullan, MD, MPH, San Juan, PR

William Schaffner, MD, Nashville, TN

Anne Schuchat, MD, Atlanta, GA

Dixie E. Snider, MD, MPH, Atlanta, GA

John W. Ward, MD, Atlanta, GA

Case Reports

Patient A. One of the campers, a previously healthy adolescent girl, received oseltamivir prophylaxis at an appropriate prophylactic dose of 75 mg daily during June 26–July 5, despite having no reported exposure to an ill person. After completing the initial course of oseltamivir on July 5, she was exposed to an ill cabin mate (patient C) and administered a second 10-day course of chemoprophylaxis at the same dosage beginning on July 7. On July 8, she experienced cough and headache without fever, and on July 9 she experienced chills, worsening headache, and loose stools. Despite these symptoms, her oseltamivir dose was not increased to a therapeutic treatment dose. On July 10, the last day of the first camp session, she traveled away from camp with three family members while ill, returning on July 12, afebrile and with a cough, to attend the second session. On July 12, a rapid influenza detection test was positive for influenza A. The family declined treatment with zanamivir because of concern over side effects, and the patient's oseltamivir dose was doubled to 75 mg, twice daily, an appropriate therapeutic treatment dose. Patient A was isolated with other ill campers and staff members until July 16, and she recovered uneventfully. The camp physician observed that the camper became ill while taking prophylaxis and became concerned that antiviral resistance might have occurred. Therefore, a nasopharyngeal swab specimen was obtained on July 14 and sent to the state laboratory for testing. On July 22, the laboratory confirmed the presence of 2009 pandemic influenza A (H1N1) virus by real-time reverse transcription–polymerase chain reaction (rRT-PCR). On August 19, CDC testing of the same clinical specimen detected the H275Y (3,4) and I223V mutations (6). Because viral isolation was unsuccessful, a neuraminidase inhibition assay was not performed. No illness was reported among her family members.

Patient B. A second previously healthy adolescent girl, who resided in the same cabin as patient A, began oseltamivir chemoprophylaxis at a dose of 75 mg daily on July 7 after exposure to patient C. On July 10, patient B left camp for a home visit during the break between camp sessions. The next day, while at home, she experienced onset of fever (101.9°F [38.8°C]), sore throat, and cough. She continued to engage in normal activities while ill, including visiting a shopping mall and movie theater. She returned to camp for the second session on July 12 with fever, headache, cough, malaise, and myalgias. On July 12, a rapid influenza detection test was positive for influenza A. Oseltamivir was discontinued, and zanamivir treatment (two 5 mg inhalations, twice daily) was begun. A nasopharyngeal swab specimen was obtained July 14 and sent to the state laboratory for testing. On July 20, the presence of 2009 pandemic influenza A (H1N1) virus was

confirmed by rRT-PCR. On August 14, CDC testing of viral RNA detected H275Y and I223V mutations (3,4,6). Viral isolation was unsuccessful, and a neuraminidase inhibition assay was not performed. Patient B was isolated at the camp during July 12–18. Her fever resolved by July 14, and by July 17 she was asymptomatic. No illnesses were identified among close contacts potentially exposed during her weekend home visit.

Further Transmission Investigation

After identification of the oseltamivir-resistant pandemic H1N1 virus, the state health department and local health departments interviewed the families of the two campers and reviewed camp medical records to determine whether the campers might have transmitted virus to others. Retrospective review of camp records revealed that, during June 26–July 22, six other campers were diagnosed with illness while on oseltamivir chemoprophylaxis (75 mg once daily for 10 days). A single specimen from one of these six campers (not patient C) was obtained July 14 and sent to the state laboratory for testing by rRT-PCR, but no influenza virus was isolated. No evidence of pandemic H1N1 virus infection outside the camp linked to either patient A or patient B was found. CDC tested by pyrosequencing 59 specimens of pandemic H1N1 virus, collected during June 29–August 14 as part of routine surveillance conducted by sentinel sites throughout North Carolina. None of the 59 specimens had the H275Y or I223V mutations.

Reported by: *M Garrison, Buncombe County Health Center; L Weldon, Henderson County Dept of Public Health; P Brantley, L Wolf, PhD, M Davies, MD, J-M Maillard, MD, Z Moore, MD, North Carolina Dept of Health and Human Svcs. T Sheu, V Deyde, PhD, L Gubareva, PhD, AM Fry, MD, Influenza Div, National Center for Immunization and Respiratory Diseases; A Fleischauer, PhD, Career Epidemiology Field Officer Program, Coordinating Office for Terrorism Preparedness and Emergency Response; NJ Dailey, MD, EIS Officer, CDC.*

Editorial Note: This report describes confirmed oseltamivir-resistant 2009 pandemic influenza A (H1N1) virus infection in two previously healthy adolescents who were cabin mates and recipients of oseltamivir in a mass chemoprophylaxis program during an outbreak of ILI at a summer camp. This is the first report of oseltamivir resistance in symptomatic close contacts with confirmed infection. Two possible mechanisms of transmission seem most likely. One possibility is that oseltamivir-resistant virus was transmitted from patient A to patient B. The onset of illness for patient B occurred 4 days after the onset of illness for patient A, consistent with reported intervals for secondary transmission among household members with seasonal influenza (7). Alternatively, both patient A and patient B might have acquired oseltamivir-resistant virus infection from exposure to another ill person (e.g., an unknown camper or staff member, or patient C), or each might have developed

oseltamivir resistance independently. Whether the H275Y and I223V mutations occurred independently, or whether virus with one or both of these mutations circulated more widely in the camp could not be determined.

Although six other persons had illness while receiving oseltamivir chemoprophylaxis, aside from the specimens collected from patients A and B, only one specimen was obtained from any other ill person, and the pandemic H1N1 virus was not detected in that specimen. Neither mutation was found in 59 surveillance specimens from sentinel sites in the state, suggesting that the mutations were not widespread in North Carolina. The H275Y mutation has been characterized previously among seasonal influenza A (H1N1) viruses and is associated with resistance to oseltamivir (3). The I223V mutation has not been reported previously in 2009 pandemic influenza A (H1N1); because the neuraminidase inhibition assay could not be performed, the mutation's functional significance is unknown.

These cases highlight a potentially adverse outcome from oseltamivir chemoprophylaxis. In two randomized clinical trials (with 962 and 812 participants, respectively), the efficacy of oseltamivir chemoprophylaxis for preventing clinical seasonal influenza among persons within households ranged from 68% (for laboratory-confirmed infection that included serologic outcomes of infection) to 89% (for laboratory-confirmed clinical influenza) (8,9). No evidence of oseltamivir-resistant virus was reported in these studies. However, the World Health Organization has reported multiple instances of oseltamivir-resistant 2009 pandemic influenza A (H1N1) viruses being isolated from persons who developed pandemic H1N1 infection while receiving oseltamivir chemoprophylaxis (2). Resistance to oseltamivir also might develop during subtherapeutic dosing. In this report, patient A was symptomatic while on a chemoprophylaxis dose of oseltamivir for 4 days. One possibility is that she developed resistance while on a subtherapeutic dosage of 75 mg once a day for chemoprophylaxis, rather than the appropriate treatment dose of 75 mg twice a day.

CDC recommendations regarding use of antivirals during the H1N1 pandemic were updated on September 8. Use of antiviral medications for postexposure chemoprophylaxis should be reserved for persons at higher risk for influenza-related complications who have had contact with someone likely to have been infected with influenza. An emphasis on early treatment once a patient has developed symptoms, rather than chemoprophylaxis, should reduce opportunities for development of oseltamivir resistance (5). Chemoprophylaxis should not be used for prevention of illness among healthy persons after exposures in community settings. Persons who are taking antiviral medications for prevention should be instructed to contact a health-care provider if illness develops. Persons under antiviral treatment should be instructed to contact a

health-care provider if symptoms worsen. Other preventative measures (e.g., hand hygiene and cough etiquette) can reduce the risk for influenza virus transmission (5).

Chemoprophylaxis failure is known to occur even without antiviral resistance (8,9). Accordingly, not all failures need to be accompanied by testing for resistance; testing should be considered for individual cases in consultation with the state health department. However, if symptoms develop during chemoprophylaxis, providers should consider the possibility of antiviral resistance and consider alternate treatment options. Because the 2009 pandemic influenza A (H1N1) virus is resistant to adamantanes (1), limited treatment options will be available if widespread oseltamivir resistance develops. Zanamivir is not licensed for treatment of children aged <7 years and is contraindicated among persons with underlying airway disease.

References

1. CDC. Drug susceptibility of swine-origin influenza A (H1N1) viruses, April 2009. *MMWR* 2009;58:433–5.
2. World Health Organization. Pandemic (H1N1) 2009: update 60. Geneva, Switzerland: World Health Organization; 2009. Available at http://www.who.int/csr/don/2009_08_04/en/index.html.
3. Deyde VM, Gubareva LV. Influenza genome analysis using pyrosequencing method: current applications for a moving target. *Expert Rev Mol Diagn* 2009;9:493–509.
4. World Health Organization. Influenza A (H1N1) NA-H274 detailed pyrosequencing protocol for antiviral susceptibility testing. Geneva, Switzerland: World Health Organization; 2009. Available at http://www.who.int/csr/resources/publications/swineflu/NA_DetailedPyrosequencing_20090513.pdf.
5. CDC. Updated interim recommendations for the use of antiviral medications in the treatment and prevention of influenza for the 2009–2010 season. Atlanta, GA: CDC; September 8, 2009. Available at <http://www.cdc.gov/h1n1flu/recommendations.htm>.
6. Garten RJ, Davis CT, Russell CA, et al. Antigenic and genetic characteristics of swine-origin 2009 A(H1N1) influenza viruses circulating in humans. *Science* 2009;325:197–201.
7. Cowling BJ, Fang VJ, Riley S, Malik Peiris JS, Leung GM. Estimation of the serial interval of influenza. *Epidemiology* 2009;20:344–7.
8. Welliver R, Monto AS, Carewicz O, et al. Effectiveness of oseltamivir in preventing influenza in household contacts: a randomized controlled trial. *JAMA* 2001;285:748–54.
9. Hayden FG, Belshe R, Villanueva C, et al. Management of influenza in households: a prospective, randomized comparison of oseltamivir treatment with or without postexposure prophylaxis. *J Infect Dis* 2004;189:440–9.

Receipt of Influenza Vaccine During Pregnancy Among Women With Live Births – Georgia and Rhode Island, 2004–2007

Pregnant women are at increased risk for complications from influenza (1–3). Since 2004, the Advisory Committee on Immunization Practices (ACIP) and American College of Obstetricians and Gynecologists (ACOG) Committee on Obstetric Practice have recommended that all pregnant women be vaccinated with the trivalent inactivated vaccine during any trimester of pregnancy (4,5). To assess the percentage of women who were vaccinated during pregnancy among women with recent live births, CDC analyzed data from the Pregnancy Risk Assessment and Monitoring System (PRAMS) from Georgia and Rhode Island, the two states that collected this information on the PRAMS survey. This report summarizes the results, which showed that in Georgia, the prevalence of influenza vaccination during the woman's most recent pregnancy increased from 10.4% in 2004 to 15.5% in 2006. In Rhode Island, vaccination prevalence increased from 21.9% in 2004 to 33.4% in 2007. During 2006 in Georgia, the most common reasons for not receiving vaccination were, "I don't normally get the flu vaccination" (69.4%), and "my physician did not mention anything about a flu vaccine during my pregnancy" (44.5%). Increased efforts are needed to assess vaccine coverage during pregnancy and to educate providers and pregnant women about ACIP and ACOG recommendations on providing intramuscular, inactivated influenza vaccine during any trimester of pregnancy.

PRAMS is a population-based surveillance system that collects data on a wide range of maternal behaviors and experiences before, during, and after pregnancy. PRAMS surveys currently are administered by 37 states, New York City, and one tribal-state partnership in South Dakota. Each month, participating states or entities use birth certificate data to select a stratified random sample of 100–300 women with recent live births. A questionnaire is mailed to the women 2–6 months after delivery. The participating entities use a standard questionnaire, to which they can add questions. From 2004 to 2007, Georgia and Rhode Island included questions about influenza vaccination on their surveys. Responses from Georgia for 3 years (2004–2006; N = 5,231) and Rhode Island for 4 years (2004–2007; N = 5,499) were analyzed. Variables included receipt of influenza vaccination in women during pregnancy, demographics, and health-care service use indicators. Response rates for the years of data examined for Georgia were 70.0% for 2004, 70.2% for 2005, and 70.8% for 2006; rates for


Rhode Island were 75.5% for 2004, 75.1% for 2005, 72.5% for 2006, and 72.1% for 2007.

Women whose influenza vaccination status was missing (229 for Georgia and 163 for Rhode Island) were excluded. PRAMS data were weighted to take into account complex survey design, nonresponse, and noncoverage for each state. Data were analyzed to estimate influenza vaccination prevalence and 95% confidence intervals. Chi-square tests were used to determine statistical significance and statistical software were used to account for the complex sampling strategy.

Surveys conducted by both states inquired about influenza vaccination coverage by asking the question, "Did you get a flu vaccination during your most recent pregnancy?" The Georgia survey included a follow-up question for women who reported not being vaccinated to assess their reasons. The question included was, "What were your reasons for not getting a flu vaccination during your most recent pregnancy?" A list of reasons with a choice of yes/no response included items on receipt of provider advice, perceptions of vaccine safety, and timing of pregnancy. The Rhode Island survey included a question on provider advice, "At any time during your pregnancy, did a doctor, nurse, or other health-care worker offer you a flu vaccination or tell you to get one?"

In both states, most of the increase in influenza vaccination coverage was observed from 2004 to 2005 (Figure); in Georgia, coverage increased 62.5%, from 10.4% to 16.9%, and in Rhode Island, coverage increased 37.4%, from 21.9% to 30.1%. Vaccination prevalence remained mostly stable during 2005–2006, but with a further 10.0% increase observed in Rhode Island from 2006 to 2007. Prevalence of influenza

FIGURE. Influenza vaccination coverage during most recent pregnancy among women with recent live births* — Pregnancy Risk Assessment and Monitoring System, Georgia and Rhode Island, 2004–2007



* Based on response to the question, "Did you get a flu vaccination during your most recent pregnancy?"

† 2007 data for Georgia were not available. Percentages are weighted.

§ 95% confidence interval.

vaccination during pregnancy in the two states varied by state and demographically (Table 1).

In Rhode Island, the prevalence of women who reported receiving advice about influenza vaccine or an offer of vaccination increased from 33.0% during 2004 to 47.7% during 2007 ($p < 0.001$). In 2007, among respondents who reported receiving vaccination advice from a health-care provider, the prevalence of those who also were vaccinated was 65.7%. In 2007, Rhode Island data showed that among women who did not report receiving advice from their health-care provider about influenza vaccine, only 4.6% reported receiving influenza vaccination.

In Georgia, previous vaccination history, provider advice, perceptions of safety, and timing of pregnancy were among the reasons unvaccinated women cited for not getting the influenza vaccine (Table 2). The top reasons cited were "I don't normally

TABLE 1. Influenza vaccination coverage during most recent pregnancy among women with recent live births, by selected characteristics — Pregnancy Risk Assessment Monitoring System, Georgia and Rhode Island, 2006 and 2007.

Characteristic	Georgia 2006 (n = 1,958*)		Rhode Island 2007 (n = 1,328*)	
	%†	(95% CI)§	%	(95% CI)
Total	15.5	(13.2–17.8)	33.4	(30.4–36.4)
Maternal race/ethnicity				
Black, non-Hispanic	12.6	(10.0–15.2)	29.8	(19.9–39.7)
White, non-Hispanic	16.9	(13.3–20.6)	30.4	(26.7–34.1)
Hispanic	18.9	(10.8–27.0)	42.5	(36.1–49.0)
Other	12.0	(0.1–23.9)¶	35.5	(23.0–48.0)
Maternal age (yrs)				
<20	20.3	(12.9–27.7)	29.2	(19.6–38.8)
20–24	11.5	(7.8–15.4)	38.1	(31.2–45.0)
25–29	12.4	(8.3–16.5)	31.0	(25.4–36.6)
30–34	21.6	(15.9–27.3)	34.0	(28.0–40.0)
≥35	15.9	(9.2–22.6)	33.0	(26.0–40.0)
Marital status				
Married	15.9	(12.8–19.1)	35.3	(31.3–39.3)
Not married	15.0	(11.5–18.6)	30.7	(26.2–35.2)
Maternal education				
<High school	11.5	(5.4–17.6)	36.1	(28.4–43.8)
High school	11.7	(8.1–15.4)	27.8	(22.5–33.1)
>High school	17.2	(13.7–20.8)	36.1	(31.9–40.4)
Parity				
1	18.2	(14.3–22.1)	29.3	(25.0–33.6)
2	15.2	(11.1–19.3)	36.1	(30.7–41.6)
≥3	11.8	(7.8–15.8)	34.6	(28.1–41.1)
Medicaid paid for prenatal care				
Yes	12.3	(9.4–15.2)	39.9	(31.8–48.0)
No	18.5	(14.9–22.1)	32.5	(29.3–35.8)
Prenatal care initiation				
1st trimester	15.8	(13.1–18.5)	33.6	(30.3–36.9)
After 1st trimester	15.0	(9.8–20.2)	27.3	(19.6–35.0)

* Sample sizes are unweighted.

† Percentages are weighted.

§ Confidence interval.

¶ <60 respondents; estimates might not be stable.

TABLE 2. Reasons given by women with recent live births for not receiving influenza vaccination among those who were not vaccinated (n = 1,648*) — Pregnancy Risk Assessment Monitoring System, Georgia, 2006

Reason†	%‡	(95% CI¶)
I don't normally get a flu vaccination	69.4	(66.2–72.7)
My physician did not mention anything about a flu vaccine during my pregnancy	44.5	(40.9–48.0)
I was worried that the flu vaccination might harm my baby	28.1	(24.9–31.3)
I was worried about side effects of the flu vaccination for me	27.1	(23.9–30.3)
I was in my first trimester during the flu season (November–February)	25.2	(22.0–28.4)
I wasn't pregnant during the flu season (November–February)	24.2	(21.1–27.3)
Other reason	6.2	(4.3–8.0)

* Unweighted sample sizes.

† Respondents could select more than one reason.

‡ Percentages are weighted.

¶ Confidence interval.

get the flu vaccination" (69.4%) and "my physician did not mention anything about a flu vaccine during my pregnancy" (44.5%); 28.1% were worried about the safety of the influenza vaccine for their infant and 27.1% were worried about the safety for themselves.

Reported by: *IB Ahluwalia, PhD, L Harrison, MPH, D Jamieson, MD, Div of Reproductive Health, National Center on Chronic Disease Prevention and Health Promotion; SA Rasmussen, MD, Div of Birth Defects and Developmental Disabilities, National Center on Birth Defects and Developmental Disabilities, CDC.*

Editorial Note: Despite evidence that maternal vaccination with influenza vaccine protects infants from influenza-like illness during the first 6 months of life (6), recent national data show that pregnant women have the lowest rates of coverage among all adult populations recommended to receive influenza vaccination (7). During 2004–2007, influenza vaccination prevalence increased significantly in both states among women with recent live births. The increases in coverage partially could be related to changes in ACIP and ACOG recommendations in 2004, when the recommendation for pregnant women changed from vaccine administration to women who would be in their second or third trimester during influenza season to administration of vaccine any time during pregnancy (4,5). Also, increased media reporting about vaccination of high-risk populations during the 2004–2005 influenza vaccine shortage might have increased awareness among pregnant women and their providers, perhaps resulting in an increase in influenza vaccination prevalence (8).

Interventions focusing on providers and pregnant women might address barriers to influenza vaccination experienced by both (3,9). In July 2006, Rhode Island passed a law* requiring the Rhode Island health department to purchase vaccine and distribute it to physicians who enroll in the Immunize for Life

adult immunization program.† By enacting specific legislation, Rhode Island increased vaccine availability for health-care providers and perhaps alerted providers and pregnant women about the importance of immunizing pregnant women with influenza vaccine. The Rhode Island experience with vaccine distribution might be a useful example for other states on the effectiveness of working with health-care providers to supply influenza vaccine for pregnant women.

Approximately 25% of women in Georgia cited being in their first trimester as a reason for not getting the influenza vaccine, identifying a need to educate pregnant women about the importance of getting the influenza vaccine, even during the earliest phases of pregnancy. These women and their physicians might not have been aware of current ACIP and ACOG recommendations that women who are pregnant during influenza season should be vaccinated, irrespective of trimester. Providers should be educated about these recommendations, about influenza risks for pregnant women, and about interventions to prevent severe illness in this population (3–5,9).

The findings in this report are subject to at least three limitations. First, PRAMS data on influenza vaccination were only available from two states, and these findings might not be generalizable to all women with live births in the United States. Second, PRAMS data are self-reported by women 2–4 months postpartum and therefore they might be subject to recall bias. Finally, information on provider recommendations was assessed by maternal report; data from health-care providers regarding their practice related to influenza vaccine might have shown different results.

Because the seasonal influenza vaccine is unlikely to provide protection against pandemic influenza A (H1N1) infection (10), ACIP recommends that pregnant women receive both vaccine formulations during the 2009–10 influenza season. The

* Routine childhood and adult immunization vaccines. Title 23, Chapter 23-1, Sect. 23-1-44 (2006). Available at <http://www.rilin.state.ri.us/statutes/title23/23-1/23-1-44.htm>.

† Additional information available at <http://www.health.ri.gov/immunization/immunizeforlife.php>.

trivalent inactivated seasonal influenza vaccine is now available and the influenza A (H1N1) 2009 monovalent vaccine is expected to become available in mid-October (10).

Acknowledgments

This report is based, in part, on contributions by H Kim, PhD, and R Cain, Rhode Island Dept of Health Center for Health Data and Analysis; C Hoban, PhD, and D Goodman, PhD, Georgia Dept of Community Health, Office of Epidemiology, Evaluation, and Health Information; A Harrison, Science Applications International Corporation Contractor; and S Gupta, MPH, Div of Reproductive Health, National Center for Chronic Disease Prevention and Health Promotion, CDC.

References

1. Jamieson DJ, Theiler RN, Rasmussen SA. Emerging infections and pregnancy. *Emerg Infect Dis* 2006;12:1638–43.
2. Rasmussen SA, Jamieson DJ, Bresee JS. Pandemic influenza and pregnant women. *Emerg Infect Dis* 2008;14:95–100.
3. Naleway AL, Smith WJ, Mullooly JP. Delivering influenza vaccine to pregnant women. *Epidemiol Rev* 2006;28:47–53.
4. CDC. Prevention and control of influenza: recommendations of the Advisory Committee on Immunization Practices (ACIP), 2008. *MMWR* 2008;57(No. RR-7).
5. American College of Obstetricians and Gynecologists' Committee on Obstetric Practice. ACOG committee opinion number 305, November 2004. Influenza vaccination and treatment during pregnancy. *Obstet Gynecol* 2004;104:1125–6.
6. Zaman K, Roy E, Arifeen SE, Rahman M, et al. Effectiveness of maternal influenza immunization in mothers and infants. *N Engl J Med* 2008;359:1555–64.
7. Lu P, Bridges CB, Euler GL, Singleton JA. Influenza vaccination of recommended adult populations, U.S., 1989–2005. *Vaccine* 2008;26:1786–93.
8. Brewer NT, Hallman WK. Subjective and objective risk as predictors of influenza vaccination during the vaccine shortage of 2004–2005. *Clin Infect Dis* 2006;43:1379–86.
9. CDC. Influenza vaccination in pregnancy: practices among obstetrician-gynecologists—United States, 2003–04 influenza season. *MMWR* 2005;54:1050–2.
10. CDC. Use of influenza A (H1N1) 2009 monovalent vaccine: recommendations of the Advisory Committee on Immunization Practices (ACIP), 2009. *MMWR* 2009;58(No. RR-10).

National Laboratory Inventories for Wild Poliovirus Containment — Western Pacific Region, 2008

In the future, when wild poliovirus (WPV) transmission is interrupted worldwide, facilities holding WPV materials will represent the only remaining repository of the virus. Maintaining the number of such facilities at a minimum and at an appropriate biosafety standard (laboratory containment) reduces the risk for a facility-associated reintroduction of WPV. In May 1999, the World Health Assembly (WHA) urged

all member states to begin the process leading to laboratory containment of WPV (1). The World Health Organization (WHO) global action plan for laboratory containment of WPV issued in 1999 indicated a staged approach that begins with a national survey of all biomedical facilities (Phase I); the purpose of the survey is to alert institutions and facilities to the need for containment, encourage reduction of WPV materials, and develop a national inventory of facilities holding such materials. The survey and inventory provide a facility database for use in all subsequent steps toward global poliovirus containment. In May 2008, WHA urged all WHO member states to complete Phase I activities outlined in the *WHO Global Action Plan for Laboratory Containment of Wild Polioviruses* (2,3). In the WHO Western Pacific Region (WPR), Phase I surveys of 77,260 laboratories in the 37 countries and areas of WPR were conducted during 1999–2008. A total of 45 laboratories were identified as holding WPV materials in 2008. This report describes completion of Phase I containment activities by WPR countries, and updates a previous report on Phase I completion in the European Region and global progress (4).

Specific guidelines for Phase I activities in the WPR were issued in 1999 that advised member states and areas* to conduct national surveys of biomedical facilities,† communicate the need for WPV containment, develop a database of facilities, and compile a national inventory of laboratories identified to possess WPV infectious or potentially infectious materials (WPV materials).§ Facilities were to be advised to dispose of unneeded WPV materials and to handle retained WPV materials under appropriate biosafety conditions. In 1999, the Regional Commission for the Certification of Poliomyelitis Eradication in the Western Pacific (WPR RCC) established progress toward completion of Phase I containment as a requirement for certification of the region as polio-free (5).

* American Samoa, Australia, Brunei Darussalam, Cambodia, China, Commonwealth of the Cook Islands, Fiji, French Polynesia, Guam, Hong Kong (China), Japan, Kiribati, Lao People's Democratic Republic, Macao (China), Malaysia, Marshall Islands, Micronesia, Mongolia, Nauru, New Caledonia, New Zealand, Niue, Northern Mariana Islands, Palau, Papua New Guinea, Philippines, Pitcairn Islands, Republic of Korea, Samoa, Singapore, Solomon Islands, Republic of Tokelau, Tonga, Tuvalu, Vanuatu, Vietnam, and Federated States of Wallis and Futuna.

† Facilities holding WPV infectious or potentially infectious materials (WPV materials) include diagnostic laboratories with frozen stool specimens, institutions with current or past research on polioviruses, teaching and industrial facilities that use poliovirus as a test virus, and vaccine manufacturers.

§ Infectious materials include clinical materials from persons with confirmed WPV infections; environmental sewage, or water samples in which WPV is present; and replication products containing WPV. Potentially infectious materials include feces, respiratory secretions, environmental sewage, and untreated water samples of unknown origin or collected for any purpose at a time and in a geographic area where presence of WPVs was suspected, and the replication products of such materials. Replication products include cell culture isolates, reference stocks, and laboratory derivatives in poliovirus-permissive cells or animals. For the purposes of containment, vaccine-derived poliovirus materials are treated as equivalent to WPV materials (3).

Preliminary information on Phase I activities was reported by each member state to the WHO regional office annually. When WPR was certified as polio free in October 2000, all member states/areas had initiated Phase I, but only four member states/areas had completed it (Figure).

Strategies for identifying and surveying biomedical facilities differed among WPR countries according to population size, administrative and health infrastructure, economic development, and political structure. In small member states (e.g., Brunei Darussalam and Macao [China]), laboratory surveys were easier to complete by the small number of facilities, the majority of which were under government jurisdiction. In member states with less developed laboratory infrastructures (e.g., Cambodia and the 21 Pacific island countries and areas), health staff at district levels identified the facilities in the country and assessed freezer capacity and power supply to exclude facilities not capable of preserving polioviruses. Containment officials at the national level then focused survey efforts on the facilities assumed to be at higher risk for retaining WPV materials (i.e., those with research or teaching functions). In member states with more developed laboratory infrastructures (e.g., Australia and Republic of Korea), the database of facilities was compiled from preexisting lists of licensed laboratories supplemented with member lists of professional institution associations (e.g., biosafety associations and microbiological societies), and lists for laboratory accreditation or quality control. In the majority of countries, the surveys were completed by calling or visiting nonresponding laboratories. Completeness of the surveys was assessed by a systematic quality-assurance procedure provided by WHO.

By 2006, Phase I was complete in all countries except China and Japan, which have vast biomedical laboratory infrastructures. During 2000–2003, Phase I surveys in China were conducted in facilities operating under the jurisdiction of five Chinese ministries/agencies.[‡] During 2005–2008, a comprehensive approach was initiated which included 1) surveying nearly 50,000 biomedical laboratories under the jurisdiction of the Ministry of Health identified by compiling information at the county, prefecture, and provincial levels; 2) performing a pilot survey in selected provinces of biomedical facilities among 46 ministries and agencies other than the Ministry of Health; and 3) expanding surveys to all provinces and ministries based on the lessons learned from the pilot. Survey completion was facilitated by regulations on safe handling of pathogenic agents issued after confirmed laboratory-acquired SARS infections during the 2003 epidemic (6,7).

In Japan, a nationwide survey was implemented during 2000–2002 by the Ministry of Health, Labor, and Welfare (MHLW) covering 7,865 facilities with an overall return rate of 53.8%. An expanded survey conducted during 2004–2005 achieved a response rate of >99% from the 12,142 facilities under the jurisdiction of the MHLW and 1,367 facilities under the Ministry of Education, Culture, Sport, Science, and Technology, the two ministries overseeing facilities considered most likely to have WPV materials. A further total of 560 facilities were surveyed under the jurisdiction of the remaining ministries. The number of biomedical laboratories surveyed within each facility was not reported. An additional targeted survey of 80 high-risk facilities identified through a search of published poliovirus research was conducted in 2008 to further validate the earlier surveys. Among the 88 laboratories in these facilities, 82 (93%) had been surveyed previously; one laboratory of the six that were not surveyed previously reported holding WPV materials.**

In the WPR, a total of 77,260 biomedical facilities responded to the Phase I surveys for all countries, including 55,688 facilities in China and 14,069 facilities in Japan (Table). Of all biomedical facilities responding to the surveys, 89% (68,831) were clinical diagnostic laboratories, primarily under the jurisdiction of ministries of health; only 32 (0.05%) of these were found to hold WPV materials, 27 of which are laboratories within the Global Polio Laboratory Network (one in Australia, 25 in China, and one in Japan). Of the regional total, 3,838 facilities were listed as being at high risk for retaining WPV materials, 94% of which were located in Australia, China, Japan, and Republic of Korea (Table); 11 (0.3%) of these reported having WPV materials.†† Any facility that had reported retention of WPV materials in a given survey was resurveyed annually. The number of facilities/laboratories retaining WPV materials decreased during the course of Phase I implementation, resulting in a decline from 147 facilities in 1999 to 107 in 2006 to 45 in 2008 (two in Australia, 27 in China, 15 in Japan, and one in the Republic of Korea) (Figure).


All member states documented laboratory containment activities through standardized reports reviewed by the WHO regional office, a panel of experts external to the process, and the RCC. For the two member states with the most complex laboratory infrastructure (China and Japan), the external review of the process included site visits to critical academic and research institutions. In December 2008, the WPR RCC accepted the final reports from China and Japan and

[‡] Ministry of Health, Ministry of Education, State Environmental Protection Administration, State Drug Administration, and Chinese Academy of Science.

** Another facility was added to the national inventory after the survey, when newly requested reference strains were transferred from Japan's National Institute of Infectious Diseases.

†† The other two facilities holding WPV materials were regulatory and production facilities.

FIGURE. Number of biomedical facilities reporting wild poliovirus (WPV) materials* and number of World Health Organization (WHO) member states and areas† having completed Phase I of the WPV containment process, by year — WHO Western Pacific Region, 1999–2008



* WPV infectious and potentially infectious materials. Additional information available at <http://www.polioeradication.org/content/publications/who-vb-03-729.pdf>.

† Of the 37 member states/areas, the 21 Pacific island countries and areas are presented as a block and include American Samoa, Commonwealth of the Cook Islands, Fiji, French Polynesia, Guam, Kiribati, Marshall Islands, Micronesia, Nauru, New Caledonia, Niue, Northern Mariana Islands, Palau, Pitcairn Islands, Samoa, Solomon Islands, Republic of Tokelau, Tonga, Tuvalu, Vanuatu, and Federated States of Wallis and Futuna. The 16 other member states and areas are Australia, Brunei Darussalam, Cambodia, China, Hong Kong (China), Japan, Lao People's Democratic Republic, Macao (China), Malaysia, Mongolia, New Zealand, Papua New Guinea, Philippines, Republic of Korea, Singapore, and Vietnam.

declared Phase I WPV laboratory containment complete for the region.

Reported by: *National containment coordinators, Western Pacific Region member states; B Bayutas, S Roesel, World Health Organization Western Pacific Regional Office, Manila, Philippines. C Wolff, Polio Eradication Dept, World Health Organization, Geneva, Switzerland. Task Force for Global Health, Decatur, Georgia. Global Immunization Div, Div of Viral Diseases, National Center for Immunization and Respiratory Diseases, CDC.*

Editorial Note: WPR has joined the WHO European Region in completing Phase I WPV laboratory containment activities (4). The WHO Region of the Americas (AMR), the remaining polio-free region, did not initiate containment activities at the time of certification in 1994 because the policies were not yet developed. The AMR RCC has accepted the final Phase I reports from 24 (69%) of 35 member states, including the United States. The other 11 AMR countries have reported completion of Phase I and are anticipated to submit final reports by the end of 2009. In the three polio-endemic WHO regions (8),^{§§} 43 (62%) of 69 polio-free countries and areas have completed Phase I activities.

Multiple challenges were faced in the 10 years required to complete Phase I in the WPR. Prioritization for WPV

containment activities weakened in the ministries of health of many countries after regional certification. China and Japan had to access laboratories under the jurisdiction of a wide range of government agencies in addition to the ministries of health. During 2001–2006, identification of vaccine-derived polioviruses (9), which are considered equivalent as WPV for containment purposes, required several WPR countries to resurvey some facilities.

Despite the challenges, successful completion of Phase I activities is achievable even in countries with highly complex laboratory infrastructures. The only polio-free countries with comparable laboratory infrastructures remaining to complete Phase I activities are Egypt and South Africa. India, which remains polio-endemic in 2009, also has a complex laboratory infrastructure and will require similar efforts. Tangential benefits of Phase I were noted in the WPR and other regions.

Authorities of many member states found that the national survey and inventory process led to a better understanding of the laboratory infrastructure of the country, a strengthened process for laboratory registration, and an increased awareness of the importance of maintaining biosafety standards.

A major accomplishment in the WPR was a progressive voluntary reduction in number of facilities retaining WPV from 147 provisionally reported in 1999 to 45 in 2008, certified by the ministries of health. Authorities in the four relevant WPR countries have indicated the intention to reduce further the number of facilities holding WPV materials.

Subsequent phases of WPV containment are outlined in the newly developed *WHO Global Action Plan to Minimize Poliovirus Facility-Associated Risk in the Post-Eradication/Post-OPV Era* (10). A revised draft edition will be available for public review and comment before the end of 2009.^{¶¶} This action plan includes containment of oral poliovirus vaccine (OPV) Sabin strains as well, and establishes the goal of reducing the number of facilities holding WPV worldwide to <20 in the post-eradication/post-OPV era, including vaccine manufacturers. Nonviable poliovirus reagents can replace live polioviruses in national surveillance and diagnostic facilities. Phase II of the action plan begins after evident interruption of WPV

^{§§} African Region, Eastern Mediterranean Region, and South-East Asia Region.

^{¶¶} Additional information available at <http://www.polioeradication.org>.

TABLE. Number of biomedical facilities surveyed for the presence of wild poliovirus (WPV) materials,* by country/area and type of facility, 1999–2008, and number retaining WPV materials in 2008 — World Health Organization Western Pacific Region

Country/Area	Diagnostic	Teaching/ Research	Industrial	Other/Unknown	Total	Retaining WPV materials
Australia	2,026	197	0	85	2,308	2 [†]
Brunei Darussalam	12	2	0	0	14	
Cambodia	0	10	0	233	243	
China	52,502	1,704	379	1,103	55,688	27 [†]
Hong Kong (China)	190	19	47	0	256	
Japan	10,865	1,280	1,285	639	14,069	15 [†]
Lao People's Democratic Republic	23	2	0	1	26	
Macao (China)	9	3	1	1	14	
Malaysia	431	43	0	81	555	
Mongolia	53	6	0	12	71	
New Zealand	60	44	3	2	109	
Pacific island countries and areas [§]	17	4	0	6	27	
Papua New Guinea	19	2	0	5	26	
Philippines	2,114	40	0	609	2,763	
Republic of Korea	299	427	37	4	767	1
Singapore	83	43	19	38	183	
Vietnam	128	12	1	0	141	
Total	68,831	3,838	1,772	2,819	77,260	45[†]

* WPV infectious and potentially infectious materials. Additional information available at <http://www.polioeradication.org/content/publications/who-vb-03-729.pdf>.

[†] Includes 27 laboratories in the Global Polio Laboratory Network: Australia (one), China (25), and Japan (one).

[§] The 21 Pacific island countries and areas are presented as a block and include American Samoa, Commonwealth of the Cook Islands, Fiji, French Polynesia, Guam, Kiribati, Marshall Islands, Micronesia, Nauru, New Caledonia, Niue, Northern Mariana Islands, Palau, Pitcairn Islands, Samoa, Solomon Islands, Republic of Tokelau, Tonga, Tuvalu, Vanuatu, and Federated States of Wallis and Futuna.

transmission in one of the four remaining endemic countries, during which member states are requested to establish long-term national policies and regulations for destruction and/or containment of WPV materials. Completion of Phase I in all countries of two WHO regions and the majority of countries in the other four regions, as of the end of 2008, provides a solid base for subsequent polio containment phases.

References

1. World Health Assembly. 52nd session resolution WHA52.8. Eradication of poliomyelitis. Geneva, Switzerland: World Health Organization; 1999. Available at http://apps.who.int/gb/archive/pdf_files/wha52/ew8.pdf.
2. World Health Assembly. 61st session resolution WHA61.1. Poliomyelitis: mechanism for management of potential risks to eradication. Geneva, Switzerland: World Health Organization; 2008. Available at http://apps.who.int/gb/ebwha/pdf_files/wha61-rec1/a61_rec1-part2-en.pdf.
3. World Health Organization. WHO global action plan for laboratory containment of wild polioviruses. 2nd edition. Geneva, Switzerland: World Health Organization; 2004. Available at <http://www.polioeradication.org/content/publications/who-vb-03-729.pdf>.
4. CDC. National laboratory inventory for global poliovirus containment—European Region, June 2006. *MMWR* 2006;55:916–8.
5. World Health Organization Regional Committee for the Western Pacific. 50th session resolution WPR/RC50.R2. Eradication of poliomyelitis in the region. Manila, Philippines: World Health Organization Regional Committee for the Western Pacific; 1999. Available at http://www.wpro.who.int/rcm/en/archives/rc50/rc_resolutions/wpr_rc50_r02.htm.
6. Wang M, Du L, Zhou DH, et al. [Study on the epidemiology and measures for control on severe acute respiratory syndrome in Guangzhou city][Chinese]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2003;24:353–7.
7. Wilder-Smith A, Low JG. Risk of respiratory infections in health care workers: lessons on infection control emerge from the SARS outbreak. *Southeast Asian J Trop Med Public Health*. 2005;36:481–8.

8. CDC. Progress toward interruption of wild poliovirus transmission—worldwide, 2008. *MMWR* 2009;58:308–12.
9. CDC. Update on vaccine-derived polioviruses. *MMWR* 2006;55:1093–7.
10. World Health Organization. WHO global action plan to minimize poliovirus facility-associated risk in the post-eradication/post-OPV era [draft]. Geneva, Switzerland: World Health Organization; 2009. Available at http://www.polioeradication.org/content/publications/gapiiworkingdraft_07.pdf.

Notice to Readers

National Child Passenger Safety Week — September 12–18, 2009

In 2007, a total of 606 children aged <8 years died and approximately 75,000 were treated in emergency departments for occupant injuries sustained in motor-vehicle crashes in the United States (1,2). National Child Passenger Safety Week, September 12–18, 2009, highlights the importance of the correct installation and use of child restraints.

The use of booster seats has been found to reduce the risk for injury by 59% in children aged 4–7 years, compared with use of adult seat belts alone (3). The National Highway Traffic Safety Administration (NHTSA) and CDC recommend placing infants and children in age-, weight-, and height-appropriate child restraints until they are aged ≥8 years or are

57 inches tall, at which time they can use adult seat belts (4,5). Although no recent data are available on consistent compliance with this recommendation during a specified period, an older study, CDC's Second Injury Control and Risk Survey (ICARIS-2), a national, cross-sectional, random-digit-dial telephone survey conducted July 2001–February 2003, found that 46% of parents of children aged 4–7 years reported their children had used adult seat belts all of the time during the preceding 30 days (6).

Although the use of child restraints is mandatory in all 50 states and the District of Columbia, the age at which children can transition to adult safety belts varies by state. Twenty-three states allow children to use adult seat belts by age ≤ 7 years, with four states allowing adult seat belt use for children at age 5 years and one state allowing adult seat belt use for children aged 4 years (7).

Information about National Child Passenger Safety Week activities and child passenger safety is available from NHTSA at <http://www.nhtsa.gov/cps> and from CDC at http://www.cdc.gov/motorvehiclesafety/child_passenger_safety/childseat-spot.html.

References

1. National Highway Traffic Safety Administration. Fatality Analysis Reporting System (FARS) encyclopedia. Available at <http://www-fars.nhtsa.dot.gov/main/index.aspx>.
2. CDC. WISQARS nonfatal injury reports. Available at <http://webappa.cdc.gov/sasweb/ncipc/nfirates.html>.
3. Durbin DR, Elliott MR, Winston FK. Belt-positioning booster seats and reduction in risk of injury among children in vehicle crashes. *JAMA* 2003;28:2835–40.
4. National Highway Traffic Safety Administration. Child passenger safety. A parent's primer. Available at <http://www.nhtsa.dot.gov/staticfiles/DOT/NHTSA/Traffic%20Injury%20Control/Articles/Associated%20Files/4StepsFlyer.pdf>.
5. CDC. Child passenger safety fact sheet. Available at http://www.cdc.gov/motorvehiclesafety/child_passenger_safety/cps-factsheet.html.
6. CDC. Second injury control and risk survey (ICARIS-2). Available at <http://www.cdc.gov/ncipc/osp/icaris2.htm>.
7. Insurance Institute for Highway Safety, Highway Loss Data Institute. Child restraint/belt laws. Available at http://www.iihs.org/laws/state_laws/restrain.html.

TABLE I. Provisional cases of infrequently reported notifiable diseases (<1,000 cases reported during the preceding year) — United States, week ending September 5, 2009 (35th week)*


Disease	Current week	Cum 2009	5-year weekly average†	Total cases reported for previous years					States reporting cases during current week (No.)
				2008	2007	2006	2005	2004	
Anthrax	—	—	0	—	1	1	—	—	
Botulism:									
foodborne	—	12	1	17	32	20	19	16	
infant	—	31	2	109	85	97	85	87	
other (wound and unspecified)	1	17	1	19	27	48	31	30	CA (1)
Brucellosis	3	63	2	80	131	121	120	114	PA (1), FL (1), CA (1)
Chancroid	—	19	0	25	23	33	17	30	
Cholera	—	4	0	5	7	9	8	6	
Cyclosporiasis§	2	99	3	139	93	137	543	160	FL (2)
Diphtheria	—	—	—	—	—	—	—	—	
Domestic arboviral diseases§,¶:									
California serogroup	—	9	5	62	55	67	80	112	
eastern equine	—	1	1	4	4	8	21	6	
Powassan	—	1	0	2	7	1	1	1	
St. Louis	—	7	1	13	9	10	13	12	
western equine	—	—	—	—	—	—	—	—	
Ehrlichiosis/Anaplasmosis§,**:									
<i>Ehrlichia chaffeensis</i>	3	473	20	1,137	828	578	506	338	NY (1), SC (1), CA (1)
<i>Ehrlichia ewingii</i>	—	3	0	9	—	—	—	—	
<i>Anaplasma phagocytophilum</i>	6	341	17	1,026	834	646	786	537	NY (5), NE (1)
undetermined	—	75	5	180	337	231	112	59	
<i>Haemophilus influenzae</i> ††									
invasive disease (age <5 yrs):									
serotype b	—	16	0	30	22	29	9	19	
nonserotype b	1	141	2	244	199	175	135	135	MN (1)
unknown serotype	1	146	3	163	180	179	217	177	MD (1)
Hansen disease§	—	45	1	80	101	66	87	105	
Hantavirus pulmonary syndrome§	—	6	1	18	32	40	26	24	
Hemolytic uremic syndrome, postdiarrheal§	—	124	8	330	292	288	221	200	
Hepatitis C viral, acute	7	1,323	15	878	845	766	652	720	MI (1), MD (1), WV (1), KY (1), OK (1), WA (1), CA (1)
HIV infection, pediatric (age <13 years)§§	—	—	2	—	—	—	380	436	
Influenza-associated pediatric mortality§,¶¶	1	113	0	90	77	43	45	—	MS (1)
Listeriosis	6	434	21	759	808	884	896	753	NY (3), MI (1), OK (1), CA (1)
Measles***	—	55	1	140	43	55	66	37	
Meningococcal disease, invasive†††:									
A, C, Y, and W-135	2	183	4	330	325	318	297	—	MN (1), OK (1)
serogroup B	—	96	2	188	167	193	156	—	
other serogroup	—	18	0	38	35	32	27	—	
unknown serogroup	4	317	8	616	550	651	765	—	PA (1), MD (1), CA (2)
Mumps	7	252	13	454	800	6,584	314	258	NY (2), NYC (3), IL (1), NV (1)
Novel influenza A virus infections	—	§§§	0	2	4	N	N	N	
Plague	—	6	0	3	7	17	8	3	
Poliomyelitis, paralytic	—	—	—	—	—	—	1	—	
Polio virus infection, nonparalytic§	—	—	—	—	—	N	N	N	
Psittacosis§	—	7	0	8	12	21	16	12	
Q fever total§,¶¶¶:									
acute	1	55	3	124	171	169	136	70	
chronic	1	46	1	110	—	—	—	—	NE (1)
Rabies, human	—	9	—	14	—	—	—	—	
Rubella, human	—	1	0	2	1	3	2	7	
Rubella****	—	4	0	16	12	11	11	10	
Rubella, congenital syndrome	—	1	—	—	—	1	1	—	
SARS-CoV§,††††	—	—	—	—	—	—	—	—	
Smallpox§	—	—	—	—	—	—	—	—	
Streptococcal toxic-shock syndrome§	2	99	1	157	132	125	129	132	CT (2)
Syphilis, congenital (age <1 yr)	—	117	8	434	430	349	329	353	
Tetanus	—	6	1	19	28	41	27	34	
Toxic-shock syndrome (staphylococcal)§	2	54	2	71	92	101	90	95	NY (1), CA (1)
Trichinellosis	—	12	0	39	5	15	16	5	
Tularemia	1	42	4	123	137	95	154	134	AR (1)
Typhoid fever	3	228	11	449	434	353	324	322	CA (3)
Vancomycin-intermediate <i>Staphylococcus aureus</i> §	3	50	1	63	37	6	2	—	NY (1), FL (2)
Vancomycin-resistant <i>Staphylococcus aureus</i> §	—	—	—	—	2	1	3	1	
Vibriosis (noncholera <i>Vibrio</i> species infections)§	14	314	11	492	549	N	N	N	OH (1), MD (1), FL (5), AZ (3), WA (2), CA (2)
Yellow fever	—	—	—	—	—	—	—	—	

See Table I footnotes on next page.

TABLE I. (Continued) Provisional cases of infrequently reported notifiable diseases (<1,000 cases reported during the preceding year) — United States, week ending September 5, 2009 (35th week)*

—: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts.
 * Incidence data for reporting year 2008 and 2009 are provisional, whereas data for 2004, 2005, 2006, and 2007 are finalized.
 † Calculated by summing the incidence counts for the current week, the 2 weeks preceding the current week, and the 2 weeks following the current week, for a total of 5 preceding years. The total sum of incident cases is then divided by 25 weeks. Additional information is available at <http://www.cdc.gov/epo/dphsi/phs/files/5yearweeklyaverage.pdf>.
 § Not reportable in all states. Data from states where the condition is not reportable are excluded from this table, except starting in 2007 for the domestic arboviral diseases and influenza-associated pediatric mortality, and in 2003 for SARS-CoV. Reporting exceptions are available at <http://www.cdc.gov/epo/dphsi/phs/infdis.htm>.
 ¶ Includes both neuroinvasive and nonneuroinvasive. Updated weekly from reports to the Division of Vector-Borne Infectious Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases (ArboNET Surveillance). Data for West Nile virus are available in Table II.
 ** The names of the reporting categories changed in 2008 as a result of revisions to the case definitions. Cases reported prior to 2008 were reported in the categories: Ehrlichiosis, human monocytic (analogous to *E. chaffeensis*); Ehrlichiosis, human granulocytic (analogous to *Anaplasma phagocytophilum*), and Ehrlichiosis, unspecified, or other agent (which included cases unable to be clearly placed in other categories, as well as possible cases of *E. ewingii*).
 †† Data for *H. influenzae* (all ages, all serotypes) are available in Table II.
 §§ Updated monthly from reports to the Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention. Implementation of HIV reporting influences the number of cases reported. Updates of pediatric HIV data have been temporarily suspended until upgrading of the national HIV/AIDS surveillance data management system is completed. Data for HIV/AIDS, when available, are displayed in Table IV, which appears quarterly.
 ¶¶ Updated weekly from reports to the Influenza Division, National Center for Immunization and Respiratory Diseases. One hundred and twelve influenza-associated pediatric deaths occurring during the 2008–09 influenza season have been reported.
 *** No measles cases were reported for the current week.
 ††† Data for meningococcal disease (all serogroups) are available in Table II.
 §§§ CDC discontinued reporting of individual confirmed and probable cases of novel influenza A (H1N1) viruses infections on July 24, 2009. CDC will report the total number of novel influenza A (H1N1) hospitalizations and deaths weekly on the CDC H1N1 influenza website (<http://www.cdc.gov/h1n1flu>).
 ¶¶¶ In 2008, Q fever acute and chronic reporting categories were recognized as a result of revisions to the Q fever case definition. Prior to that time, case counts were not differentiated with respect to acute and chronic Q fever cases.
 **** No rubella cases were reported for the current week.
 †††† Updated weekly from reports to the Division of Viral and Rickettsial Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases.

FIGURE I. Selected notifiable disease reports, United States, comparison of provisional 4-week totals September 5, 2009, with historical data



* Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

Notifiable Disease Data Team and 122 Cities Mortality Data Team
 Patsy A. Hall
 Deborah A. Adams Rosaline Dhara
 Willie J. Anderson Michael S. Wodajo
 Jose Aponte Pearl C. Sharp
 Lenee Blanton

TABLE II. Provisional cases of selected notifiable diseases, United States, weeks ending September 5, 2009, and August 30, 2008 (35th week)*

Reporting area	Chlamydia†					Coccidioidomycosis					Cryptosporidiosis				
	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008
		Med	Max				Med	Max				Med	Max		
United States	11,558	22,433	25,700	753,367	791,568	186	156	472	7,398	4,349	108	124	401	4,084	4,809
New England	688	759	1,655	27,100	24,759	—	0	1	1	1	—	5	30	198	285
Connecticut	221	224	1,306	7,809	7,095	N	0	0	N	N	—	0	23	23	41
Maine§	34	48	75	1,634	1,689	N	0	0	N	N	—	0	5	22	30
Massachusetts	386	334	945	13,313	11,911	N	0	0	N	N	—	2	13	73	116
New Hampshire	1	39	63	1,162	1,378	—	0	1	1	1	—	1	4	39	47
Rhode Island§	—	63	244	2,392	1,898	—	0	0	—	—	—	0	3	4	7
Vermont§	46	21	53	790	788	N	0	0	N	N	—	1	7	37	44
Mid. Atlantic	2,613	2,913	6,734	103,410	98,562	—	0	0	—	—	15	13	28	484	481
New Jersey	293	415	838	13,952	15,045	N	0	0	N	N	—	0	3	8	29
New York (Upstate)	728	579	4,563	20,745	18,114	N	0	0	N	N	10	4	13	138	157
New York City	1,153	1,136	3,130	40,202	37,865	N	0	0	N	N	—	1	8	45	72
Pennsylvania	439	824	1,072	28,511	27,538	N	0	0	N	N	5	7	19	293	223
E.N. Central	1,593	3,494	4,382	113,700	129,441	—	0	4	22	34	11	28	120	908	1,246
Illinois	446	1,088	1,367	35,040	39,159	N	0	0	N	N	—	2	11	77	129
Indiana	365	418	713	15,219	14,459	N	0	0	N	N	—	3	17	128	125
Michigan	720	851	1,332	30,854	30,580	—	0	3	11	25	2	5	13	167	166
Ohio	62	798	1,300	21,173	30,826	—	0	2	11	9	9	9	59	278	333
Wisconsin	—	349	494	11,414	14,417	N	0	0	N	N	—	8	40	258	493
W.N. Central	418	1,320	1,658	43,513	44,712	—	0	1	7	1	8	17	47	638	646
Iowa	129	192	256	6,527	5,852	N	0	0	N	N	—	4	12	149	205
Kansas	1	154	549	5,289	6,159	N	0	0	N	N	—	1	5	50	55
Minnesota	—	258	342	7,983	9,665	—	0	0	—	—	—	4	33	182	132
Missouri	288	509	645	17,546	16,365	—	0	1	7	1	3	3	12	118	120
Nebraska§	—	101	219	3,423	3,565	N	0	0	N	N	5	2	5	64	76
North Dakota	—	25	60	772	1,202	N	0	0	N	N	—	0	10	7	2
South Dakota	—	57	81	1,973	1,904	N	0	0	N	N	—	2	6	68	56
S. Atlantic	1,831	4,136	5,453	131,289	161,296	—	0	1	5	3	24	21	49	680	579
Delaware	91	84	180	3,192	2,484	—	0	1	1	1	1	0	1	6	10
District of Columbia	126	127	226	4,586	4,661	—	0	0	—	—	—	0	2	2	9
Florida	580	1,413	1,597	49,125	47,905	N	0	0	N	N	17	8	35	247	254
Georgia	4	753	1,909	20,138	28,222	N	0	0	N	N	6	6	23	256	159
Maryland§	299	430	772	14,292	15,533	—	0	1	4	2	—	1	5	29	25
North Carolina	—	0	1,193	—	22,309	N	0	0	N	N	—	1	16	58	17
South Carolina§	—	550	1,424	16,142	17,186	N	0	0	N	N	—	1	6	32	34
Virginia§	683	616	926	21,371	20,826	N	0	0	N	N	—	1	5	40	53
West Virginia	48	68	101	2,443	2,170	N	0	0	N	N	—	0	2	10	18
E.S. Central	1,641	1,751	2,209	61,888	56,390	—	0	0	—	—	5	3	10	127	106
Alabama§	14	475	624	15,693	16,993	N	0	0	N	N	—	1	4	36	48
Kentucky	437	256	458	9,057	7,824	N	0	0	N	N	2	1	4	39	23
Mississippi	565	442	841	16,464	13,388	N	0	0	N	N	—	0	3	8	11
Tennessee§	625	575	809	20,674	18,185	N	0	0	N	N	3	1	5	44	24
W.S. Central	820	2,883	5,300	100,336	100,239	—	0	1	1	3	27	11	271	305	793
Arkansas§	239	273	418	9,723	9,653	N	0	0	N	N	2	1	10	30	45
Louisiana	290	420	1,134	14,457	14,640	—	0	1	1	3	2	1	6	29	40
Oklahoma	291	178	2,735	9,559	8,966	N	0	0	N	N	5	2	16	80	55
Texas§	—	1,967	2,520	66,597	66,980	N	0	0	N	N	18	7	258	166	653
Mountain	537	1,449	2,145	47,625	49,676	163	105	369	5,730	2,936	1	9	25	310	397
Arizona	67	467	735	15,695	16,502	163	104	365	5,662	2,858	—	1	4	25	59
Colorado	—	376	728	10,949	11,749	N	0	0	N	N	—	2	10	94	74
Idaho§	—	68	313	2,373	2,690	N	0	0	N	N	1	1	7	53	42
Montana§	—	54	88	1,903	2,059	N	0	0	N	N	—	0	4	27	38
Nevada§	276	167	455	6,780	6,551	—	1	3	40	43	—	0	4	14	11
New Mexico§	163	177	540	5,910	5,169	—	0	2	8	23	—	2	10	69	134
Utah	—	103	251	2,679	3,975	—	0	2	20	10	—	0	6	13	24
Wyoming§	31	34	97	1,336	981	—	0	1	—	2	—	0	2	15	15
Pacific	1,417	3,645	4,689	124,506	126,493	23	41	172	1,632	1,371	17	11	24	434	276
Alaska	—	96	199	3,071	3,153	N	0	0	N	N	—	0	1	4	3
California	1,279	2,803	3,599	96,945	98,369	23	41	172	1,632	1,371	9	6	20	256	161
Hawaii	—	119	247	3,942	3,895	N	0	0	N	N	—	0	1	1	1
Oregon§	115	201	631	6,308	6,671	N	0	0	N	N	2	3	8	121	50
Washington	23	414	571	14,240	14,405	N	0	0	N	N	6	1	6	52	61
American Samoa	—	0	0	—	73	N	0	0	N	N	N	0	0	N	N
C.N.M.I.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Guam	—	2	8	—	107	—	0	0	—	—	—	0	0	—	—
Puerto Rico	—	130	332	5,016	4,896	N	0	0	N	N	N	0	0	N	N
U.S. Virgin Islands	—	9	17	290	473	—	0	0	—	—	—	0	0	—	—

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2008 and 2009 are provisional. Data for HIV/AIDS, AIDS, and TB, when available, are displayed in Table IV, which appears quarterly.

† Chlamydia refers to genital infections caused by *Chlamydia trachomatis*.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending September 5, 2009, and August 30, 2008 (35th week)*

Reporting area	Giardiasis					Gonorrhea					Haemophilus influenzae, invasive All ages, all serotypes†				
	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008
		Med	Max				Med	Max				Med	Max		
United States	219	321	641	10,833	11,487	2,989	5,358	7,164	178,307	223,791	25	57	124	2,013	1,978
New England	5	27	55	828	1,036	88	95	301	3,246	3,464	2	3	16	133	114
Connecticut	—	5	14	162	224	43	46	275	1,486	1,607	2	0	12	42	27
Maine§	5	3	12	127	109	3	2	9	90	63	—	0	2	14	9
Massachusetts	—	10	27	318	436	42	39	112	1,345	1,469	—	2	5	64	55
New Hampshire	—	3	10	104	105	—	2	6	70	73	—	0	2	7	9
Rhode Island§	—	1	8	35	61	—	6	19	225	226	—	0	7	3	6
Vermont§	—	3	15	82	101	—	1	4	30	26	—	0	1	3	8
Mid. Atlantic	43	62	116	2,007	2,105	586	588	1,138	20,720	21,975	6	12	25	428	369
New Jersey	—	7	17	173	347	63	87	123	2,904	3,616	—	2	7	82	63
New York (Upstate)	27	24	81	815	696	207	106	664	3,854	4,058	2	3	20	103	107
New York City	—	16	30	494	558	204	210	577	7,451	6,926	—	2	11	82	63
Pennsylvania	16	16	46	525	504	112	190	267	6,511	7,375	4	4	10	161	136
E.N. Central	22	45	90	1,451	1,727	530	1,079	1,627	35,233	46,176	—	9	28	366	319
Illinois	—	9	25	270	476	175	335	494	10,750	13,633	—	3	9	102	100
Indiana	N	0	11	N	N	132	146	252	5,049	5,851	—	1	22	49	56
Michigan	1	12	22	390	372	195	284	493	9,957	11,253	—	0	3	16	17
Ohio	21	16	31	534	552	28	251	455	6,566	11,146	—	2	6	73	99
Wisconsin	—	8	19	257	327	—	91	140	2,911	4,293	—	2	20	126	47
W.N. Central	13	25	143	1,008	1,267	114	286	393	9,358	11,311	5	3	15	112	145
Iowa	—	6	18	199	203	17	34	53	1,114	1,019	—	0	0	—	2
Kansas	—	2	7	70	107	2	35	83	1,328	1,498	—	0	2	11	17
Minnesota	—	0	106	250	386	—	43	65	1,314	2,132	5	0	10	40	43
Missouri	12	7	25	316	334	95	129	183	4,429	5,409	—	1	4	37	53
Nebraska§	1	3	9	114	138	—	23	53	887	963	—	0	4	19	21
North Dakota	—	0	16	9	10	—	2	7	43	76	—	0	4	5	9
South Dakota	—	1	7	50	89	—	7	20	243	214	—	0	0	—	—
S. Atlantic	39	70	109	2,460	1,861	533	1,177	2,042	37,640	56,789	11	13	31	514	506
Delaware	—	0	3	18	28	28	16	37	631	762	—	0	1	3	6
District of Columbia	—	0	5	16	44	53	50	89	1,827	1,723	—	0	2	—	5
Florida	33	37	59	1,287	775	186	416	486	14,276	16,082	2	4	10	174	131
Georgia	—	13	67	650	464	3	251	876	6,840	10,464	3	3	9	113	104
Maryland§	2	5	9	163	180	88	121	212	3,823	4,128	3	1	6	64	74
North Carolina	N	0	0	N	N	—	0	470	—	9,812	—	1	17	57	54
South Carolina§	4	2	8	57	81	—	171	413	5,070	6,297	2	1	5	36	46
Virginia§	—	8	31	238	239	161	147	308	4,821	7,001	—	1	6	42	68
West Virginia	—	1	3	31	50	14	10	23	352	520	1	0	3	25	18
E.S. Central	3	8	20	232	308	456	514	714	17,795	20,485	—	3	9	118	103
Alabama§	1	3	12	109	178	6	143	216	4,432	6,705	—	0	4	26	16
Kentucky	N	0	0	N	N	108	84	135	2,631	3,098	—	0	5	18	6
Mississippi	N	0	0	N	N	179	142	252	5,138	4,879	—	0	1	4	11
Tennessee§	2	4	13	123	130	163	160	273	5,594	5,803	—	2	6	70	70
W.S. Central	13	9	22	298	267	290	865	1,383	29,227	34,539	—	2	22	79	90
Arkansas§	3	2	8	91	88	87	83	134	2,997	3,155	—	0	2	13	11
Louisiana	4	3	8	96	99	96	146	420	4,662	6,343	—	0	1	12	8
Oklahoma	6	4	18	111	80	107	70	613	3,328	3,299	—	1	20	53	64
Texas§	N	0	0	N	N	—	551	725	18,240	21,742	—	0	1	1	7
Mountain	19	26	62	883	998	119	176	313	5,652	7,756	1	5	11	171	222
Arizona	4	3	10	134	84	16	56	88	1,784	2,336	1	1	7	60	88
Colorado	—	9	27	311	340	—	57	152	1,616	2,302	—	1	6	53	42
Idaho§	4	3	10	112	125	—	2	13	67	121	—	0	1	4	12
Montana§	—	2	10	71	61	—	1	6	48	78	—	0	1	1	2
Nevada§	10	2	8	75	72	75	30	91	1,213	1,549	—	0	2	14	14
New Mexico§	1	1	7	58	76	27	24	52	746	939	—	0	3	16	32
Utah	—	4	15	91	210	—	5	15	126	347	—	1	2	20	29
Wyoming§	—	1	4	31	30	1	1	7	52	84	—	0	1	3	3
Pacific	62	50	130	1,666	1,918	273	550	765	19,436	21,296	—	2	8	92	110
Alaska	—	2	10	64	59	—	15	24	512	356	—	0	3	13	15
California	47	34	57	1,131	1,285	236	468	658	16,309	17,489	—	0	3	22	38
Hawaii	—	0	2	9	31	—	12	21	416	418	—	0	3	22	14
Oregon§	3	7	17	224	302	33	20	48	682	822	—	1	3	32	41
Washington	12	7	74	238	241	4	46	80	1,517	2,211	—	0	2	3	2
American Samoa	—	0	0	—	—	—	0	0	—	3	—	0	0	—	—
C.N.M.I.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Guam	—	0	0	—	—	—	1	15	—	45	—	0	0	—	—
Puerto Rico	1	2	15	63	132	—	4	24	165	194	—	0	1	2	1
U.S. Virgin Islands	—	0	0	—	—	—	2	7	80	91	N	0	0	N	N

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2008 and 2009 are provisional.

† Data for *H. influenzae* (age <5 yrs for serotype b, nonserotype b, and unknown serotype) are available in Table I.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending September 5, 2009, and August 30, 2008 (35th week)*

Reporting area	Hepatitis (viral, acute), by type†										Legionellosis				
	A				B										
	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008
	Med	Max				Med	Max				Med	Max			
United States	22	36	89	1,215	1,801	31	65	197	2,067	2,539	48	51	117	1,823	1,941
New England	2	2	8	54	90	—	1	4	24	56	3	2	18	85	136
Connecticut	2	0	4	16	18	—	0	3	10	22	3	1	5	40	26
Maine§	—	0	5	1	5	—	0	2	8	10	—	0	2	4	6
Massachusetts	—	1	3	29	46	—	0	2	3	14	—	1	5	25	58
New Hampshire	—	0	1	3	9	—	0	2	3	4	—	0	2	8	24
Rhode Island§	—	0	2	3	10	—	0	0	—	4	—	0	14	4	17
Vermont§	—	0	1	2	2	—	0	1	—	2	—	0	1	4	5
Mid. Atlantic	3	5	13	151	209	1	7	17	218	306	17	15	61	716	626
New Jersey	—	1	5	23	51	—	1	6	54	89	1	3	14	119	75
New York (Upstate)	3	1	4	36	41	—	1	11	38	42	11	5	29	234	188
New York City	—	2	6	50	72	—	1	4	41	69	—	2	20	123	85
Pennsylvania	—	1	4	42	45	1	3	8	85	106	5	6	25	240	278
E.N. Central	2	5	17	172	247	1	8	21	258	344	10	9	29	323	432
Illinois	—	1	12	77	91	—	1	7	35	131	—	1	13	26	55
Indiana	—	0	3	12	13	—	1	18	45	23	—	1	5	25	37
Michigan	—	1	5	46	89	—	3	8	89	95	3	2	10	77	123
Ohio	2	1	4	30	29	1	1	13	66	81	7	4	17	190	196
Wisconsin	—	0	3	7	25	—	0	4	23	14	—	0	6	5	21
W.N. Central	—	2	16	85	200	—	3	16	109	56	—	2	7	64	90
Iowa	—	1	3	24	95	—	0	3	20	14	—	0	2	15	12
Kansas	—	0	1	7	12	—	0	2	5	6	—	0	1	3	1
Minnesota	—	0	12	14	26	—	0	11	20	7	—	0	3	8	9
Missouri	—	0	3	19	25	—	1	5	52	23	—	1	5	28	50
Nebraska§	—	0	3	19	39	—	0	2	11	5	—	0	2	8	16
North Dakota	—	0	2	—	—	—	0	1	—	1	—	0	3	1	—
South Dakota	—	0	1	2	3	—	0	1	1	—	—	0	1	1	2
S. Atlantic	5	7	14	272	262	6	18	32	605	614	12	9	22	312	308
Delaware	—	0	1	3	6	U	0	1	U	U	—	0	5	11	9
District of Columbia	U	0	0	U	U	U	0	0	U	U	—	0	2	4	11
Florida	3	4	8	127	99	5	6	11	203	214	7	3	7	110	93
Georgia	—	1	4	43	35	1	3	9	97	118	—	1	5	33	25
Maryland§	—	0	4	27	33	—	1	5	48	55	5	2	10	74	90
North Carolina	—	0	4	25	47	—	2	19	130	51	—	0	7	39	14
South Carolina§	2	0	3	27	8	—	1	4	29	49	—	0	1	5	8
Virginia§	—	0	6	19	29	—	1	10	52	75	—	1	5	32	37
West Virginia	—	0	1	1	5	—	0	19	46	52	—	0	3	4	21
E.S. Central	3	1	5	30	57	5	7	11	206	261	3	2	11	81	87
Alabama§	—	0	2	7	8	2	2	7	62	73	—	0	1	7	13
Kentucky	1	0	2	7	21	1	2	7	53	63	1	1	3	36	41
Mississippi	1	0	1	8	4	—	1	2	18	32	—	0	1	2	1
Tennessee§	1	0	2	8	24	2	2	6	73	93	2	1	8	36	32
W.S. Central	—	3	43	103	172	4	10	99	325	505	—	1	21	44	57
Arkansas§	—	0	1	4	6	1	1	5	36	39	—	0	2	3	9
Louisiana	—	0	2	3	8	—	1	4	33	63	—	0	2	4	8
Oklahoma	—	0	6	3	7	3	2	17	69	74	—	0	6	3	3
Texas§	—	3	37	93	151	—	6	76	187	329	—	1	19	34	37
Mountain	—	3	7	109	162	1	3	7	88	139	1	2	8	73	56
Arizona	—	2	6	52	83	—	1	4	33	55	1	0	4	34	14
Colorado	—	1	5	34	28	—	0	2	15	24	—	0	2	7	6
Idaho§	—	0	1	3	16	—	0	2	6	6	—	0	1	1	3
Montana§	—	0	1	5	1	—	0	0	—	2	—	0	2	4	4
Nevada§	—	0	3	6	7	1	0	3	21	30	—	0	2	9	8
New Mexico§	—	0	1	5	15	—	0	2	5	8	—	0	2	2	5
Utah	—	0	2	4	9	—	0	3	5	9	—	0	4	15	16
Wyoming§	—	0	0	—	3	—	0	2	3	5	—	0	1	1	—
Pacific	7	7	17	239	402	13	6	36	234	258	2	4	12	125	149
Alaska	—	0	1	3	3	—	0	1	2	9	—	0	1	1	1
California	7	5	17	188	325	10	5	28	171	176	2	3	9	100	115
Hawaii	—	0	1	5	14	—	0	1	4	6	—	0	1	1	5
Oregon§	—	0	2	12	22	—	0	4	26	32	—	0	2	8	13
Washington	—	1	4	31	38	3	1	8	31	35	—	0	4	15	15
American Samoa	—	0	0	—	—	—	0	0	—	—	N	0	0	N	N
C.N.M.I.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Guam	—	0	0	—	—	—	0	0	—	—	—	0	0	—	—
Puerto Rico	—	0	2	17	19	—	0	3	12	40	—	0	0	—	—
U.S. Virgin Islands	—	0	0	—	—	—	0	0	—	—	—	0	0	—	—

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2008 and 2009 are provisional.

† Data for acute hepatitis C, viral are available in Table I.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending September 5, 2009, and August 30, 2008 (35th week)*

Reporting area	Lyme disease					Malaria					Meningococcal disease, invasive† All groups				
	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008
		Med	Max				Med	Max				Med	Max		
United States	410	486	1,637	17,950	23,066	12	23	46	740	778	6	17	48	614	863
New England	10	91	327	2,505	8,678	—	0	5	27	39	—	0	4	20	23
Connecticut	—	0	105	—	2,984	—	0	4	5	10	—	0	1	2	1
Maine§	10	8	73	467	312	—	0	1	1	1	—	0	1	3	4
Massachusetts	—	24	125	1,041	3,729	—	0	4	16	19	—	0	3	11	15
New Hampshire	—	13	73	750	1,255	—	0	1	2	3	—	0	1	1	2
Rhode Island§	—	0	78	54	119	—	0	1	1	2	—	0	1	2	1
Vermont§	—	5	35	193	279	—	0	1	2	4	—	0	1	1	—
Mid. Atlantic	346	246	1,401	11,230	9,198	1	5	17	167	209	1	2	5	69	96
New Jersey	—	35	263	2,621	2,767	—	0	3	—	52	—	0	2	8	13
New York (Upstate)	229	87	1,368	2,891	2,982	1	1	10	36	21	—	0	2	17	25
New York City	—	2	33	58	551	—	3	11	95	109	—	0	2	11	19
Pennsylvania	117	53	601	5,660	2,898	—	1	4	36	27	1	1	4	33	39
E.N. Central	7	19	174	1,459	1,828	1	3	8	101	111	—	3	8	99	149
Illinois	—	1	10	80	96	—	1	4	44	59	—	1	6	25	52
Indiana	—	1	4	33	30	—	0	1	7	5	—	0	3	24	22
Michigan	5	1	11	74	54	—	0	3	17	13	—	0	5	18	25
Ohio	2	1	5	32	28	1	1	6	29	22	—	0	3	26	32
Wisconsin	—	15	160	1,240	1,620	—	0	2	4	12	—	0	1	6	18
W.N. Central	1	5	336	163	438	2	1	7	37	44	1	1	9	49	76
Iowa	—	1	11	64	89	—	0	3	6	5	—	0	1	6	15
Kansas	—	0	4	15	6	—	0	2	3	4	—	0	2	7	4
Minnesota	—	1	326	67	329	—	0	7	13	19	1	0	4	10	21
Missouri	—	0	2	4	3	1	0	2	9	9	—	0	3	18	23
Nebraska§	1	0	3	12	8	1	0	1	5	7	—	0	1	5	10
North Dakota	—	0	10	—	—	—	0	0	—	—	—	0	3	1	1
South Dakota	—	0	1	1	3	—	0	1	1	—	—	0	1	2	2
S. Atlantic	36	63	205	2,366	2,701	2	6	17	233	195	1	3	9	113	119
Delaware	5	12	62	698	592	—	0	1	3	2	—	0	1	2	1
District of Columbia	—	0	5	18	53	—	0	2	5	2	—	0	0	—	—
Florida	4	1	10	48	40	2	2	7	69	34	—	1	4	42	41
Georgia	—	0	6	39	31	—	1	5	49	45	—	0	2	21	14
Maryland§	27	27	130	1,113	1,355	—	1	8	51	52	1	0	1	6	12
North Carolina	—	1	14	56	10	—	0	5	21	21	—	0	5	18	11
South Carolina§	—	0	3	18	18	—	0	1	2	7	—	0	1	10	19
Virginia§	—	12	61	312	498	—	1	4	31	30	—	0	2	9	16
West Virginia	—	0	17	64	104	—	0	1	2	2	—	0	2	5	5
E.S. Central	—	0	2	18	37	—	0	3	23	13	—	0	3	20	39
Alabama§	—	0	1	2	8	—	0	3	6	3	—	0	1	5	5
Kentucky	—	0	1	1	4	—	0	2	8	4	—	0	1	4	7
Mississippi	—	0	0	—	1	—	0	1	1	1	—	0	1	2	9
Tennessee§	—	0	2	15	24	—	0	3	8	5	—	0	1	9	18
W.S. Central	5	1	21	34	68	1	1	8	34	49	1	1	12	58	93
Arkansas§	—	0	0	—	—	—	0	1	3	—	—	0	2	5	13
Louisiana	—	0	1	—	2	1	0	1	3	2	—	0	3	11	19
Oklahoma	—	0	2	—	—	—	0	2	2	2	1	0	3	6	12
Texas§	5	1	21	34	66	—	1	7	26	45	—	1	9	36	49
Mountain	—	1	13	31	41	1	0	4	21	20	—	1	4	49	46
Arizona	—	0	2	3	7	1	0	2	5	9	—	0	2	13	6
Colorado	—	0	1	3	2	—	0	3	8	3	—	0	2	15	9
Idaho§	—	0	2	9	6	—	0	1	1	—	—	0	1	5	4
Montana§	—	0	13	2	4	—	0	3	4	—	—	0	2	4	4
Nevada§	—	0	2	12	10	—	0	1	—	4	—	0	2	4	7
New Mexico§	—	0	1	1	8	—	0	1	—	2	—	0	1	3	7
Utah	—	0	1	—	2	—	0	2	3	2	—	0	1	1	7
Wyoming§	—	0	1	1	2	—	0	0	—	—	—	0	2	4	2
Pacific	5	3	13	144	77	4	3	10	97	98	2	3	14	137	222
Alaska	—	0	1	2	5	—	0	1	2	4	—	0	2	3	6
California	5	3	12	125	39	3	2	8	72	70	2	2	8	92	163
Hawaii	N	0	0	N	N	—	0	1	1	2	—	0	1	3	4
Oregon§	—	0	3	12	26	—	0	2	9	4	—	0	6	26	26
Washington	—	0	12	5	7	1	0	3	13	18	—	0	6	13	23
American Samoa	N	0	0	N	N	—	0	0	—	—	—	0	0	—	—
C.N.M.I.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Guam	—	0	0	—	—	—	0	2	—	1	—	0	0	—	—
Puerto Rico	N	0	0	N	N	—	0	1	2	2	—	0	1	—	2
U.S. Virgin Islands	N	0	0	N	N	—	0	0	—	—	—	0	0	—	—

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2008 and 2009 are provisional.

† Data for meningococcal disease, invasive caused by serogroups A, C, Y, and W-135; serogroup B; other serogroup; and unknown serogroup are available in Table I.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending September 5, 2009, and August 30, 2008 (35th week)*

Reporting area	Pertussis					Rabies, animal					Rocky Mountain spotted fever				
	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008
		Med	Max				Med	Max				Med	Max		
United States	87	270	1,697	8,643	6,022	57	68	138	2,432	2,909	7	33	179	980	1,592
New England	—	15	29	386	683	10	8	14	224	272	—	0	2	8	4
Connecticut	—	1	4	26	39	8	3	10	101	130	—	0	0	—	—
Maine†	—	1	10	64	23	2	1	5	36	36	—	0	2	4	1
Massachusetts	—	8	21	224	537	—	0	0	—	—	—	0	1	3	1
New Hampshire	—	1	7	53	21	—	0	7	24	29	—	0	0	—	1
Rhode Island†	—	0	5	11	55	—	0	3	27	24	—	0	2	—	1
Vermont†	—	0	2	8	8	—	1	4	36	53	—	0	1	1	—
Mid. Atlantic	6	23	64	755	699	20	14	27	428	632	1	1	29	49	102
New Jersey	—	4	12	128	144	—	0	0	—	—	—	0	2	—	70
New York (Upstate)	2	5	41	136	265	20	8	20	310	343	1	0	29	10	12
New York City	1	0	21	53	49	—	0	2	—	12	—	0	4	21	9
Pennsylvania	3	13	33	438	241	—	5	17	118	277	—	0	2	18	11
E.N. Central	17	52	238	1,759	985	8	2	19	177	184	—	1	6	55	118
Illinois	—	12	45	268	190	3	1	9	71	77	—	1	6	33	86
Indiana	—	5	158	171	35	—	0	6	17	4	—	0	3	4	6
Michigan	4	11	26	452	157	—	1	5	49	63	—	0	2	5	3
Ohio	13	20	57	774	514	5	0	7	40	40	—	0	3	13	23
Wisconsin	—	3	11	94	89	N	0	0	N	N	—	0	0	—	—
W.N. Central	17	34	872	1,237	499	4	5	17	198	209	—	4	25	207	349
Iowa	—	6	21	122	76	—	0	5	24	16	—	0	2	4	7
Kansas	—	4	12	132	40	—	1	6	56	50	—	0	1	2	—
Minnesota	—	0	808	165	153	1	0	11	40	34	—	0	1	2	—
Missouri	14	20	51	678	159	3	1	5	47	47	—	4	24	189	324
Nebraska†	3	4	32	108	51	—	0	2	—	28	—	0	2	10	15
North Dakota	—	0	24	17	1	—	0	9	4	17	—	0	1	—	—
South Dakota	—	0	10	15	19	—	0	4	27	17	—	0	0	—	3
S. Atlantic	31	27	71	1,074	580	3	25	111	1,057	1,211	—	13	42	365	526
Delaware	—	0	2	10	11	—	0	0	—	—	—	0	3	14	26
District of Columbia	—	0	2	2	3	—	0	0	—	—	—	0	0	—	6
Florida	23	8	32	392	173	—	0	95	121	138	—	0	2	5	9
Georgia	—	3	11	106	61	—	0	71	262	272	—	1	6	34	64
Maryland†	1	3	10	76	74	—	6	14	226	312	—	1	3	27	66
North Carolina	—	0	65	204	79	N	0	4	N	N	—	8	36	225	222
South Carolina†	2	3	17	151	83	—	0	0	—	—	—	0	9	15	28
Virginia†	3	3	24	109	89	—	10	23	364	424	—	2	9	42	98
West Virginia	2	0	5	24	7	3	2	6	84	65	—	0	1	3	7
E.S. Central	2	14	33	548	216	1	2	7	70	131	3	4	19	170	236
Alabama†	—	4	19	202	28	—	0	0	—	—	—	1	6	38	61
Kentucky	2	6	15	178	58	1	1	4	36	32	—	0	1	1	1
Mississippi	—	1	4	40	76	—	0	2	—	2	—	0	1	7	9
Tennessee†	—	3	14	128	54	—	1	4	34	97	3	2	15	124	165
W.S. Central	—	55	389	1,748	960	—	0	13	45	74	3	1	161	105	224
Arkansas†	—	4	38	155	58	—	0	5	23	41	3	0	61	47	44
Louisiana	—	2	8	90	59	—	0	0	—	—	—	0	1	2	5
Oklahoma	—	0	45	36	30	—	0	13	21	31	—	0	98	44	142
Texas†	—	41	304	1,467	813	—	0	1	1	2	—	0	6	12	33
Mountain	1	17	31	571	593	—	1	9	57	61	—	1	3	19	30
Arizona	—	4	10	144	161	N	0	0	N	N	—	0	2	4	8
Colorado	—	5	12	193	109	—	0	0	—	—	—	0	0	—	1
Idaho†	1	1	5	54	22	—	0	2	—	7	—	0	1	1	1
Montana†	—	0	4	12	74	—	0	4	16	7	—	0	2	8	3
Nevada†	—	0	3	10	26	—	0	1	4	9	—	0	1	1	2
New Mexico†	—	1	10	37	32	—	0	2	16	23	—	0	1	1	3
Utah	—	3	19	113	158	—	0	6	4	4	—	0	1	1	5
Wyoming†	—	0	5	8	11	—	0	4	17	11	—	0	2	3	7
Pacific	13	19	98	565	807	11	5	12	176	135	—	0	1	2	3
Alaska	—	1	21	31	108	—	0	2	10	12	N	0	0	N	N
California	—	5	19	143	366	10	4	12	152	116	—	0	1	2	—
Hawaii	2	0	3	22	8	—	0	0	—	—	N	0	0	N	N
Oregon†	4	3	16	175	122	1	0	3	14	7	—	0	0	—	3
Washington	7	6	76	194	203	—	0	0	—	—	—	0	0	—	—
American Samoa	—	0	0	—	—	N	0	0	N	N	N	0	0	N	N
C.N.M.I.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Guam	—	0	0	—	—	—	0	0	—	—	N	0	0	N	N
Puerto Rico	—	0	1	1	—	—	1	3	27	45	N	0	0	N	N
U.S. Virgin Islands	—	0	0	—	—	N	0	0	N	N	N	0	0	N	N

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2008 and 2009 are provisional.

† Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending September 5, 2009, and August 30, 2008 (35th week)*

Reporting area	Salmonellosis					Shiga toxin-producing <i>E. coli</i> (STEC) [†]					Shigellosis				
	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008
		Med	Max				Med	Max				Med	Max		
United States	788	900	2,323	27,342	30,111	101	86	255	2,573	3,214	164	317	1,268	9,929	12,954
New England	—	30	303	1,316	1,642	—	3	51	137	176	—	3	31	150	167
Connecticut	—	0	277	277	491	—	0	51	51	47	—	0	26	26	40
Maine [§]	—	2	7	83	103	—	0	3	14	11	—	0	1	2	18
Massachusetts	—	20	39	631	813	—	1	6	41	85	—	2	15	101	92
New Hampshire	—	3	42	198	104	—	1	3	23	14	—	0	3	8	4
Rhode Island [§]	—	2	11	87	67	—	0	1	—	7	—	0	1	8	10
Vermont [§]	—	1	5	40	64	—	0	6	8	12	—	0	2	5	3
Mid. Atlantic	77	91	182	2,952	3,812	51	6	16	268	338	21	56	79	1,878	1,647
New Jersey	—	11	41	226	912	—	1	5	24	103	—	13	35	391	555
New York (Upstate)	54	24	66	869	866	6	3	9	90	109	11	5	23	156	432
New York City	—	18	49	712	844	—	1	5	39	35	—	9	23	267	529
Pennsylvania	23	29	66	1,145	1,190	45	1	5	115	91	10	24	61	1,064	131
E.N. Central	35	90	141	3,144	3,479	7	12	74	420	509	7	65	132	1,841	2,483
Illinois	—	25	50	810	1,020	—	1	10	65	84	—	13	25	362	710
Indiana	—	8	50	243	404	—	1	7	39	61	—	1	21	38	490
Michigan	8	18	29	647	656	2	3	43	98	92	3	5	24	158	88
Ohio	27	28	52	1,027	878	5	3	15	100	120	4	36	80	931	937
Wisconsin	—	12	30	417	521	—	3	10	118	152	—	11	42	352	258
W.N. Central	39	51	109	1,825	1,936	11	12	38	494	567	14	15	49	601	636
Iowa	—	7	16	284	306	—	2	14	116	149	—	2	12	47	114
Kansas	—	6	19	225	311	—	1	7	30	31	—	3	11	147	25
Minnesota	15	12	51	439	504	8	2	19	148	111	4	2	14	58	214
Missouri	18	12	31	416	516	3	2	10	81	122	8	3	40	324	171
Nebraska [§]	6	5	41	267	163	—	2	6	64	118	2	0	3	19	5
North Dakota	—	0	30	40	31	—	0	28	3	1	—	0	9	3	32
South Dakota	—	3	22	154	105	—	0	12	52	35	—	0	1	3	75
S. Atlantic	275	262	440	7,444	7,284	7	13	30	415	569	29	47	85	1,558	2,179
Delaware	1	2	8	70	104	—	0	2	10	9	—	1	8	72	7
District of Columbia	—	0	5	20	47	—	0	1	1	5	—	0	2	6	15
Florida	179	110	197	3,513	2,993	3	3	7	112	99	9	8	24	304	616
Georgia	51	39	96	1,408	1,423	1	1	4	49	65	12	13	30	451	806
Maryland [§]	15	16	26	483	567	3	2	8	58	93	6	6	14	250	66
North Carolina	—	25	104	778	697	—	2	21	74	59	—	6	27	251	99
South Carolina [§]	26	14	54	439	655	—	0	3	20	31	2	4	14	81	427
Virginia [§]	—	20	88	577	654	—	2	16	73	178	—	5	59	137	116
West Virginia	3	4	23	156	144	—	0	3	18	30	—	0	3	6	27
E.S. Central	26	56	140	1,758	2,147	3	4	12	144	187	8	19	58	571	1,352
Alabama [§]	5	15	40	426	610	—	1	4	33	48	—	4	11	97	321
Kentucky	8	10	18	335	298	1	2	7	52	59	3	2	25	143	207
Mississippi	—	14	57	509	713	—	0	1	6	4	—	1	4	28	273
Tennessee [§]	13	14	62	488	526	2	2	5	53	76	5	11	48	303	551
W.S. Central	164	110	1,333	2,894	4,129	4	4	139	96	233	39	60	967	1,758	2,853
Arkansas [§]	24	12	38	400	470	—	1	5	25	34	1	8	20	234	368
Louisiana	12	18	43	599	729	—	0	1	—	7	1	4	17	108	487
Oklahoma	29	14	102	422	477	2	0	82	19	22	6	5	61	186	83
Texas [§]	99	55	1,204	1,473	2,453	2	2	55	52	170	31	42	889	1,230	1,915
Mountain	19	57	107	1,903	2,250	3	10	40	311	372	10	24	54	788	626
Arizona	13	20	45	677	700	—	1	4	50	47	9	16	41	591	307
Colorado	—	13	34	440	501	—	3	18	103	101	—	2	11	62	70
Idaho [§]	1	4	10	128	117	3	2	15	58	77	—	0	2	7	8
Montana [§]	—	2	7	73	80	—	0	3	15	27	—	0	5	13	4
Nevada [§]	4	4	13	179	158	—	0	4	20	13	1	1	11	47	150
New Mexico [§]	1	5	25	200	405	—	0	2	19	41	—	2	12	57	60
Utah	—	6	15	163	237	—	1	5	41	56	—	0	3	11	24
Wyoming [§]	—	1	6	43	52	—	0	2	5	10	—	0	1	—	3
Pacific	153	126	537	4,106	3,432	15	9	31	288	263	36	27	82	784	1,011
Alaska	—	1	6	52	41	—	0	1	—	5	—	0	1	1	—
California	117	94	516	3,122	2,482	8	5	15	156	120	35	20	75	633	876
Hawaii	2	5	13	163	181	—	0	1	3	11	—	0	4	24	31
Oregon [§]	2	8	15	274	308	1	1	6	41	44	—	1	10	27	50
Washington	32	12	85	495	420	6	3	16	88	83	1	3	11	99	54
American Samoa	—	0	1	—	2	—	0	0	—	—	—	0	2	3	1
C.N.M.I.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Guam	—	0	2	—	10	—	0	0	—	—	—	0	1	—	14
Puerto Rico	2	9	40	244	454	1	0	0	1	—	—	0	2	7	21
U.S. Virgin Islands	—	0	0	—	—	—	0	0	—	—	—	0	0	—	—

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2008 and 2009 are provisional.

† Includes *E. coli* O157:H7; Shiga toxin-positive, serogroup non-O157; and Shiga toxin-positive, not serogrouped.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending September 5, 2009, and August 30, 2008 (35th week)*

Reporting area	Streptococcal diseases, invasive, group A				<i>Streptococcus pneumoniae</i> , invasive disease, nondrug resistant†					
	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008
		Med	Max				Med	Max		
United States	27	101	239	3,822	4,087	18	36	122	1,172	1,221
New England	—	5	28	223	296	—	1	12	41	60
Connecticut	—	0	21	63	83	—	0	11	—	—
Maine§	—	0	2	13	20	—	0	1	3	1
Massachusetts	—	2	10	91	140	—	1	4	28	44
New Hampshire	—	1	4	34	20	—	0	2	8	8
Rhode Island§	—	0	2	9	21	—	0	2	—	7
Vermont§	—	0	3	13	12	—	0	1	2	—
Mid. Atlantic	6	19	43	776	839	3	5	33	180	153
New Jersey	—	3	6	103	153	—	1	4	31	46
New York (Upstate)	4	7	25	258	261	2	2	17	85	68
New York City	—	4	12	145	151	1	0	31	64	39
Pennsylvania	2	6	18	270	274	N	0	2	N	N
E.N. Central	3	17	42	731	779	3	6	18	177	222
Illinois	—	5	12	204	210	—	1	5	23	63
Indiana	—	3	23	116	102	—	0	13	25	24
Michigan	—	3	11	119	133	—	1	5	47	57
Ohio	3	4	13	185	214	3	1	6	52	41
Wisconsin	—	2	10	107	120	—	1	4	30	37
W.N. Central	1	6	37	318	311	3	2	11	107	66
Iowa	—	0	0	—	—	—	0	0	—	—
Kansas	—	1	5	37	32	N	0	1	N	N
Minnesota	1	0	34	146	150	2	0	10	60	18
Missouri	—	1	8	69	72	—	0	4	29	29
Nebraska§	—	1	3	35	31	1	0	1	8	7
North Dakota	—	0	4	11	8	—	0	3	4	6
South Dakota	—	0	3	20	18	—	0	2	6	6
S. Atlantic	12	22	48	869	832	2	6	16	216	239
Delaware	—	0	1	9	6	—	0	0	—	—
District of Columbia	—	0	3	11	12	N	0	0	N	N
Florida	2	6	12	212	189	—	1	6	49	46
Georgia	5	5	13	206	185	1	2	6	54	63
Maryland§	4	3	12	138	145	1	1	4	51	46
North Carolina	—	2	12	81	104	N	0	0	N	N
South Carolina§	1	1	5	53	51	—	1	6	32	42
Virginia§	—	3	9	125	108	—	0	4	18	37
West Virginia	—	1	4	34	32	—	0	3	12	5
E.S. Central	1	4	10	147	144	2	2	7	61	63
Alabama§	N	0	0	N	N	N	0	0	N	N
Kentucky	—	1	5	28	31	N	0	0	N	N
Mississippi	N	0	0	N	N	—	0	2	14	8
Tennessee§	1	3	9	119	113	2	1	6	47	55
W.S. Central	3	9	79	318	359	4	6	46	198	189
Arkansas§	—	0	2	14	8	—	0	4	20	11
Louisiana	—	0	3	11	14	—	0	3	13	11
Oklahoma	—	3	20	108	81	1	1	7	40	49
Texas§	3	5	59	185	256	3	4	34	125	118
Mountain	1	10	22	329	422	1	4	16	169	193
Arizona	1	3	7	110	146	1	2	10	88	89
Colorado	—	3	9	107	106	—	1	4	32	42
Idaho§	—	0	2	7	12	—	0	2	7	3
Montana§	N	0	0	N	N	N	0	0	N	N
Nevada§	—	0	1	5	8	—	0	1	—	3
New Mexico§	—	2	7	59	103	—	0	4	15	27
Utah	—	1	6	40	41	—	0	5	27	28
Wyoming§	—	0	1	1	6	—	0	1	—	1
Pacific	—	3	9	111	105	—	0	4	23	36
Alaska	—	1	3	21	26	—	0	3	17	23
California	N	0	0	N	N	N	0	0	N	N
Hawaii	—	3	8	90	79	—	0	2	6	13
Oregon§	N	0	0	N	N	N	0	0	N	N
Washington	N	0	0	N	N	N	0	0	N	N
American Samoa	—	0	0	—	30	N	0	0	N	N
C.N.M.I.	—	—	—	—	—	—	—	—	—	—
Guam	—	0	0	—	—	—	0	0	—	—
Puerto Rico	N	0	0	N	N	N	0	0	N	N
U.S. Virgin Islands	—	0	0	—	—	N	0	0	N	N

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2008 and 2009 are provisional.

† Includes cases of invasive pneumococcal disease, in children aged <5 years, caused by *S. pneumoniae*, which is susceptible or for which susceptibility testing is not available (NNDS event code 11717).

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending September 5, 2009, and August 30, 2008 (35th week)*

Reporting area	<i>Streptococcus pneumoniae</i> , invasive disease, drug resistant†										Syphilis, primary and secondary				
	All ages				Aged <5 years										
	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008
	Med	Max				Med	Max				Med	Max			
United States	23	60	276	1,984	2,203	2	9	21	307	336	102	262	452	8,746	8,489
New England	—	1	48	34	46	—	0	5	3	7	2	5	15	225	216
Connecticut	—	0	48	—	—	—	0	5	—	—	—	1	5	41	21
Maine§	—	0	2	9	15	—	0	1	1	1	—	0	1	1	9
Massachusetts	—	0	1	2	—	—	0	1	2	—	2	4	11	159	153
New Hampshire	—	0	3	5	—	—	0	0	—	—	—	0	2	12	13
Rhode Island§	—	0	6	7	18	—	0	1	—	4	—	0	5	12	14
Vermont§	—	0	2	11	13	—	0	0	—	2	—	0	2	—	6
Mid. Atlantic	1	3	14	117	228	—	0	3	20	20	39	35	51	1,255	1,122
New Jersey	—	0	0	—	—	—	0	0	—	—	2	4	13	153	151
New York (Upstate)	1	1	10	52	48	—	0	2	10	6	3	2	8	85	93
New York City	—	0	4	3	92	—	0	2	—	1	29	23	40	786	693
Pennsylvania	—	1	8	62	88	—	0	2	10	13	5	6	12	231	185
E.N. Central	2	11	41	439	471	—	1	7	63	64	20	23	44	703	783
Illinois	N	0	0	N	N	N	0	0	N	N	9	7	19	203	311
Indiana	—	3	32	151	161	—	0	6	21	20	3	2	10	108	96
Michigan	—	0	2	19	17	—	0	1	2	2	8	3	18	165	129
Ohio	2	7	18	269	293	—	1	4	40	42	—	6	17	197	208
Wisconsin	—	0	0	—	—	—	0	0	—	—	—	1	4	30	39
W.N. Central	1	2	161	94	155	—	0	3	20	31	1	6	11	208	284
Iowa	—	0	0	—	—	—	0	0	—	—	—	0	2	16	13
Kansas	—	1	5	39	58	—	0	2	13	3	1	0	3	19	23
Minnesota	—	0	156	—	23	—	0	3	—	23	—	2	6	40	71
Missouri	1	1	5	43	67	—	0	1	5	2	—	3	6	115	167
Nebraska§	—	0	0	—	—	—	0	0	—	—	—	0	3	14	10
North Dakota	—	0	3	10	2	—	0	0	—	—	—	0	1	3	—
South Dakota	—	0	2	2	5	—	0	2	2	3	—	0	1	1	—
S. Atlantic	17	26	53	946	902	2	4	14	140	147	14	64	262	2,154	1,828
Delaware	—	0	2	15	3	—	0	0	—	—	—	0	3	22	10
District of Columbia	N	0	0	N	N	N	0	0	N	N	2	3	9	116	94
Florida	10	15	36	550	513	—	2	13	86	96	3	20	31	658	692
Georgia	5	8	25	290	300	2	1	5	47	43	—	14	227	488	402
Maryland§	—	0	1	4	4	—	0	0	—	1	9	6	16	211	227
North Carolina	N	0	0	N	N	N	0	0	N	N	—	9	21	361	171
South Carolina§	—	0	0	—	—	—	0	0	—	—	—	2	6	78	57
Virginia§	N	0	0	N	N	N	0	0	N	N	—	7	16	216	167
West Virginia	2	2	13	87	82	—	0	3	7	7	—	0	2	4	8
E.S. Central	1	5	25	196	233	—	1	3	29	42	12	22	36	774	723
Alabama§	N	0	0	N	N	N	0	0	N	N	—	8	17	288	299
Kentucky	—	1	5	55	56	—	0	2	7	9	3	1	10	46	58
Mississippi	—	0	3	3	28	—	0	1	2	8	3	4	18	152	103
Tennessee§	1	3	23	138	149	—	0	3	20	25	6	8	19	288	263
W.S. Central	1	2	6	72	74	—	0	3	14	12	7	49	80	1,640	1,452
Arkansas§	1	1	5	40	13	—	0	3	9	3	6	4	35	151	110
Louisiana	—	1	5	32	61	—	0	1	5	9	—	11	40	303	403
Oklahoma	N	0	0	N	N	N	0	0	N	N	1	1	7	42	51
Texas§	—	0	0	—	—	—	0	0	—	—	—	32	48	1,144	888
Mountain	—	2	7	84	92	—	0	3	17	11	2	9	18	293	434
Arizona	—	0	0	—	—	—	0	0	—	—	1	4	9	132	223
Colorado	—	0	0	—	—	—	0	0	—	—	—	1	4	58	104
Idaho§	N	0	1	N	N	N	0	1	N	N	—	0	2	3	2
Montana§	—	0	1	—	—	—	0	0	—	—	—	0	7	—	—
Nevada§	—	1	4	33	44	—	0	2	7	5	1	1	7	65	56
New Mexico§	—	0	0	—	—	—	0	0	—	—	—	1	5	33	30
Utah	—	1	6	42	47	—	0	3	9	6	—	0	2	—	16
Wyoming§	—	0	2	9	1	—	0	1	1	—	—	0	1	2	3
Pacific	—	0	1	2	2	—	0	1	1	2	5	44	66	1,494	1,647
Alaska	—	0	0	—	—	—	0	0	—	—	—	0	0	—	1
California	N	0	0	N	N	N	0	0	N	N	3	40	59	1,360	1,488
Hawaii	—	0	1	2	2	—	0	1	1	2	—	0	3	19	16
Oregon§	N	0	0	N	N	N	0	0	N	N	1	1	4	32	12
Washington	N	0	0	N	N	N	0	0	N	N	1	2	7	83	130
American Samoa	N	0	0	N	N	N	0	0	N	N	—	0	0	—	—
C.N.M.I.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Guam	—	0	0	—	—	—	0	0	—	—	—	0	0	—	—
Puerto Rico	—	0	0	—	—	—	0	0	—	—	—	3	16	142	101
U.S. Virgin Islands	—	0	0	—	—	—	0	0	—	—	—	0	0	—	—

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2008 and 2009 are provisional.

† Includes cases of invasive pneumococcal disease caused by drug-resistant *S. pneumoniae* (DRSP) (NNDSS event code 11720).

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending September 5, 2009, and August 30, 2008 (35th week)*

Reporting area	West Nile virus disease†														
	Varicella (chickenpox)					Neuroinvasive					Nonneuroinvasive§				
	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008	Current week	Previous 52 weeks		Cum 2009	Cum 2008
	Med	Max				Med	Max				Med	Max			
United States	60	451	1,035	12,161	20,567	—	1	70	98	417	—	0	55	97	483
New England	—	8	46	194	1,139	—	0	2	—	3	—	0	0	—	3
Connecticut	—	0	21	—	587	—	0	2	—	3	—	0	0	—	3
Maine¶	—	0	11	5	175	—	0	0	—	—	—	0	0	—	—
Massachusetts	—	0	1	1	—	—	0	1	—	—	—	0	0	—	—
New Hampshire	—	4	11	141	181	—	0	0	—	—	—	0	0	—	—
Rhode Island¶	—	0	1	4	—	—	0	1	—	—	—	0	0	—	—
Vermont¶	—	2	17	43	196	—	0	0	—	—	—	0	0	—	—
Mid. Atlantic	15	38	58	1,049	1,637	—	0	6	2	28	—	0	4	—	11
New Jersey	N	0	0	N	N	—	0	2	—	2	—	0	1	—	3
New York (Upstate)	N	0	0	N	N	—	0	3	1	13	—	0	2	—	3
New York City	—	0	0	—	—	—	0	2	—	5	—	0	1	—	4
Pennsylvania	15	38	58	1,049	1,637	—	0	1	1	8	—	0	1	—	1
E.N. Central	25	154	254	4,184	4,977	—	0	8	2	15	—	0	3	1	12
Illinois	—	33	73	835	720	—	0	4	1	2	—	0	1	—	6
Indiana	—	2	19	207	—	—	0	1	1	2	—	0	1	—	—
Michigan	1	48	90	1,314	2,094	—	0	4	—	3	—	0	2	—	2
Ohio	24	42	91	1,444	1,598	—	0	3	—	6	—	0	1	1	—
Wisconsin	—	14	55	384	565	—	0	2	—	2	—	0	0	—	4
W.N. Central	4	22	114	669	811	—	0	6	5	33	—	0	6	20	108
Iowa	N	0	0	N	N	—	0	1	—	2	—	0	1	1	2
Kansas	—	5	22	176	318	—	0	2	—	8	—	0	2	4	12
Minnesota	—	0	0	—	—	—	0	0	—	2	—	0	2	1	6
Missouri	4	10	51	436	459	—	0	3	1	5	—	0	1	—	2
Nebraska¶	N	0	0	N	N	—	0	1	1	3	—	0	3	6	26
North Dakota	—	0	108	57	—	—	0	0	—	2	—	0	0	—	35
South Dakota	—	0	4	—	34	—	0	2	3	11	—	0	2	8	25
S. Atlantic	15	57	146	1,433	3,353	—	0	2	3	13	—	0	3	—	13
Delaware	—	0	4	8	30	—	0	0	—	—	—	0	0	—	1
District of Columbia	—	0	3	8	18	—	0	2	—	2	—	0	1	—	1
Florida	9	28	67	927	1,181	—	0	0	—	3	—	0	0	—	—
Georgia	N	0	0	N	N	—	0	1	2	2	—	0	1	—	3
Maryland¶	N	0	0	N	N	—	0	2	—	3	—	0	2	—	5
North Carolina	N	0	0	N	N	—	0	0	—	2	—	0	0	—	1
South Carolina¶	—	4	54	154	590	—	0	1	1	—	—	0	0	—	1
Virginia¶	—	0	119	28	1,027	—	0	0	—	—	—	0	0	—	1
West Virginia	6	9	32	308	507	—	0	0	—	1	—	0	0	—	—
E.S. Central	—	12	28	358	861	—	0	5	17	33	—	0	5	10	45
Alabama¶	—	12	28	356	851	—	0	1	—	9	—	0	2	—	5
Kentucky	N	0	0	N	N	—	0	1	—	—	—	0	0	—	—
Mississippi	—	0	1	2	10	—	0	5	16	16	—	0	5	9	33
Tennessee¶	N	0	0	N	N	—	0	2	1	8	—	0	1	1	7
W.S. Central	1	94	747	3,259	6,200	—	0	7	25	48	—	0	5	9	42
Arkansas¶	—	3	47	96	491	—	0	1	1	6	—	0	0	—	2
Louisiana	1	1	7	76	56	—	0	3	5	12	—	0	5	5	16
Oklahoma	N	0	0	N	N	—	0	1	2	2	—	0	0	—	5
Texas¶	—	86	721	3,087	5,653	—	0	6	17	28	—	0	2	4	19
Mountain	—	33	83	936	1,499	—	0	12	32	56	—	0	17	42	131
Arizona	—	0	0	—	—	—	0	7	9	30	—	0	8	3	21
Colorado	—	13	44	367	608	—	0	3	9	12	—	0	9	24	42
Idaho¶	N	0	0	N	N	—	0	1	2	3	—	0	2	6	32
Montana¶	—	2	20	105	229	—	0	1	2	—	—	0	1	1	5
Nevada¶	N	0	0	N	N	—	0	2	7	5	—	0	1	5	6
New Mexico¶	—	2	20	134	160	—	0	1	2	3	—	0	1	1	1
Utah	—	13	31	330	492	—	0	1	—	3	—	0	1	—	18
Wyoming¶	—	0	1	—	10	—	0	1	1	—	—	0	2	2	6
Pacific	—	2	7	79	90	—	0	34	12	188	—	0	15	15	118
Alaska	—	1	6	49	44	—	0	0	—	—	—	0	0	—	—
California	—	0	0	—	—	—	0	33	12	184	—	0	15	14	104
Hawaii	—	1	4	30	46	—	0	0	—	—	—	0	0	—	—
Oregon¶	N	0	0	N	N	—	0	1	—	2	—	0	0	—	13
Washington	N	0	0	N	N	—	0	0	—	2	—	0	1	1	1
American Samoa	N	0	0	N	N	—	0	0	—	—	—	0	0	—	—
C.N.M.I.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Guam	—	2	3	—	55	—	0	0	—	—	—	0	0	—	—
Puerto Rico	1	8	23	317	433	—	0	0	—	—	—	0	0	—	—
U.S. Virgin Islands	—	0	0	—	—	—	0	0	—	—	—	0	0	—	—

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not reportable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2008 and 2009 are provisional. Data for HIV/AIDS, AIDS, and TB, when available, are displayed in Table IV, which appears quarterly.

† Updated weekly from reports to the Division of Vector-Borne Infectious Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases (ArboNET Surveillance). Data for California serogroup, eastern equine, Powassan, St. Louis, and western equine diseases are available in Table I.

§ Not reportable in all states. Data from states where the condition is not reportable are excluded from this table, except starting in 2007 for the domestic arboviral diseases and influenza-associated pediatric mortality, and in 2003 for SARS-CoV. Reporting exceptions are available at <http://www.cdc.gov/epo/dphsi/phs/infdis.htm>.

¶ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE III. Deaths in 122 U.S. cities,* week ending September 5, 2009 (35th week)

Reporting area	All causes, by age (years)							Reporting area	All causes, by age (years)						
	All Ages	≥65	45-64	25-44	1-24	<1	P&† Total		All Ages	≥65	45-64	25-44	1-24	<1	P&† Total
New England	444	291	100	34	4	15	30	S. Atlantic	1,181	711	300	100	35	35	55
Boston, MA	106	60	29	7	2	8	4	Atlanta, GA	107	63	23	11	4	6	4
Bridgeport, CT	23	17	2	4	—	—	4	Baltimore, MD	132	72	32	20	3	5	5
Cambridge, MA	15	11	4	—	—	—	1	Charlotte, NC	107	64	29	9	2	3	7
Fall River, MA	21	17	3	1	—	—	1	Jacksonville, FL	120	71	35	9	3	2	2
Hartford, CT	49	31	12	5	—	1	2	Miami, FL	73	51	12	7	2	1	8
Lowell, MA	22	18	2	2	—	—	1	Norfolk, VA	51	31	15	—	2	3	4
Lynn, MA	4	2	2	—	—	—	—	Richmond, VA	53	26	19	7	1	—	2
New Bedford, MA	20	16	1	3	—	—	1	Savannah, GA	67	41	19	4	1	2	4
New Haven, CT	20	13	5	2	—	—	5	St. Petersburg, FL	54	31	13	4	4	2	5
Providence, RI	63	41	15	5	—	2	3	Tampa, FL	209	127	51	17	8	6	11
Somerville, MA	4	1	3	—	—	—	—	Washington, D.C.	201	129	51	11	5	5	1
Springfield, MA	29	14	9	3	1	2	1	Wilmington, DE	7	5	1	1	—	—	2
Waterbury, CT	26	19	5	2	—	—	1	E.S. Central	771	487	199	50	17	18	52
Worcester, MA	42	31	8	—	1	2	6	Birmingham, AL	122	74	37	8	1	2	9
Mid. Atlantic	1,747	1,193	380	113	31	30	59	Chattanooga, TN	75	53	17	3	2	—	1
Albany, NY	46	29	12	3	1	1	1	Knoxville, TN	109	76	24	6	1	2	6
Allentown, PA	36	25	8	2	—	1	2	Lexington, KY	50	26	16	5	1	2	2
Buffalo, NY	64	40	17	4	1	2	3	Memphis, TN	156	94	37	14	4	7	15
Camden, NJ	33	17	11	2	1	2	—	Mobile, AL	94	56	29	5	1	3	5
Elizabeth, NJ	18	11	4	3	—	—	1	Montgomery, AL	19	15	3	—	1	—	2
Erie, PA	37	30	5	2	—	—	3	Nashville, TN	146	93	36	9	6	2	12
Jersey City, NJ	20	10	7	2	1	—	3	W.S. Central	1,192	757	301	70	37	26	67
New York City, NY	985	688	210	56	14	17	22	Austin, TX	105	66	26	8	2	3	6
Newark, NJ	U	U	U	U	U	U	U	Baton Rouge, LA	U	U	U	U	U	U	U
Paterson, NJ	8	5	2	1	—	—	—	Corpus Christi, TX	U	U	U	U	U	U	U
Philadelphia, PA	247	144	69	23	8	3	13	Dallas, TX	156	91	47	9	3	6	13
Pittsburgh, PA§	U	U	U	U	U	U	U	El Paso, TX	94	67	21	2	2	2	3
Reading, PA	31	23	5	2	1	—	1	Fort Worth, TX	U	U	U	U	U	U	U
Rochester, NY	123	93	18	7	2	3	6	Houston, TX	357	213	97	23	18	6	11
Schenectady, NY	9	7	2	—	—	—	—	Little Rock, AR	72	45	21	3	2	1	1
Scranton, PA	29	23	3	2	1	—	1	New Orleans, LA	U	U	U	U	U	U	U
Syracuse, NY	13	9	2	1	—	1	3	San Antonio, TX	247	162	53	15	8	8	22
Trenton, NJ	22	17	3	1	1	—	—	Shreveport, LA	40	24	10	5	1	—	4
Utica, NY	9	7	1	1	—	—	—	Tulsa, OK	121	89	26	5	1	—	7
Yonkers, NY	17	15	1	1	—	—	—	Mountain	1,171	737	278	95	33	28	54
E.N. Central	1,343	927	302	63	21	30	68	Albuquerque, NM	107	66	27	10	3	1	3
Akron, OH	51	33	15	1	—	2	1	Boise, ID	48	37	7	3	—	1	9
Canton, OH	31	23	6	—	2	—	1	Colorado Springs, CO	141	95	29	10	3	4	3
Chicago, IL	U	U	U	U	U	U	U	Denver, CO	89	57	18	7	1	6	7
Cincinnati, OH	75	41	22	3	3	6	9	Las Vegas, NV	300	172	92	25	7	4	11
Cleveland, OH	208	160	39	6	—	3	9	Ogden, UT	31	19	6	3	3	—	4
Columbus, OH	121	78	26	9	1	7	4	Phoenix, AZ	153	86	41	13	7	6	6
Dayton, OH	142	99	33	5	2	3	9	Pueblo, CO	20	11	5	3	1	—	—
Detroit, MI	U	U	U	U	U	U	U	Salt Lake City, UT	118	87	16	9	2	4	6
Evansville, IN	39	26	10	2	1	—	3	Tucson, AZ	164	107	37	12	6	2	5
Fort Wayne, IN	55	41	11	1	1	1	3	Pacific	1,583	1,080	345	90	39	26	119
Gary, IN	15	10	1	4	—	—	—	Berkeley, CA	14	11	1	1	—	1	—
Grand Rapids, MI	57	41	11	1	2	2	3	Fresno, CA	97	59	24	10	4	—	9
Indianapolis, IN	180	118	48	8	4	2	9	Glendale, CA	36	28	6	1	—	1	4
Lansing, MI	35	27	5	2	1	—	—	Honolulu, HI	59	45	10	—	3	1	7
Milwaukee, WI	92	50	29	10	1	2	1	Long Beach, CA	67	41	16	5	3	2	8
Peoria, IL	U	U	U	U	U	U	U	Los Angeles, CA	206	126	58	11	5	6	19
Rockford, IL	43	29	9	4	1	—	1	Pasadena, CA	21	15	5	1	—	—	3
South Bend, IN	55	44	8	2	—	1	1	Portland, OR	116	87	24	3	1	1	4
Toledo, OH	85	60	18	5	1	1	7	Sacramento, CA	240	163	51	15	7	4	13
Youngstown, OH	59	47	11	—	1	—	7	San Diego, CA	155	103	36	5	6	3	11
W.N. Central	448	301	84	36	17	9	15	San Francisco, CA	130	89	24	13	2	2	9
Des Moines, IA	—	—	—	—	—	—	—	San Jose, CA	187	128	44	11	4	—	20
Duluth, MN	35	28	4	2	1	—	2	Santa Cruz, CA	23	18	1	2	—	1	2
Kansas City, KS	23	16	3	4	—	—	—	Seattle, WA	78	54	17	4	2	1	2
Kansas City, MO	89	57	14	8	7	3	2	Spokane, WA	48	34	10	3	—	1	2
Lincoln, NE	35	28	4	2	—	1	2	Tacoma, WA	106	79	18	5	2	2	6
Minneapolis, MN	63	39	14	5	4	1	6	Total¶	9,880	6,484	2,289	651	234	217	519
Omaha, NE	U	U	U	U	U	U	U								
St. Louis, MO	60	38	13	4	4	1	3								
St. Paul, MN	49	37	7	5	—	—	—								
Wichita, KS	94	58	25	6	1	3	—								

U: Unavailable. —: No reported cases.

* Mortality data in this table are voluntarily reported from 122 cities in the United States, most of which have populations of >100,000. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included.

† Pneumonia and influenza.

§ Because of changes in reporting methods in this Pennsylvania city, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks.

¶ Total includes unknown ages.

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format. To receive an electronic copy each week, visit *MMWR*'s free subscription page at <http://www.cdc.gov/mmwr/mmwrsubscribe.html>. Paper copy subscriptions are available through the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Data are compiled in the National Center for Public Health Informatics, Division of Integrated Surveillance Systems and Services. Address all inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop E-90, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333 or to mmwrq@cdc.gov.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.