

## Influenza-Related Hospitalizations and Poverty Levels — United States, 2010–2012

James L. Hadler, MD<sup>1</sup>; Kimberly Yousey-Hindes, MPH<sup>1</sup>; Alejandro Pérez, MPH<sup>2</sup>; Evan J. Anderson, MD<sup>3,4</sup>; Marisa Bargsten, MPH<sup>5</sup>; Susan R. Bohm, MS<sup>6</sup>; Mary Hill, MPH<sup>7</sup>; Brenna Hogan, MPH<sup>8</sup>; Matt Laidler, MPH, MA<sup>9</sup>; Mary Lou Lindegren, MD<sup>10</sup>; Krista L. Lung, MPH<sup>11</sup>; Elizabeth Mermel, MS<sup>12</sup>; Lisa Miller, MD<sup>13</sup>; Craig Morin, MPH<sup>14</sup>; Erin Parker, MPH<sup>15</sup>; Shelley M. Zansky, PhD<sup>16</sup>; Sandra S. Chaves, MD<sup>2</sup>

Annual influenza vaccine is recommended for all persons aged  $\geq 6$  months in the United States, with recognition that some persons are at risk for more severe disease (1). However, there might be previously unrecognized demographic groups that also experience higher rates of serious influenza-related disease that could benefit from enhanced vaccination efforts. Socioeconomic status (SES) measures that are area-based can be used to define demographic groups when individual SES data are not available (2). Previous surveillance data analyses in limited geographic areas indicated that influenza-related hospitalization incidence was higher for persons residing in census tracts that included a higher percentage of persons living below the federal poverty level (3–5). To determine whether this association occurs elsewhere, influenza hospitalization data collected in 14 FluSurv-NET sites covering 27 million persons during the 2010–11 and 2011–12 influenza seasons were analyzed. The age-adjusted incidence of influenza-related hospitalizations per 100,000 person-years in high poverty ( $\geq 20\%$  of persons living below the federal poverty level) census tracts was 21.5 (95% confidence interval [CI]: 20.7–22.4), nearly twice the incidence in low poverty ( $< 5\%$  of persons living below the federal poverty level) census tracts (10.9, 95% CI: 10.3–11.4). This relationship was observed in each surveillance site, among children and adults, and across racial/ethnic groups. These findings suggest that persons living in poorer census tracts should be targeted for enhanced influenza vaccination outreach and clinicians serving these persons should be made aware of current recommendations for use of antiviral agents to treat influenza (6).

Influenza causes annual epidemics in the United States resulting in an estimated 4,900–27,000 deaths and 114,000–624,000 hospitalizations per year (7). Influenza vaccination recommendations have evolved from focusing on persons at

higher risk for severe disease and influenza-associated complications to a recommendation for vaccination of all persons aged  $\geq 6$  months (1). In addition to the recommendation for universal influenza vaccination, enhancing vaccination efforts in specific demographic groups that experience higher rates of serious influenza-related disease can reduce their vulnerability and minimize disparities.

Influenza surveillance data have generally not included individual measures of SES. Thus, any potential association

### INSIDE

- 106 HIV-Related Risk Behaviors Among Male High School Students Who Had Sexual Contact with Males — 17 Large Urban School Districts, United States, 2009–2013
- 110 Occupational HIV Transmission Among Male Adult Film Performers — Multiple States, 2014
- 115 Progress Toward Strengthening National Blood Transfusion Services — 14 Countries, 2011–2014
- 120 Interim Guidelines for Prevention of Sexual Transmission of Zika Virus — United States, 2016
- 122 Update: Interim Guidelines for Health Care Providers Caring for Pregnant Women and Women of Reproductive Age with Possible Zika Virus Exposure — United States, 2016
- 128 Notes from the Field: Circulating Vaccine-Derived Poliovirus Outbreaks — Five Countries, 2014–2015
- 130 Announcement
- 132 QuickStats

Continuing Education examination available at [http://www.cdc.gov/mmwr/cme/conted\\_info.html#weekly](http://www.cdc.gov/mmwr/cme/conted_info.html#weekly).



between influenza, particularly severe disease, and SES was unmeasured until area-based SES measures began to be used. During 1998–2005, the Public Health Disparities Geocoding Project recognized the potential for using area-based SES measures to describe and monitor the association between SES and reportable disease incidence. After comparing numerous possible area-based SES measures to describe SES disparities for selected health outcomes, the census tract poverty level of case residence was recommended to be used as a variable in addition to age, sex, and race/ethnicity in routine surveillance data analyses (2). During 2003–2005, the 10 Emerging Infections Programs (EIP)\* established active surveillance for influenza-related hospitalizations. Analyses of data from New Haven County, Connecticut, and eight counties in Tennessee indicated that, during multiple influenza seasons, including those when influenza A(H1N1)pdm09 predominated, influenza-related hospitalization incidence was consistently higher for children and adults residing in census tracts with higher percentages of persons living below the federal poverty level (3–5).

To assess the association between census tract-level poverty and influenza hospitalization at a national level, participating sites in the Influenza Hospitalization Surveillance Network (FluSurv-NET), including all 10 EIP sites, participated in a multisite analysis. FluSurv-NET is a national sentinel

surveillance system established in 2009 that conducts population-based surveillance for laboratory-confirmed influenza-associated hospitalizations annually during October–April. For this analysis, data were gathered from 78 counties in 14 FluSurv-NET states† representing approximately 9% of the U.S. population. Each site geocoded the residential address of all influenza-associated hospitalizations for the 2010–11 and 2011–12 influenza seasons. Geocoded addresses were assigned to census tracts. Census tract poverty level, defined as the percentage of households in the census tract living below the federal poverty level, was determined from the 2008–2012 American Community Survey 5-Year Estimates.§ Census tracts were categorized by their percentage of households living below the poverty level (<5%, 5%–9%, 10%–19%, ≥20%), and age-adjusted (2000 U.S. Standard Population) influenza-related hospitalization incidence overall and for each FluSurv-NET site was calculated, stratified by census tract poverty status. County and census tract-specific denominators were determined from the 2010 U.S. Census.

In total, 7,936 (96%) of 8,304 influenza-related hospitalizations were coded to census tract, including 5,624 in 2010–2011 and 2,312 in 2011–2012. For both seasons combined, the age-adjusted incidence of influenza-related hospitalizations per 100,000 person-years in high poverty (≥20% of persons

\*California, Colorado, Connecticut, Georgia, Maryland, Minnesota, New Mexico, New York, Oregon, and Tennessee. <http://www.cdc.gov/nceizd/dpei/eip/index.html>.

†Ten EIP states plus Michigan, Ohio, Rhode Island, and Utah.

§<http://www.census.gov/programs-surveys/acs>.

The *MMWR* series of publications is published by the Center for Surveillance, Epidemiology, and Laboratory Services, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30329-4027.

**Suggested citation:** [Author names; first three, then et al., if more than six.] [Report title]. *MMWR Morb Mortal Wkly Rep* 2016;65:[inclusive page numbers].

#### Centers for Disease Control and Prevention

Thomas R. Frieden, MD, MPH, *Director*  
 Harold W. Jaffe, MD, MA, *Associate Director for Science*  
 Joanne Cono, MD, ScM, *Director, Office of Science Quality*  
 Chesley L. Richards, MD, MPH, *Deputy Director for Public Health Scientific Services*  
 Michael F. Iademarco, MD, MPH, *Director, Center for Surveillance, Epidemiology, and Laboratory Services*

#### MMWR Editorial and Production Staff (Weekly)

Sonja A. Rasmussen, MD, MS, *Editor-in-Chief*  
 Charlotte K. Kent, PhD, MPH, *Executive Editor*  
 Jacqueline Gindler, MD, *Editor*  
 Teresa F. Rutledge, *Managing Editor*  
 Douglas W. Weatherwax, *Lead Technical Writer-Editor*  
 Soumya Dunworth, PhD, Teresa M. Hood, MS,  
*Technical Writer-Editors*

Martha F. Boyd, *Lead Visual Information Specialist*  
 Maureen A. Leahy, Julia C. Martinroe,  
 Stephen R. Spriggs, Moua Yang, Tong Yang,  
*Visual Information Specialists*  
 Quang M. Doan, MBA, Phyllis H. King,  
 Teresa C. Moreland, Terraye M. Starr,  
*Information Technology Specialists*

#### MMWR Editorial Board


Timothy F. Jones, MD, *Chairman*  
 Matthew L. Boulton, MD, MPH  
 Virginia A. Caine, MD  
 Katherine Lyon Daniel, PhD  
 Jonathan E. Fielding, MD, MPH, MBA  
 David W. Fleming, MD

William E. Halperin, MD, DrPH, MPH  
 King K. Holmes, MD, PhD  
 Robin Ikeda, MD, MPH  
 Rima F. Khabbaz, MD  
 Phyllis Meadows, PhD, MSN, RN  
 Jewel Mullen, MD, MPH, MPA

Jeff Niederdeppe, PhD  
 Patricia Quinlisk, MD, MPH  
 Patrick L. Remington, MD, MPH  
 Carlos Roig, MS, MA  
 William L. Roper, MD, MPH  
 William Schaffner, MD

living below the federal poverty level) neighborhoods was 21.5 (95% CI: 20.7–22.4), nearly twice the incidence in low poverty (<5% of persons living below the federal poverty level) neighborhoods (10.9, 95% CI: 10.3–11.4), with a gradient of increasing incidence as census tract poverty category increased (Figure 1). This relationship was observed in all 14 surveillance sites (Table), within groups defined by age (0–4 years, 5–17 years, 18–49 years, 50–64 years, and ≥65 years), within each racial/ethnic group (Figure 2) and during each influenza season. The relationship also was observed for age-adjusted rates for hospitalizations requiring intensive care, for those requiring mechanical ventilation, and for deaths during or within 30 days of hospitalization. The incidence rate ratios for census tracts with ≥20% versus <5% of households living below the federal poverty level were 1.96 [95% CI: 1.7–2.3] for hospitalizations requiring intensive care; 2.03 [95% CI: 1.6–2.5] for hospitalizations requiring mechanical ventilation; and 1.82 [95% CI: 1.3–2.7] for deaths occurring within 30 days of hospitalization. The overall percentage of hospitalized influenza patients who were vaccinated was inversely associated with census tract poverty level, from a high of 48% in the census tracts with the lowest poverty levels to a low of 35% in the census tracts with the highest poverty levels, a finding driven by differences in vaccination rates among persons aged ≥65 years, who accounted for 94% of hospitalized cases in the lowest poverty census tracts compared with 80% in the highest.

**FIGURE 1. Age-adjusted incidence of influenza-related hospitalizations per 100,000 person-years,\* by census tract poverty level — FluSurv-NET, 14 states, 2010–2012**



\* With 95% confidence intervals.

## Discussion

The association of higher census tract-level poverty with higher influenza-related hospitalization rates appears to be robust, occurring across counties in 14 states, within pediatric and adult age groups, and across racial/ethnic groups. Possible contributing factors are lower vaccination rates in residents of poorer census tracts, poverty-related crowding with higher rates of influenza transmission, and higher prevalence of medical conditions predisposing persons to influenza complications in poorer areas. However, differences in vaccination rates cannot fully explain all the age-specific differences by census tract poverty observed: only hospitalized influenza patients aged ≥65 years had a large enough difference in vaccination rates to fully explain the findings. Regardless of the causes, to reduce poverty-associated disparities in influenza-related hospitalizations, there is a need to increase influenza vaccination levels in higher poverty neighborhoods and to more fully implement recommendations on the use of antivirals in the outpatient setting (6). This will require enhanced efforts by public health agencies and health care providers to address missed opportunities for vaccination and system barriers (8), as well as a better understanding of personal barriers (9) to influenza vaccination in these neighborhoods. In addition, it will require evaluation of use of antivirals and efforts to improve them.

Healthy People 2020 includes a public health infrastructure objective (PHI 7.3) to increase the percentage of population-based health objectives for which national data are available by socioeconomic status (10). Public health surveillance data often lack information on the SES of individual persons, making it difficult to describe and monitor health disparities based on SES. When residential address data are available, it is possible to geocode the address, link it to census tract SES data and conduct analyses. The EIP, recognizing the potential to use geocoded addresses to examine possible SES-related disparities at a broad interstate level, formed a health equity workgroup to explore standardizing its use in all EIP states (11). This analysis demonstrates that multisite analyses using census tract SES can provide national-level data that are relevant to prevention efforts.

The findings in this report are subject to at least two limitations. First, a total of 4% of cases were unable to be geocoded and thus were not included in the analysis. Second, the data in the report were from two influenza seasons during which influenza A(H3N2) viruses predominated. The findings could be different during an A(H1N1) season or during an influenza pandemic. However, in the single site studies that stimulated this analysis (3–5), the association of higher census tract poverty with higher influenza-related hospitalization incidence was consistent throughout all seasons examined, regardless of the dominant circulating strain.


**TABLE. Age-adjusted incidence of influenza-related hospitalizations per 100,000 person-years, by census tract poverty level and state — FluSurv-NET, 14 states, 2010–2012**

State	Census tract poverty level*			
	0%–4%	5%–9%	10%–19%	≥20%
	Incidence (95% CI)	Incidence (95% CI)	Incidence (95% CI)	Incidence (95% CI)
California	11.5 (10.0–13.2)	14.5 (13.0–16.3)	15.9 (14.0–18.0)	21.4 (18.8–24.4)
Colorado	13.6 (12.8–17.0)	16.5 (14.1–19.1)	22.8 (20.1–25.8)	25.0 (21.9–28.3)
Connecticut	13.0 (11.4–14.8)	16.1 (13.7–18.7)	24.6 (21.1–28.4)	33.5 (29.0–38.6)
Georgia	9.1 (7.3–11.1)	7.9 (6.5–9.5)	10.3 (9.0–11.7)	12.9 (11.2–14.7)
Maryland	9.3 (8.0–10.7)	14.0 (12.1–16.1)	18.6 (16.1–21.3)	26.0 (22.7–29.6)
Michigan	8.5 (3.9–16.0)	11.9 (7.9–17.3)	14.9 (10.7–20.2)	20.5 (14.8–27.7)
Minnesota	7.3 (6.1–8.7)	9.7 (8.2–11.4)	15.1 (13.1–17.4)	24.1 (20.6–28.1)
New Mexico	7.5 (4.7–11.5)	10.1 (7.1–14.0)	14.6 (12.1–17.5)	15.9 (13.6–18.5)
New York	9.9 (8.1–11.9)	10.1 (8.5–11.9)	10.8 (9.0–12.8)	28.6 (24.9–32.7)
Ohio	7.8 (6.1–10.0)	10.0 (7.9–12.5)	11.2 (9.3–13.5)	21.5 (18.6–24.7)
Oregon	10.7 (7.9–14.2)	9.6 (7.7–11.9)	13.5 (11.4–15.8)	18.4 (15.2–22.0)
Tennessee	6.4 (4.5–8.7)	7.3 (5.5–9.5)	7.3 (5.6–9.5)	12.4 (9.9–15.3)
Utah	23.5 (19.0–28.7)	26.5 (22.2–31.3)	30.9 (26.4–36.1)	38.9 (32.8–45.7)
Rhode Island	14.1 (10.2–19.1)	14.6 (10.9–19.0)	14.4 (10.2–19.6)	26.0 (21.1–31.6)

Abbreviation: CI = confidence interval.

\* Percentage of population living below the federal poverty level.

**FIGURE 2. Age-adjusted incidence of influenza-related hospitalizations per 100,000 person-years,\* by racial/ethnic group and census tract poverty level,† — FluSurv-NET, 14 states, 2010–2012**



\* With 95% confidence intervals.

† Percentage of population living below the federal poverty level.

Using census tract-based SES measures as variables for surveillance data analysis can contribute to achieving the Healthy People 2020 public health infrastructure goal of having national population-based data available by SES. It is important from

an influenza control perspective that local vaccination efforts be emphasized in demographic groups found to have a higher incidence of more severe and costly complications of influenza, including hospitalization, intensive care and mechanical

## References

## Summary

## What is already known on this topic?

Measures of socioeconomic status are infrequently used in public health surveillance. Several studies in small U.S. geographic areas found that higher census tract-level poverty is associated with higher population-level rates of influenza-related hospitalization, a finding with possible implications for influenza control efforts.

## What is added by this report?

A collaborative initiative among 14 states that examined the association between census tract-level poverty and incidence of influenza-related hospitalization found increasing rates of influenza-related hospitalization with increasing census tract poverty. This finding was present during two influenza seasons, among all 14 sites, all age and racial/ethnic groups, and for more severe outcomes of hospitalization (intensive care, respiratory support, and death).

## What are the implications for public health practice?

Persons who live in high poverty census tracts represent a demographic group at higher risk for severe influenza outcomes. Persons in poorer neighborhoods should be a focus for enhanced influenza vaccination outreach and early use of antiviral treatment. Analysis of surveillance data using census tract-level measures of socioeconomic status can provide new perspectives and directions for prevention of diseases of public health importance.

ventilation. Based on the consistency of the findings in this study across FluSurv-NET sites, persons who live in high poverty census tracts are one such demographic group. Enhanced influenza outreach to improve influenza vaccination coverage for persons living in poorer neighborhoods and efforts to increase use of antivirals by clinicians serving these neighborhoods could reduce poverty-related disparities in severe influenza outcomes.

1. Grohskopf LA, Sokolow LZ, Olsen SJ, Bresee JS, Broder KR, Karron RA. Prevention and control of influenza with vaccines: recommendations of the Advisory Committee on Immunization Practices, United States, 2015–16 influenza season. *MMWR Morb Mortal Wkly Rep* 2015;64:818–25. <http://dx.doi.org/10.15585/mmwr.mm6430a3>.
2. Krieger N, Chen JT, Waterman PD, Rehkopf DH, Subramanian SV. Painting a truer picture of US socioeconomic and racial/ethnic health inequalities: the Public Health Disparities Geocoding Project. *Am J Public Health* 2005;95:312–23. <http://dx.doi.org/10.2105/AJPH.2003.032482>.
3. Yousey-Hindes KM, Hadler JL. Neighborhood socioeconomic status and influenza hospitalizations among children: New Haven County, Connecticut, 2003–2010. *Am J Public Health* 2011;101:1785–9. <http://dx.doi.org/10.2105/AJPH.2011.300224>.
4. Tam K, Yousey-Hindes K, Hadler JL. Influenza-related hospitalization of adults associated with low census tract socioeconomic status and female sex in New Haven County, Connecticut, 2007–2011. *Influenza Other Respir Viruses* 2014;8:274–81. <http://dx.doi.org/10.1111/irv.12231>.
5. Sloan C, Chandrasekhar R, Mitchel E, Schaffner W, Lindegren ML. Socioeconomic disparities and influenza hospitalizations, Tennessee, USA. *Emerg Infect Dis* 2015;21:1602–10. <http://dx.doi.org/10.3201/eid2109.141861>.
6. Fiore AE, Fry A, Shay D, Gubareva L, Bresee JS, Uyeki TM. Antiviral agents for the treatment and chemoprophylaxis of influenza—recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep* 2011;60(No. RR-01).
7. Reed C, Chaves SS, Daily KP, et al. Estimating influenza disease burden from population-based surveillance data in the United States. *PLoS One* 2015;10:e0118369.
8. Bryant WK, Ompad DC, Sisco S, et al.; Project VIVA Intervention Working Group. Determinants of influenza vaccination in hard-to-reach urban populations. *Prev Med* 2006;43:60–70. <http://dx.doi.org/10.1016/j.ypmed.2006.03.018>.
9. Chen JY, Fox SA, Cantrell CH, Stockdale SE, Kagawa-Singer M. Health disparities and prevention: racial/ethnic barriers to flu vaccinations. *J Community Health* 2007;32:5–20. <http://dx.doi.org/10.1007/s10900-006-9031-7>.
10. US Department of Health and Human Services. Healthy People 2020 topics and objectives: public health infrastructure. <http://www.healthypeople.gov/2020/topics-objectives/topic/public-health-infrastructure>.
11. Hadler JL, Vugia DJ, Bennett NM, Moore MR. Emerging Infections Program efforts to address health equity. *Emerg Infect Dis* 2015;21:1589–94. <http://dx.doi.org/10.3201/eid2109.150275>.

<sup>1</sup>Connecticut Emerging Infections Program, Yale School of Public Health, New Haven, Connecticut; <sup>2</sup>Influenza Division, National Center for Immunization and Respiratory Diseases, CDC; <sup>3</sup>Departments of Pediatrics and Medicine, Emory University School of Medicine, Atlanta, Georgia; <sup>4</sup>Atlanta Veterans Affairs Medical Center, Atlanta, Georgia; <sup>5</sup>New Mexico Department of Health; <sup>6</sup>Division of Communicable Disease, Michigan Department of Health and Human Services; <sup>7</sup>Salt Lake County Health Department, Salt Lake City, Utah; <sup>8</sup>Maryland Department of Health and Mental Hygiene; <sup>9</sup>Oregon Public Health Division, Oregon Health Authority; <sup>10</sup>Department of Pediatrics, Department of Health Policy, Vanderbilt University School of Medicine, Nashville, Tennessee; <sup>11</sup>Ohio Department of Health; <sup>12</sup>Rhode Island Department of Health; <sup>13</sup>Colorado Department of Public Health and Environment; <sup>14</sup>Virginia Department of Health; <sup>15</sup>California Emerging Infections Program; <sup>16</sup>New York Emerging Infections Program, New York State Department of Health.

Corresponding author: James L. Hadler, [hadler-epi@att.net](mailto:hadler-epi@att.net), 203-507-0911.